

List of Figures

Figure 1.1: Atomic arrangement of self-organized gold chains on Ge (001). (a) Topography from STM, showing parallel alignment of Au-induced chains, spaced by 1.6 nm (bias -1.0 V, 0.4 nA). The elevated chain architecture exposes a single-atom ridge, including a small buckling, which hosts the 1D electron liquid under study. (b) Schematic representation of chain formation and atomic modification possibilities. Epitaxy of Au leads to self-organized chain formation at elevated temperatures.³⁷19

Figure 1.2: Snapshots from simulations of breaking gold (upper row) and copper contacts (lower row). The gold contact exhibits the formation of a single-atomic chain, while the copper contact breaks after a single-atom contact is formed. Out of the six metals studied (Cu, Ag, Au, Ni, Pd, and Pt) only Au and Pt form chains.³²20

Figure 1.3: Snapshots of the molecular dynamics simulations showing the formation of mixed LACs from alloys: a) Au₄₀Ag₆₀; and b) Au₆₀Ag₄₀. Yellow (grey) indicates gold (silver) atoms.⁵⁰26

Figure 3.1: Atomic arrangements and unit cell for LAC of Au-Ag and Au-Pt with four different compositions labelled as A, B,C and D. The horizontal line with the arrow shows unit cell considered for the calculation for each composition, Yellow sphere shows Au atoms, grey spheres shows Pd (Ag or Pt) atoms in LAC.51

Figure 3.2: Electronic band structure and Density of states for Au-Pd LAC for different compositions described in Fig 3.158

Figure 3.3: Electronic band structure and Density of states for Au-Ag LAC for different compositions described in Fig 3.159

Figure 3.4: Electronic band structure and Density of states for Au-Pt LAC for different compositions described in Fig.3.1, are exhibited.60

Figure 3.5: Computed real part $\epsilon_1(\omega)$ and imaginary part of dielectric function, $\epsilon_2(\omega)$, reflectance spectra, $R(\omega)$ absorption spectra, $\alpha(\omega)$ and EELS spectra $L(\omega)$ of Au-Pd bimetallic LAC with different compositions A,B,C and D defined in Figure 3.1, are displayed as function of energy.....62

Figure 3.6: Computed real part $\epsilon_1(\omega)$ and imaginary part of dielectric function, $\epsilon_2(\omega)$, reflectance spectra, $R(\omega)$ absorption spectra, $\alpha(\omega)$ and EELS spectra $L(\omega)$ of Au-

Ag bimetallic LAC with different compositions A,B,C and D defined in Figure 3.1, are displayed as function of energy.....63

Figure 3.7: Computed real part $\epsilon_1(\omega)$ and imaginary part of dielectric function, $\epsilon_2(\omega)$, reflectance spectra, $R(\omega)$ absorption spectra, $\alpha(\omega)$ and EELS spectra $L(\omega)$ of Au-Pt bimetallic LAC with different compositions A,B,C and D defined in Figure 3.1, are displayed as function of energy.....64

Figure 4.1: Variation of cohesive energy per atom, E_{coh} and Tension with average z-projected interatomic distance, dz for 4-types of Au-Pd, Au-Ag and Au-Pt chain structures.....69

Figure 4.2: Optimized 2-types of chain structures of Au-Pd with bond lengths and bond angles. Yellow and Brown spheres present Au and Pd atoms, respectively.....70

Figure 4.3: Optimized 4-types of chain structures of Au-Ag with bond lengths and bond angles. Yellow and Grey spheres present Au and Pt atoms, respectively.71

Figure 4.4: Optimized 4-types of chain structures of Au-Pt with bond lengths and bond angles. Yellow and Green spheres present Au and Ag atoms, respectively.72

Figure 4.5: Electronic band structure and density of states for the type-C Au-Pd, type-A Au-Ag and the type-D Au-Pt atomic chains depicted in Figure 275

Figure 4.6: Real and imaginary parts of the dielectric function, reflectivity and electron energy loss spectrum for the type-C Au-Pd, type-A Au-Ag and type-D Au-Pt atomic chains76

Figure 5.1: a) and b) Au-Pd, Au-Pt and Au-Ag chains over NiAl(110) substrate ; c) and d) Au-Pd, Au-Pt and Au-Ag chains over Cu(110) substrate. The copper atoms are represented by reddish yellow spheres. Al atoms are shown by light blue spheres while light grey spheres display Ni atoms in NiAl substrate. The chains atoms are shown by blue(Pd(Pt, Ag)) and yellow (Au) spheres, respectively, when are placed chain on substrates.80

Figure 5.2: (a) Enlarged forms of chains without substrate are shown. Grey atoms represent Pd (Pt, Ag), while Au atoms are shown by yellow spheres. (b) Two different orientation of double chains placed on Cu (110) substrate. of Bimetallic chains placed on Cu(110) substrate. Relaxations of chain and substrate system yields three different possible configurations termed as A, B & C.82

Figure 5.3: Projected density of states (PDOS) are plotted as function of energy for chain-NiAl(110), chain-Cu(110) systems. The PDOS for free standing chains of Au-Pd, Au-Pt and Au-Ag are also displayed in the figure.89

Figure 5.4: PDOS for different configuration of Au-Pd double zigzag chains on Cu (110) substrate (left panel) and corresponding free-standing chains of Au-Pd(right panel). The structures are exhibited in Fig.5.2.....91

Figure 5.5: PDOS for different configurations of Au-Pt chains on Cu (110) substrate (left panel) and for corresponding free-standing chains of Au-Pt(right panel), whose geometries are as shown in Fig. 5.2.....92

Figure 5.6: PDOS for different configuration of Au-Ag chains on Cu (110) substrate (left panel) and for corresponding free-standing chains of Au-Ag(right panel). The geometries of structures are shown in Fig. 5.2.93