

LIST OF TABLES

- Table 1.** Few Asclepiadaceae family members along with their metabolites and properties
- Table 2.** Murashige and Skoog's (MS, 1962) basal stock composition
- Table 3.** Preparation of PGR stock solutions
- Table 4.** Effect of different antibiotics on inhibition of bacterial growth
- Table 5.** Individual and synergistic effect of different PGRs on callus growth and morphology in leaf explants of *L. reticulata* (6 weeks).
- Table 6.** Effect of different PGRs on induction of somatic embryos using leaf explants of *L. reticulata* (6 weeks).
- Table 7.** Effect of PGRs on shoot regeneration from nodal explants of *L. reticulata* (8 weeks)
- Table 8.** Peak areas of different samples at 254 nm for hexane extracts
- Table 9.** Peak areas of different samples at 366 nm for hexane extracts
- Table 10.** Peak areas of different samples at 525 nm for hexane extracts
- Table 11.** Peak areas of different samples at 254 nm for ethyl acetate extracts
- Table 12.** Peak areas of different samples at 366 nm for ethyl acetate extracts
- Table 13.** Peak areas of different samples at 525 nm for ethyl acetate extracts
- Table 14.** Peak areas of different samples at 254 nm for methanol extracts
- Table 15.** Peak areas of different samples at 366 nm for methanol extracts
- Table 16.** Peak areas of different samples at 525 nm for methanol extracts
- Table 17.** Intra-day and inter-day precisions for method validation
- Table 18.** HPTLC method validation parameters for *p*-coumaric acid
- Table 19.** Quantification of *p*-Coumaric acid in *in vivo* and *in vitro* samples of *L. reticulata* using HPTLC
- Table 20.** Effect of different concentrations of YE on *p*-coumaric acid content
- Table 21.** Effect of different PGRs on shoot regeneration from leaf explants of *T. indica* (8 weeks)
- Table 22.** Effect of different PGRs on somatic embryogenesis from leaf explants of *T. indica* (8 weeks)
- Table 23.** Effect of cytokinins on shoot regeneration from nodal explants of *T. indica* (8 weeks)
- Table 24.** Effect of IBA and NAA on rooting in *T. indica* shoots (4 weeks)
- Table 25.** Peak areas of different samples at 254 nm for hexane extracts

Table 26. Peak areas of different samples at 366 nm for hexane extracts

Table 27. Peak areas of different samples at 525 nm for hexane extracts

Table 28. Peak areas of different samples at 254 nm for ethyl acetate extracts

Table 29. Peak areas of different samples at 366 nm for ethyl acetate extracts

Table 30. Peak areas of different samples at 525 nm for ethyl acetate extracts

Table 31. Peak areas of different samples at 254 nm for methanol extracts

Table 32. Peak areas of different samples at 366 nm for methanol extracts

Table 33. Peak areas of different samples at 525 nm for methanol extracts

Table 34. Quantification of lupeol in *in vivo* and *in vitro* samples of *T. indica* using HPTLC

Table 35. Quantification of lupeol in presence of YE at different time intervals

Table 36. Quantification of lupeol in presence of SA at different time intervals