

## LIST OF FIGURES

- Figure 1.** Medicinal plant species availability in different zones of Gujarat
- Figure 2.** Different aspects of plant tissue culture
- Figure 3.** *Leptadenia reticulata* (Retz.) Wight and Arn plant growing in the Botanical Garden of the University
- Figure 4.** *Tylophora indica* (Burm. F.) Merrill plant growing in the Botanical Garden of the University
- Figure 5.** *p*-Coumaric biosynthesis via phenylpropanoid pathway
- Figure 6.** Lupeol biosynthesis via MVA/MEP pathways
- Figure 7.** Antibiotic assay using disk diffusion method
- Figure 8.** Effect of individual cytokinins on callus differentiation from leaf explants of *L. reticulata*
- Figure 9.** Effect of individual auxins on callus formation from leaf explants of *L. reticulata* at the end of six weeks
- Figure 10.** Synergistic effect of PGRs on callus growth and morphology from leaf explants of *L. reticulata* at the end of six weeks
- Figure 11.** Induction of somatic embryos (SEs) from leaf explants of *L. reticulata* in presence of BA (15  $\mu$ M) + Kn (5  $\mu$ M) at the end of six weeks
- Figure 12.** Differentiation of somatic embryos (SEs) in an optimized medium with BA (15  $\mu$ M) + AdSO<sub>4</sub> (15  $\mu$ M) from leaf explants of *L. reticulata*
- Figure 13.** Microscopic observation of somatic embryos (SEs) differentiated on optimized medium BA (15  $\mu$ M) + AdSO<sub>4</sub> (15  $\mu$ M)
- Figure 14.** Differentiation of somatic embryos (SEs) in the presence of BA (10  $\mu$ M) + IAA (0.5  $\mu$ M) from leaf explants of *L. reticulata*
- Figure 15.** Development of SEs in GA<sub>3</sub> (1.0  $\mu$ M) derived through leaf explants of *L. reticulata*
- Figure 16.** Effect of individual cytokinins on shoot regeneration from nodal explants of *L. reticulata* at the end of eight weeks
- Figure 17.** Effect of synergistic cytokinins on shoot regeneration from nodal explants of *L. reticulata* at the end of eight weeks
- Figure 18.** Axillary shoot regeneration in optimized medium fortified with BA (10  $\mu$ M) + AdSO<sub>4</sub> (10  $\mu$ M) from nodal explant of *L. reticulata*

**Figure 19.** Effect of CW on axillary shoot elongation in *L. reticulata* by end of four weeks.

**Figure 20.** Elongation of axillary shoots in presence of CW at the end of four weeks

**Figure 21.** Rooting of *L. reticulata* micro shoots

**Figure 22.** Rooting of *L. reticulata* shoots in different MS medium strengths fortified with IBA

**Figure 23.** HPTLC fingerprints of hexane extract for different samples of *L. reticulata*

**Figure 24.** Chromatograms of different samples for hexane extract in *L. reticulata*

**Figure 25.** HPTLC fingerprints of ethyl acetate extract for different samples of *L. reticulata*

**Figure 26.** Chromatograms of different samples for ethyl acetate extract in *L. reticulata*

**Figure 27.** HPTLC fingerprints of methanol extract for different samples of *L. reticulata*

**Figure 28.** Chromatograms of different samples for methanol extract in *L. reticulata*

**Figure 29.** Standard curve of *p*-coumaric acid

**Figure 30.** Quantification of *p*-coumaric acid using HPTLC in samples of *L. reticulata*.

**Figure 31.** Growth determination of callus till six weeks

**Figure 32.** Effect of yeast extract on callus biomass derived through leaf explants of *L. reticulata*

**Figure 33.** TLC plates and chromatograms for methanol extract of callus samples

**Figure 34.** Effect of individual cytokinins on leaf explants of *T. indica*

**Figure 35.** Effect of individual auxins on leaf explants of *T. indica* at the end of eight weeks

**Figure 36.** Synergistic effect of PGRs on leaf explants of *T. indica* at the end of eight weeks

**Figure 37.** Synergistic effect of BA (10  $\mu$ M) and Kn (15  $\mu$ M) on optimum shoot regeneration from leaf explants of *T. indica*

**Figure 38.** Synergistic effect of PGRs on indirect shoot regeneration from leaf explants of *T. indica* at the end of eight weeks

**Figure 39.** Differentiation of somatic embryos in different combinations of PGRs from leaf explants of *T. indica* at the end of eight weeks

**Figure 40.** Somatic embryos from leaf explant of *T. indica* in MS + Kn (5  $\mu$ M) + IAA (0.1  $\mu$ M)

**Figure 41.** Microscopic observation of SEs formed in MS + Kn (5  $\mu$ M) + IAA (0.1  $\mu$ M) from leaf explant of *T. indica*

**Figure 42.** Development and elongation of plantlets in static medium fortified with different PGR combinations at the end of four weeks in *T. indica*

**Figure 43.** Development and elongation of plantlets in liquid medium fortified with different PGRs combinations at the end of four weeks in *T. indica*

**Figure 44.** Development and elongation of plantlets in MS medium fortified with GA<sub>3</sub> (1 μM) in *T. indica*

**Figure 45.** Development of plantlets in different media after four weeks

**Figure 46.** Effect of individual cytokinins on shoot regeneration from nodal explants of *T. indica* at the end of eight weeks

**Figure 47.** Synergistic effect of PGRs on optimum shoot regeneration in MS + BA (10 μM) + Kn (15 μM) from nodal explants of *T. indica*

**Figure 48.** Effect of synergistic cytokinins on shoot regeneration from nodal explants of *T. indica* at the end of eight weeks

**Figure 49.** Indirect shoot regeneration from nodal derived callus in presence of BA (20 μM) + AdSO<sub>4</sub> (5 μM) in *T. indica*

**Figure 50.** Response of nodal derived callus for shoot regeneration in different of BA + AdSO<sub>4</sub> by the end of eight weeks

**Figure 51.** Induction of indirect SEs from nodal derived callus in presence of Kn (10 μM) + AdSO<sub>4</sub> (15 μM) in *T. indica*

**Figure 52.** Response of nodal derived callus for differentiation of SEs in different concentrations of Kn + AdSO<sub>4</sub> by the end of eight weeks

**Figure 53.** Histology of SEs in *T. indica*

**Figure 54.** Rooting of *T. indica* shoots at the end of four weeks

**Figure 55.** Differentiation of SEs from immersed stem portion in ½MS medium added with NAA at the end of four weeks in *T. indica*

**Figure 56.** Formation of SEs from stem of shoots in ½MS medium supplemented with NAA

**Figure 57.** Hardening of *T. indica* shoots developed from different regenerative pathways

**Figure 58.** Growth of *T. indica* plantlets during hardening and acclimatization

**Figure 59.** Schematic representation of different regenerative pathways in *T. indica*

**Figure 60.** HPTLC fingerprints of hexane extract for different samples of *T. indica*

**Figure 61.** Chromatograms of samples for hexane extract in *T. indica*

**Figure 62.** HPTLC fingerprints of ethyl acetate extract for different samples of *T. indica*

**Figure 63.** Chromatograms of samples for ethyl acetate extract in *T. indica*

**Figure 64.** HPTLC fingerprints of methanol extract for different samples of *T. indica*

**Figure 65.** Chromatograms of samples for methanol extract in *T. indica*

**Figure 66.** Quantification of lupeol in different samples of *T. indica*

**Figure 67.** TLC plate and respective chromatogram for hexane extract of control shoot samples

**Figure 68.** Effect of yeast extract on shoot biomass derived through nodal explants of *T. indica*

**Figure 69.** TLC plates and respective chromatograms for hexane extract of shoot samples after YE treatment

**Figure 70.** Effect of salicylic acid on shoot biomass derived through nodal explants of *T. indica*

**Figure 71.** TLC plates and respective chromatograms for hexane extract of shoot samples after SA treatment