

LIST OF PLATES

No	Title	Page
Plate 2 1 (A)	Field photograph showing stabilized parabolic dunes with palaeosols Loc Luni river bank, Tilwara	32
(B)	A close up view of the above section containing char coal and bone remains	32
Plate 2 2	A view of Luni river cliff showing early Pleistocene fluvio - lacustral sequences Loc Sindari.	36
Plate 3 1 (A)	A view of wind faceted granitic yardang Loc north of Pali	51
(B)	A view of rhyolitic ridge showing intensely developed joints due to arid climate and wind deflation Loc. Dhorimanna	51
Plate 3 2 (A)	A view of sandstone plateau with ferricretic duricrust and the pedimentary slopes merging with aeolian plains Loc Khara rann, Chetral	57
(B)	Panoramic view of Luni river showing deep entrenchment Loc Bhukan.	57
Plate 3 3 (A)	A view of falling dunes (obstacle dunes). Loc. Chotan.	58
(B)	A view of shrub coppice dunes. Loc Bhaniana	58
Plate 3 4	Panoramic view of mega parabolic dune developed in the vicinity of sandstone mesa Loc. Belwa.	59
Plate 3 5 (A)	A view of transverse dunes. Loc Sujangarh.	60
(B)	A view of dunefield showing coalesced barchans Loc. Phalsund	60
Plate 4 1 (A)	A view of stabilized dunal profile with palaeosol horizons. Loc. Dhanola trijunction	77
(B)	A close view of above profile showing iron stained second palaeosol horizon lying at the base	77

Plate 4 2 (A)	A view of the nodular calcretic profile developed near Raneri.	79
(B)	A view of the calcretic profile with nodular form at the top and massive, powdery form at the base Loc Sardarsahar	79
Plate 4 3 (A)	A view of the calcrete profile with well developed nodular calcrete. Loc. Surpalia	81
(B)	A view of calcretic profile showing a cap of active dunal sand followed by palaeosol (greyish) horizon and massive calcrete. Loc Bhaleri	81
Plate 4 4 (A)	A view of calcretic profile showing coalesced nodular calcrete at the top and hardpan calcrete at the bottom. Loc. Belwa.	84
(B)	A view of quarry section showing development of nodular calcrete over weathered pediments of Jodhpur sandstone. Loc. Nimbijodhan	84
Plate 4 5 (A)	A view of dunal profile showing nodular calcretes, lime coated rhyolitic colluvium Loc. Tejsingh ki dhani	88
(B)	A view of quarry section showing the development of honeycomb calcrete in a sandy plain. Loc Girab.	88
Plate 4 6 (A)	A view of calcrete profile in a sandy plain. Loc. Vav.	93
(B)	A view of intensely calcretized gravelly sand Loc. Chetak, Bikaner.	93
Plate 4 7 (A)	A view of calcretic profile within the interdunal flat comprising nodular calcrete at top and honeycomb calcrete at bottom Loc Kuri	96
(B)	A view of hard honeycomb calcrete developed in an interdunal area. Loc Sultana.	96
Plate 4 8 (A)	A view of quarry section showing the development of nodular calcrete and honeycomb calcrete over Shumar formation. Loc. Sadewala	97
(B)	A view of nodular calcrete incorporating ferricrete breccias Loc. Mohangarh	97

Plate 4 9	Photomicrograph of calcretes showing	101
(A)	Type I Pedogenic carbonate and vesicular structure	
(B)	Type I-A Pedogenic carbonate with neocalcitans around detrital	
(C)	Type II Diffused nodules with a clear s - fabric	
(D)	Type II-A Pedogenic carbonate with a diffusive boundary	
(E)	Type III-A Pedogenic nodules without any impregnation	
(F)	Type III-B Pedogenic nodules with uniform iron impregnation.	
Plate 4 10	Photo micrographs depicting	103
(A)	Type III-C Nodules with manganans.	
(B)	Type IV Glaebules with dissolution feature along the boundary.	
(C)	Type VI Pedogenic carbonate - channel neocalcitan indicative of pedogenic carbonate accretion	
(D)	Meniscus cement around detrital grain indicative of high soil moisture regime.	
(E)	Poorly developed needle fibers, a feature of quick precipitation with partial transformation into sparites	
(F)	Typical floating texture, characteristic feature of groundwater precipitations.	
Plate 4 11	Photomicrographs of calcretes illustrating	110
(A)	In situ weathered biotite with a ferruginous halo - a feature of pedogenesis and neoformations	
(B)	An orthic nodule with glaebular halo, pointing to in situ development of nodules.	
(C)	An allothitic nodule, an indicator of palaeo erosional surfaces.	
(D)	A disorthic nodule - a feature pointing to pedoturbated soils	
(E)	Translocated skeletal grains along the channel, supporting the pedoturbations.	
(F)	Intergrain micro aggregate structure, caused due to segregation of b - fabric	
Plate 4 12	Photomicrographs of calcretes depicting	136
(A)	In situ weathering of mica and incorporation of neoformations into the b- fabric A common feature in the early stages of pedogenesis	
(B)	Altered, dissolved feldspar detrital with retention of twin lamellae - a feature of in situ weathering.	
(C)	Etched, corroded and dissolved quartz detrital in a micritic groundmass indicating high alkaline soil- microenvironment.	

	(D) Argillic cutans, an ideal feature of clay illuviation	
	(E) Bow shaped features (Bullock et al , 1985) suggestive of intense pedoturbation	
	(F) Insitu fractured quartz detrital caused due to high soil temperature gradient	
Plate 5 1	(A) A view of the massive ferricrete surface developed over Tertiary sandstones Loc. RD 105 Indira Gandhi Canal, near Sultana	157
	(B) A dismantled calcretized ferricrete profile Loc Ramgarh	157
Plate 5 2	(A) Calcretized - ferricrete capping over isolated plateaucic surfaces. Loc Nr. Bandah	159
	(B) A close view of above duricrust depicting fragments of calcretized ferricretes	159
Plate 5 3	(A) Calcretized ferricretes as capping over Plateaus Loc Khenya	162
	(B) Sheet type ferricretes developed over Lathi sandstone (Jurassic) Loc Savanta	162
Plate 5 4	(A) A view of ferricrete profile developed over sandstone Loc. Kaladongar	163
	(B) A close view of above ferricrete surface showing dissolution features	163
Plate 5 5	(A) Multi layered ferricrete sheets forming a cap over Tertiary sandstone Loc. Chetral	165
	(B) A view of ferricrete pedimentary surface Loc Near Khuyala	165
Plate 5 6	Photomicrographs of ferricrete profiles illustrating	168
	(A) Vesicular structure suggestive of pedogenic activity.	
	(B) Desilicification features in ferricretes, indicative of lateritic weathering	
	(C) Kaolinised, corroded, dissolved feldspar in mottled horizon a characteristic feature of lateritic weathering.	
	(D) Relics of original rock fabric along with small iron segregations in a clayey pallid horizon.	
	(E) Dissolution of pisoliths giving rise to concentric opaque and clayey rings caused due to dismantling of ferricrete profile	
	(F) Vesicles filled by sparites . an indication of I st phase of calcretization	

Plate 5 7	Photo micrographs of ferricrete profiles elucidating	169
	(A) Sparitic fillings in the cavities of ferricretes denoting Ist phase of calcretization	
	(B) Sparites forming pendent cements around detritals	
	(C) Sparites filling the dissolution spaces around the corroded quartz skeletal grains	
	(D) Channel filled with sparites and development of micro laminations along channel wall due to calcite replacement.	
	(E) Accumulation of cryptocrystalline silica along the fractures in mottled horizon, an indicator of microenvironmental changes within ferricrete profiles	
	(F) Micrites encompassing the ferricrete fragments, illustrative of II phase of calcretization	
Plate 5 8	Photomicrographs of calcretized ferricretes depicting	170
	(A) Neoformed sparites around ferricrete fragments developed from the II generation micrites - depicting maturity of calcretes.	
	(B) Neoformed sparites around ferricrete fragments at an intial stage of calcite replacement.	
	(C) Intense replacive action of calcites on ferricretes depicting the progression of calcretization	
Plate 7 1	Panoramic view of border roads in an active dunal tract. Loc Raniya.	210
Plate 7 2 (A)	A view of road alignment. Observe the failures (pot holes and headward gullies) developed due to poorly stabilized embankments Loc Nachana	211
	(B) A close view of headward gully depicting poor compaction and less thickness of pavement courses Loc Nachana.	211