

REFERENCES

- Agrawal, S.K. (1956) On the so-called "macrocephalus" beds of Kutch, Curr. Sci., V-25., pp 84.
- Agrawal, S.K.(1958). Kutch Mesozoic: A study of the Jurassics of Kutch with Special reference to the Jhura Dome, Jour. Paleont. Soc.Ind, Lucknow,V-2, pp 119-
- Alcock.A (1895) Materials for a Carinological fauna of India, No.1, The Brachyura Oxyrhyucha. Jour. As. Sos. Bengal. 1xiv, pt. 11, pp. 157-289.
- Alcock.A (1895a) Description of New species of Oxyrhyucha crab of Genus Parthenope, Jour. As. Sos.Bengal. 1xiv, pt. 11 no. 11, pp-134-396.
- Alcock.A (1896) Materials for a Carinological fauna of India, No. 2, The Brachyura Oxyrhyucha. Jour. As. Sos. Bengal. 1xiv, pt. 11, No II pp. 134-296.
- Alcock. A (1898) Materials for a Carinological fauna of India, No. 3, The Brachyura Cyclometopa. Part 1, The Family Xanthidae. Jour. As. Sos. Bengal. 1xiv, pt. 11, No II, pp. 67-233.
- Alcock. A (1899) Materials for a Carinological fauna of India, No. 4, The Brachyura Cyclometopa. Part II. A revision of the Cyclometopa with account of the families Portunidae, Cancridae and Corystidae. Jour. As. Sos. Bengal. 1xviii. pt.11 , no I, pp.1-104.
- Alcock. A (1899a) Natural History notes from R.I.M. Survey ship 'investigator' Ser.III. No. 3. On some new and rare species of crustacea. Jour. As. Sos. Bengal. 1xviii, pt.11,no II, pp.111- 119.
- Alcock. A (1899b) Materials for a Carinological fauna of India, No. 5, The Brachyura Primigenia, or Dormicea Jour. As. Sos. Bengal. 1xviii,pt.11, no.III, 123-169.
- Alcock. A (1900) Materials for a Carinological fauna of India, No.6, The Brachyura Catometopa, or Graspsoidea. Jour. As. Sos. Bengal. 1xix, pt.11, no III, pp.279-459.
- Alexander, J.; Bridge, J.S.; Cheel, R.J.; Leclair, S.F.; (2001) Bedform and associated sedimentary structures formed under supercritical water flow over aggrading sand beds. Sedimentolgy, V-48, 133-152.
- Allen. J.R.L., (1968) Current ripples. Amsterdam, Northen Holland.
- Allen, J.R.L (1982) Sedimentary Structures: Their Character and Physical Basis, Vol.1 Elsevier Amsterdam.
- Allen, J.R.L and Friend, P.F. (1976) Changes in intertidal dunes during two spring- neap cycles, Lifeboat Station Bank, Wells-next-the Sea, Norflok (England). Sediment. V-23, pp. 329-346.
- Allen, J.R.L., Friend, P.F., Lloyd, A., Wells, H. (1994) Morphodynamics of intertidal dunes: a year long study at lifeboat station bank, Wells-next-the Sea, Eastern England. Phil. Trans. R.Soc. Lond. V-347, pp. 291-345.

- Alpert, S.P. (1974) Systematic review of the genus *Skolithos*; Jour of Paleontology, V-49; 919-924.
- Ambraseys, N.N. (1988): Engenerring Seismology:Earthquake engeneering and structural dynamics- Journal of Inter. Assoc. Earthq. Engg.V-17: 1-105.
- Anonymous (1954) Instructor for collectors (Invertebrate animal other than insect) British Museum Natural History, London.
- Badve, R.M and Bhosle, S.P. (1987) Burrowing of the Gastropod *Umbonium*, link and its application in ichnofossil study. Current Science, V-56; pp 1164-1166.
- Badve, R.M and Ghare, M.A. (1978) Jurassic Ichnofauna of Kutch, Pt-I., Biovigyanum, V-4, pp-125- 140.
- Barfield, A.E. and Newell, G.E. (1961) The behaviour of *Macoma baltica* (L). Jour. Mar. Biol. Assoc. U.K.; V-41; 81-87.
- Barrington, E.J.W. (1965) The biology of Hemichordata and Protochordata, 1-176, Edinburgh.
- Barnes, R.D. (1980) Invertebrate Zoology (4th ed): Philadelphia, Saunders College 1089 p.
- Barwis, J.H. (1985) Tubes of Modern polychaete *Diopatra cuprea* as current velocity indicators and as analogs for *Skolithos-Monocraterion* , In Curran,H.A. (ed) Biogenic structures: Their use in interpreting depositional environments: SEPM, N0-35; 225-235.
- Bambach, R.K. (1977) Species richness in marine benthic habitats through the phanerozoic, Paleobiology, V-3; 152-167.
- Bhatt, N.Y. (1996) Ichnologic and Stratigraphic events as revealed by Mesozoic Sedimentary rocks exposed south and Southeast of Bhuj, Kutch, Gujarat State, Unpubl., Thesis., M.S.University of Baroda, Vadodara. 547 p.
- Bindra, S.S. (1927) The fauna of Karachi: A study of Genus Eurythoe. Mem. Dept. Zool. Punjab University-I, pp-1-18.
- Biswas, S.K., (1965) A new classification of the Tertiary rocks of Kutch, Bull.Geol. Min.Met. Soc. India. V-35, pp. 1-6.
- Biswas, S.K, (1971) Note on geology of Kutch. Quart. Jour. Geol. Min. Metal. Soc., V-43, pp. 223-235.
- Biswas, S.K. (1972) A time stratigraphic classification of the tertiary rocks of Kutch,- Revision and amendments. Quatr. Jour. Min. Met. Soc. Ind., V-44, 221-224.

- Biswas, S. K. (1974). Landscape of Kutch-a morphotectonic analysis. Ind. Jour. Earth. Sci., V-1, pp. 177-190.
- Biswas, S.K., (1977), Mesozoic rock stratigraphy of Kutch, Quart. Jour. Geol. Min. Met. Soc. India, V-49, pp-1-52.
- Biswas, S.K., (1978) On status of the Bhuj and Umia series of Kutch, W. India. Proc. 7th Ind coll. Micro. Strati., Madras.,
- Biswas, S.K., (1980) Structure of Kutch-Kathiawar Region, W.India. Proc. 3rd Ind. Geol. Congress, Pune. pp 255-272.
- Biswas, S.K. (1982) Rift basins in Western margin of India and their Hydrocarbon prospects with special reference to Kutch basin, Ann. Assoc. Pet. Geol. Bull. V-66; 1496-1513.
- Biswas, S.K. (1987) Regional tectonic framework, structure and evolution of India, with special reference to western margin basins of India. Tectonophy. V-135; 307-327.
- Biswas, S.K. and Deshpande, S.V., (1968), The basement of the Mesozoic sediments of Kutch, Western India. Bull. Min. Met. Soc. India, V-40, pp 1-7.
- Biswas, S.K. and Deshpande, S.V., (1970), Geological and tectonic map of the Kutch. Bull. Oil & Natu. Gas. Comm. V-7, pp.115-116.
- Biswas, S.K. and Deshpande, S.V., (1982), A note on mode of eruption of the Deccan Trap lavas with special reference to Kutch. Jour. Geol. Soc. Ind. V-14, pp. 134-141.
- Biswas, S.K. and Khatri, K.N. (2002) A geological study of earthquake in Kutch, Gujarat, India. Jour. Geol. Soc. Ind. V-60; 131-140.
- Biswas, S.K. and Raju, A.T.R., (1973) The Rock Stratigraphic classification of the Tertiary sediments of Kutch. Bull. O.N.G.C., V-10, pp-37- 46.
- Blair, T.C. and Mc Pherson, (1994) Alluvial fan and their natural distinction from river based on morphology, hydraulic processes, sedimentary processes and facies assemblages. Jour. Sedim. Res. V-64, pp. 450-489.
- Blandford, W.T. (1867) On the geology of a portion of Cutch, Mem. Geol. Sur. Ind. Vol - VI, Pt.2., pp-17-38.
- Bluck, B. J. (1967) Sedimentology of the Beach gravels: examples from south Wales, Jour. Sedim. Petrol. V- 37, pp 128-156.
- Boccaletti, M., Dainelli, P., Angelucci, A., Arush, M.A., Cabdulqaadir, M.M., Nafissi, P., Piccoli, G. and Robba, E. (1988) Folding of the Mesozoic cover in SW Somalia: a compressional episode related to the early stages of Indian Ocean evolution. Jour of Petrol. Geol. V-11; pp157-167.

- Bock, M.J. and Miller, D.C. (1996) Fluid flow and suspended particulate as determinants of polychaete feeding behaviour, *Jour. Mar. Res.* V-54, pp. 565-588.
- Bouma, A.H. (1964) *Methods for the study of sedimentary structures*, Wiley interscience, 458 p.
- Brenchley, G.A. (1976) Predator detection and avoidance: ornamentation of tube caps of *Diopatra* sp. (Polychaeta-Onuphidae): *Mar. Biol.* V-38, pp. 179-188.
- Bromley, R.G. (1970) Borings as trace fossils and *Entobia cretacea* Portlock, as an example. In T.P. Crimes, J.G. Harper (eds.) *Trace fossils*, Seel House, press, 49-90.
- Bromley, R.G. (1975): Trace fossils at omission surfaces- In Frey, R.W. (Ed) *The Study of trace fossils*, 399-428; Springer - Verlag, New York.
- Bromley, R.G. (1981) Concepts in ichnotaxonomy illustrated by small round holes in shells. *Acta Geologica Hispanica*, V-16(1); 55-64.
- Bromley, R.G. (1990): *Trace fossils-Biology and Taphonomy*, 280p; Unwin Hayman Limited, London.
- Bromley, R.G. (1992) Bioerosion: eating rocks for fun and profit. In C.G. Maples and R.R. West (eds.) *Trace fossils. Paleontological society short course in paleontology*, V-5, 121-129
- Bromley, R.G. (1994) The paleoecology of bioerosion, in *The Paleobiology of trace fossils* (eds. S.K. Donovan), Wiley, Chichester, pp. 134-154.
- Bromley, R.G. (1996) *Trace Fossils: Biology, Taphonomy and Applications* (2ed.) Chapman and Hall, London, 361p.
- Bromley, R.G. and D'Alessandro, A. (1987) Bioerosion of the Plio-Pleistocene transgression of southern Italy. *Rivista Italiana di Paleontologia e stratigrafia*, V-93(3), 379-442.
- Bromley, R.G. and Ekdale, A.A. (1984) Chondrite: a trace fossil indicator of anoxia in sediments. *Science*, V-224, 872-874.
- Bromley, R.G. and Fursich, F.T. (1980) Comments on proposed amendments to the International Code of Zoological Nomenclature regarding ichnotaxa. *Z.N. (S.)* 1973) *Bull. Zoo. Nomen.* V37, 6-10.
- Bronn, H.G. (1837) *Lethaea geognostica oder Abbildungen und Beschreibungen der für die Gebirgs formationen bezeichnendsten Vertheilungen*: V-1; p. 1-672, Atlas, Pl. 1-47, E. Schweizerbart.
- Bull, W.B. (1968) Alluvial fans. *Jour. Geol.* V-16, pp 101-106.
- Burns, T.P. (1989) Lindeman's contradiction and the trophic structures of ecosystems. *Ecology*, V-70(5), 1355-1362.

- Cadee, G.C. (1979) Sediment reworking by the polychaete *Heteromastus filiformis* on a tidal flats in the Dutch Wadden sea, *Neth.J. Sea Res.* V-13; 441-456.
- Cammen, L (1980a) A method for measuring ingestion model for deposit feeders and its use with the polychaete *Nereis succinea*, *Estuaries*, 3: 55-60.
- Cammen, L (1980b) Ingestion rates: An empirical model for aquatic deposit feeders and detritivores, *Oecologia*, V-44; 303-310.
- Carter, H.J., (1853) Description of some of the larger forms of fossilized Foraminifera in scinde, with observation on their internal structure. *Ann. Mag. Nat. Hist. Ser. 2.* (Also *J.Bom. Br. Roy. As. Soc.*) V-5, pp 168-174.
- Carter, R.W.G. (1986). The Morphodynamics of Beach and Ridge Formation: Magilligan, Northern Ireland. *Mar. Geol.* v. 73, pp. 191-214.
- Chace, F. A. and Hobbs, H.H. (1969) The fresh water and terrestrial decapods crustaceans of the West Indies, With special reference to Dominica, United States National Museum Bulletin, V-292, 258 p.
- Chakrabarti, A. (1981) Burrow pattern of *Ocypode ceratophthalma* (Pallas) and their environmental significance. *Jour. Paleo.* V-55, 431-444.
- Chakrabarti, A (1972) Beach structures produced by crab pellets. *Sedimentology* V- 18, pp 129- 134.
- Chakrabarti, A (1981) Burrow patterns of the ocypode ceratophthalma (Pallas) and their environmental significance *Jour. Paleon.* V-55, pp 431-441.
- Chamberlain, C.K & Baer, J.L. (1973): Ophiomorpha and a new thalassinid burrow from Permian of Utah-Brigham Young University Geological Studies., 20:79-94; Brigham.
- Chance, F.A. Jr. and Hobbs, H.H. Jr. (1969) The freshwater and the terrestrial decapod crustaceans of the West Indies, with special reference to Dominica, United States National Museum Bulletin, V-292, 258.p.
- Chappgar, B.F. (1957): On the marine crabs (Decapoda: Brachyura) of Bombay state. Part-I. *Jour. Bom. Nat. Hist. Soc.*, 54(2): 399-439, pls. A, 1-11, 2 test figs.
- Chappgar, B.F. (1957): On the marine crabs (Decapoda: Brachyura) of Bombay state. Part-II. *Jour. Bom. Nat. Hist. Soc.*, 54(3): 503-549, pls. B. 12-16, 1 text, fig.
- Chhapgar, B.F. (1958): Addition to the crabs fauna of Bombay state. *Jour. Bom. Nat. Hist. Soc.*, 55(3): 582-585, pls. i, ii.
- Chapman, G (1949) The thixotropy and dilatancy of a marine soil. *Jour. Mar. Biol. Assoc. U.K.* V-28, 123-140.

- Chapman, G and Newell, G.E. (1947) The Role of the body fluid in relation to the movement of the soft-bodied invertebrates-I, The burrowing of *Arenicola*. Proc. Roy. Soc. B, V134, 431
- Chapman, G. (1950) Of the Movement of Worms. Jour Exp. Biol ;V-27; pp 29
- Chauhan, O.S. and Almeida, F (1993) Influence of Holocene sea level, regional tectonics and fluvial gravity and slope currents induced sedimentation on the Regional geomorphology of the continental slope off northwestern India. Mar. Geo. V-112, pp 313-328.
- Chauhan, O.S., Almeida, F. and Moraes C. (1993) Regional Geomorphology of the continental slope of NW India of signatures of Deep seated structures. Mar. Geol. V-15. pp283-296.
- Cladwell, R.L. and Dingle, H. (1975) Ecology and evolution of agnostic behaviour in stomatopods. Naturwissenschaften, V-62.,pp 214-222.
- Cladwell, R.L. and Dingle, H. (1976) Stomatopods. Scientific American, V-234, pp 80-89.
- Clark, R.B. (1964) The eyes and the photonegative behaviour of the *Nephtys* (Annelida, Polychaete). Jour. Exp. Biol, V-3, pp. 460-477.
- Clifton, H.E. (1973) Pebble segregation and bed lenticularity in wave-worked versus alluvial gravel. Sedimentol. V-20, pp 173-187.
- Codez, J., and Saint Seine, R. (1958) Revision des Cirripedes Acrothoraciques fossiles: Soc. geol. France. Bull., Ser., 6, V-7; p. 699-719.
- Curran, H.A. (1992). Trace fossils in Quaternary, Bahamian-style carbonate environments: the modern to fossil transition, in Maples, C.G and West, R.R. (eds) Trace fossils, short course in paleontology, No-5.
- Curran, H.A. (1994). The paleobiology of ichnocoenoses in Quaternary, Bahamian-style carbonate environments: the modern to fossil transition, in The Paleobiology of trace fossils (eds. S.K. Donovan), Wiley, Chichester, pp. 83-104.
- Dabrio, C.J. and Polo, M.D. 1981 Flow regime and bedform in a ridge and runnel system, SE. Spain; Sed. Geol. V-28, pp 97-110.
- Dam, G (1990) Paleoenvironmental significance of trace fossils from the shallow marine Lower Jurassic Neill Klint Formation, East Greenland. Paleogeogr. Paleoclim. Paleoecol. V-79, 221-248
- Dauer, D.M; Maybury, C.A; and Ewing, R.M (1981) Feeding behaviour and general ecology of several spionid polychaetes from the Chesapeake bay. Jour. Exp. Mar. Biol. Ecol. V-54; 21-38.

- De, C. (1998) Biological reworking of sediments by crabs: A case study for erosion of the Digha beach, West Bengal. *Curr. Sci.*, V-75(6); 617-620.
- DeBoers, P.L. (1979). Convolute lamination in modern sands of the estuary of the Oosterschelde, the Netherlands, formed as a result of entrapped air. *Sedimen.*, v.26, pp. 283-294.
- DeMan.J.G. 1887 : Report on the Podopathalmous crustacea of Mergui Archipelago collected for the Trustees of the Indian Museum, Calcutta by Dr. John Anderson, F.R.S., Suprintendent of the Museum Parts i-v Jour. Linn. Soc. London (Zoology) xxii, pp 1-212, pls.i-xiv
- Desai and Patel (in press) Kachchh Earthquake of 2001: Seismically triggered deformed structures in the intertidal sediment of Mandvi coast in the Gulf of Kachchh, Gujarat, *Jour. Geol. Soc. Ind*
- Desai and Patel (communicated) Trace fossil assemblages of Tectonically Uplifted Holocene Shorelines, Kachchh, Western India.
- Desai, B.G., Patel, S.J and Chappgar, B. F. (In Press) On two interesting marine crabs from Mandvi (Decapod: Braychura) from Mandvi Kutch. *Jour. Bom. Nat. Hist. Soc.*
- Dobkin, J.E. and Folk, R.L. (1970) Shape development on Tahiti-nui. *Jour. Sedi. Petrol.* V-40, 1161-1203.
- Dolan, R., Vincent, L. And Hayden, B. (1974). Crescentic coastal landforms, *Z. Geomorp.*, v.18, pp. 1-12.
- Donaldson, D. and Simpson, S. (1962) Chomatichnus, a new ichnogenus and other trace fossils of Wegber Quarry. *Liverpool & Manchester. Geol. J.* V-3; 73-81.
- Dorjes, J and Hertweck, G. (1975) Recent biocoenoses and ichnocoenoses in shallow water marine environments. In *The Study of Trace fossils*, R.W.Frey (ed)., 459-91., New York: Springer Verlag.
- Doty, M.S. (1946) Critical tide factors that are correlated with the vertical distribution of marine algae and other organism along pacific coast. *Ecology*, V-27;315-328.
- Dyer, K.R.; Christie, M.C.; Wright, E.W: (2000) The classification of the Intertidal Mudflats, *Continental shelf research*, V-20; 1039-1060.
- Ehernberg, K. (1944) Eraganzende Bemerkungen zu den seinerzeit aus dem Miozan Von Burgschleinitz beschriebenen Gangkernen und Bautes de Kapoder Krebse, *Z.Paleont.*, V-23, pp-354-359.
- Ekdale, A..A (1985) Paleoecology of marine endobenthos . *Paleo 3*, V-50 pp 63-81.
- Ekdale, A.A (1992) Muckraking and Mudslinging: the joys of deposit feeding, in Maples, C.G and West, R.R. (eds) *Trace fossils, short course in paleontology*, No-5. 145-170.

- Ekdale, A.A. and Lewis, D.W. (1991) Trace fossils and paleoenvironmental control of ichnofacies in a late Quaternary gravel and loess fan delta complex, New Zealand. *Paleogr. Paleocli. Paleoecol.* V-81; 253-279.
- Ekdale, A.A.; Bromley, R.G; and Pemberton, S.G. (1984) *Ichnology- the use of trace fossils in sedimentology and stratigraphy*, SEPM, Short course, 15.
- Elliott, T. and Gardiner, A.R. (1981) Ripple, megaripple and sand wave bedforms in the macrotidal Loughor Estuary, South Wales, U.K. *Spec. Publ. Int. Assoc. Sedim.* V-5, pp 51-64.
- Fauchald, K. and Jumars, P.A. (1979) The diet of worms: a study of polychaete feeding guilds. *Ocean. Mar. Biol. Ann. Rev.* V-17; pp 193-284.
- Fauvel, E. (1932) *Anneliad, Polychaeta of Indian museum, Calcutta.* *Mem. Ind. Mus.* XII-1, p1-62.
- Fauvel, E. (1953) *The Fauna of India (Polychaeta).* Indian press, Allahabad.
- Fenton, C.L and Fenton, M.A (1932) Boring sponges in Devonian of Iowa: *Amm. Midland Naturalist.*, V-13, pp-42-54.
- Fillon, D and Pickerill, R.K. (1990) *Ichnology of Upper Cambrian? To Ordovician Bell island and Wabana groups of eastern Newfoundland, Canada-* *Paleonto. Canadiana.* V-7, 1-119.
- Folk, R.L. (1966) A review of grain size parameter. *Sedimentology.* V-6, pp 73-93
- Frey, R.W. (1975) *The study of trace fossils*, (Ed) Springer Verlag.
- Frey, R. W. (1987) Hermit crab: Neglected Factor in Taphonomy and Paleoecology., *Palios.*, V-2, pp 313-322
- Frey, R.W. and Howard, J.D. (1969) A profile of biogenic structures in a holocene barrier island-salt marsh complex, Georgia. *Trans. Gulf. Coast Assoc. Geol. Soc.* V-19; 427-44.
- Frey, R.W. and Howard, J.D. (1988) Beaches and beach related facies, Holocene barrier island of Georgia. *Geol. Magz.* V-125, pp621-640
- Frey, R.W. and Mayou, T.V. (1971) Decapod burrows in holocene barrier island beaches and washover fans, Georgia, *Senckn. Martima.* V-3, pp 53-77
- Frey, R.W. and Pemberton, S.G. (1987) The psilonichnus ichnocoenose, and its relationship to adjacent marine and non-marine ichnocoenoses along the Georgia coast, *Bull.Can. Petrol. Geol.* V-35, pp333-357.

- Frey, R.W; Howard, J.D; and Pryor, W.A. (1978) *Ophiomorpha*: its morphologic, taxonomic, and environmental significance. *Paleogr. Paleoclim. Paleoecol.*, V-23; 199-229.
- Frey, R.W; Curran, H.A; Pemberton, S.G. (1984) Trace making activity of crabs and their environmental significance: The tchnogenus *Psilonichmus*, *Jour. Paleon.* V-58, 333-350.
- Fursich, F.T (1971): Hartgrunde und Kondensation im Dogger von calvados- Neues Jahrb. Geol.Paleont. 138: 313-342; Abhandlungen.
- Fursich, F.T (1974) Corallian (Upper Jurassic) trace fossils from England and Normady., - Stuttgarter Beitrage Zur Naturkunde, Serie-B (Geollogie und Palaontologie)., V-13; 1-51.
- Fursich, F.T. (1981) Invertebrate trace fossils from the Upper Jurassic of Portugal. *Comunicacoes. Ser. Geol. de Portugal.* V-67, pp 153-168.
- Fursich, F.T. (1998) Environmental distribution of Trace fossils in the Jurassic of Kachchh (Western India). *Facies* V-39., pp.243-272.
- Fursich, F.T; Oschmann,W; Singh,I.B; Jaitly, A. (1992) Hardgrounds, reworked concretion levels and condensed horizons in the Jurassic of western India: their signifiance for basin analysis. *Jour. Geol. Soc.Lond.* V-149,pp 313-331
- Gedeon, P.W., Menon, P.K.B., Rao, S.R.V. and Jose, K.V. (1957) On marine fauna of the Gulf of Kautch: A preliminary survey. *Jour. Bom. Nat. Hist. Soc.*, V54, pp 630-706.
- Ghare, M.A. and Kulkarni, K.G (1986) Jurassic ichnofauna of Kutch-II: Wagad region. *Biovigayanam*
- sGlennie, K.W. (1970) Desert sedimentary environments. *Developments in sedimentology*, V-14., p 222, Elsevier, Amsterdam.
- Glennie, K.W. and Evans, G (1976) A reconaissance of the Recent sediments of the Ranns of Kutch, India. *Sedimentology* V-23., pp 625-647.
- Gingras, M.K & Pemberton, S.G & Saunders, T (2001): Bathymetry, sediment texture and substrate cohessiveness; their impact on modern *Glossifungites* trace assemblages at Willapa bay, Washington - *Paleogeogr. Paleoclim. Paleoecol.* , 169: 1-24.
- Gopalkrishnan, P. (1969a) On the Holothuroidea (echinodermata) of the Gulf of Kutch. *J. Bom. Nat. Hist. Mus. Soc.* V-66, 399-400.
- Gopalkrishnan, P. (1969b) Some marine algae from the Gulf of Kutch. *Phykos* V-8, pp61-67.
- Gopalkrishnan, P. (1970) Some observation on the shore ecology of the Okha coast. *J.Mar. Biol. Ass. Ind.* V- 12, pp 15-34.

- Grant, C.W. (1837) Memoirs to illustrate a geological map of Kutch. Trans.Geol. Soc.
- Grassle, J.F. and Sanders, H.L. (1973) Life histories and the role of disturbances. Deep Sea Res. V-20, 643-659.
- Grassle, J.F and Grassle, J.P (1974) Opportunistic life histories and genetic systems in marine benthic polychaetes. Jour. Mar. Res. V-32:253-284.
- Gravely, F.H. 1927: Crustacean In : The Littoral fauna of Kutch Island in the Gulf of Manaar. Bull. Madras. Govt. Mus.(n.s.)i, no1, pp. 145-155, pls.xx- xxvi.
- Gray, J. and Lissman, H.W. (1938) Studies in animal locomotion VII. Locomotory reflexes in the earthworm. J.exp. Biol. V-15, 506.
- Greenwood. J.G 1972 The mouthparts and feeding behaviour of two species of hermit crabs. J. Nat.Hist. 6,325-337.
- Guha, D.K. (1968) Ostracods from Middle Eocene of Kutch. Gujarat State, Western India. O.N.G.C. Bull., V-5.,pp-83.
- Gupta, S.K. (1975) Silting of the Rann of Kutch during Holocene. Ind jour. earth sci., V-2(2) pp- 163-175.
- Hagenow, K.F. Von (1840) Monographie der Rugenschen Kreideversteinerungen II. Abth. Radiarien u. Annulaten: Neues Jahrb. Miner., Geog. Geolog., Petrefaktenkd., 1840, 631-672.
- Haldeman, S.S. (1840) Supplement to number one of "A monograph of the Limniades and other freshwater univalve shells of North America: containing new description of apparently new animals in different classes, and the names and characters of the subgenera in Paludina and Anculosa: 3p (Philadelphia).
- Hall, J. (1847) Paleontology of New York Vol-1, C; Van Benthuyssen, Albany, 338p.
- Hallam, A. (1975) Preservation of trace fossils, pp 55-63. In Frey, R.W. (Ed) The study of trace fossils. Springer Verlag, New york.
- Hallam, A and Swett, K (1966) Trace fossils from the Lower Cambrian pipe rocks of the north-west highlands: Scot. Jour. Geol. V-2; 101-106.
- Hamano,T., Torisawa, M., Mitsunashi, M And Hayashi, K. (1994) Burrows of Stomatopod crustacean *Oratosquilla oratoria* (DeHann,1844) in Ishikari Bay, Japan. Crust. Res. v. 23, pp 5-11.
- Hannibal, J.T. and Feldmann, R.M. (1983) Arthropod trace fossils, interpreted as echinocarid escape burrows, from the Chagrian shales (Late Devonian) of Ohio. Jour.Paleon. V-57, pp 705-716.

- Hantzschel, W. (1962) Trace fossils and problematica. In Treatise on Invertebrate Paleontology R.C.Moore (Eds)., W., 177-245. New York and Lawrence, Kansas; Geol. Soc.Ammer.& Kansas University Press.
- Hantzschel, W. (1965) Vestigia invertebratorum et problematica Fossilium catalogus I: Animalia, V-108, 142.p.
- Hantzschel, W. (1975) Trace fossils and problematica. In Treatise on Invertebrate Paleontology 2nd Edition, C.Teachert (ed) W.Boulder, Colorado and Lawrence, Kansas; Geol. Soc.Ammer.& Kansas University Press.
- Hantzschel, W. and Kraus, O. (1972) Names based on trace fossils (ichnotaxa): request for a recommendation Z.N. (s) 1973 Bull. Zoo.Nomen. V 29, 137- 141.
- Harris, C.R. (1974). A quantitative study of compaction with depth in a vesicular sand layer in an intertidal area of Dublin bay, Ireland. Jour. Sed. Petrol., v.44, pp.1024-1028.
- Hayes, M.O. (1969) Coastal Environments., N.E. Massachusetts and New Hampshire. Univ. Of Massachuettes Amherst, Mass., 462 pp.
- Hayes, M .O. 1972 Forms of sediment accumulation in the beach zone, In R.E.Mayers (Ed.)Waves on Beach,4, Academic press, New York., pp 297-356.
- Hayes, M.O. and Boothroyd, J.C. 1969 Stroms as modifying agents in the coastal environment. In Coastal Research group, Coastal environment, Northeastern Massachusetts and New Hampshire, S. E.P.M. Field trip guide book,
- Hempton, M.R. and Dewey, J.F (1983) Earthquake- induced deformational structures in young lacustrine sediments East Anatolian fault, Southeast turkey. Tectonophysics. v. 98, 7-14.
- Henderson, J.R. 1893: A Contribution to Indian Cracinology, Trans. Linn. Soc. London (Zool.) (2)V, pp. 325-409, pls. 36-38.
- Hertweck, G (1972) Georgia coast region, Sapelo island, USA: Sedimentology and Biology 5. Distribution and environmental significance of lebensspuren and insitu skeletal remains. Senkenbergiana maritima, V-4; 125-167
- Hine, A.C. 1979 Mechanics of berm Development and resulting beach growth along a barrier spit complex. Sedimentology. V 26, pp 333-351
- Hill, G.W. and Hunter, R.E. (1979) Burrows of the Ghost crab *Ocypode quadrata* (Fabricus) on the barrier islands, south-central Texas coast. Jour. Sed. Petrol. V-43, 24-30.
- Hornell, J. (1909) Report of the Goverment of Baroda on the marine zoology of Okhamandal. London.1

- Howard, J.D. and Frey, R.W. (1975) Regional Animal-sediment characteristic of Georgia estuaries. *Senckenberg. Marita.* V-7; 33-103.
- Howard, J.D. and Frey, R.W. (1984) Characteristic trace fossils in nearshore to offshore sequences, upper Cretaceous of east-central Utah. *Can. Jour. Earth. Scie.* V-21; 200-219.
- Howard, J.D. (1972): Trace fossils as criteria for recognizing shorelines in stratigraphic record- In Rigby, J.K., and Hambill, W.M.K. (Eds) *Recognition of ancient sedimentary environments.* Society of Economist Geologist and Paleontologist Mineralogy. Special Publication, 16: 215-25.
- Howard, J.D. (1966) Characteristic trace fossils in the upper Cretaceous sandstones of the Book Cliff and Wasatch Plateau. *Utah Geol. And Mineral Sur. Bull.* V-80, pp 35-53.
- Howard, J.D. and Singh, I.B. (1985) Trace Fossils in the Mesozoic sediments of Kachchh, Western India. *Paleogeogr. Paleoceno. Paleoecol.*, V-52., pp 99- 122.
- Hyt, J.H. and Henry, V.J. (1964). Development and geological significance of the soft beach sand. *Sedimen.*, v. 3, pp. 44-51.
- International Commission of Zoological Nomenclature (Stoll, N.R.; Dollfus, R. Ph., Forest, J.; Riley, N.D.; Sabrosky, C.W.; Wright, C.W.; Melville, R.V.; Eds) (1964) *Code international de Nomenclature Zoologique adopte par le XVe Congres International de Zoologie = International Code of Zoological Nomenclature adopted by the XV international congress of Zoology (2nd Ed.).* International trust for Zoological Nomenclature, London, 176p.
- International Commission of Zoological Nomenclature (Ride, W.D.L.; Sabrosky, C.W.; Bernardi, G., and Melville, R.V.; Eds.) (1985) *Code international de Nomenclature Zoologique adopte par le XXe Assemblee Generale de l'Union Internationale des Sciences Biologiques = International code of Zoological Nomenclature adopted by the XX general Assembly of the international Union of Biological sciences (3rd. Ed.).* International trust for Zoological Nomenclature and British Museum (Natural History), London., 338 p.
- Jablonski, S. and Bottjer, D. (1991) Environmental patterns in the origin of higher taxa: the post-paleozoic fossil records. *Science.* V-252, 1831-1835.
- Jain, S.P. (1978) Recent ostracods from Mandvi beach, West coast of India. *Bull. Ind. Geol. Asso.* V-11, pp 91-139.
- Jain, S.P. and Bhatia, S.B. (1976) Recent benthonic foraminifers from Mandvi, Proc. VII, ICMS, Banaras, India
- Jenkins, R.J.F. (1975) The Fossil crab *Ommatocaricus corionis* (Cresswell) and a review of related Australian species. *Memoirs, National Museum of Victoria*, V-36, 33-62.

- Johnson, D. W., (1919). Shore Process and Shoreline Development, pp. 518-519. John Wiley and Sons Inc.
- Jones, M.L. (1961) A quantitative evaluation of the benthic fauna off point Richmod, California. Univ. Calif. Pub. Zoo. V 67, pp 219-320.
- Jux, U. (1964) Kommensalen oserdevonischer Atrypen aus Bergisch (Rheinisches Schiefergebirge). Neus. Jharb. Geol. Paleo. 675-687.
- Kar, A . (1988) Evidence for neotectonism from the Indian desert. In S.Singh and R.C.Tiwari (Eds) Geomorphology and Environment. Allahabad Geogr.Soc. Allahabad., pp 300-310.
- Kar, A . (1993). Neotectonic influence on morphologic Variations along the coastline of Kachchh India, Geomorph., v.8, pp. 199-219.
- Kar, A. (1999) Cluster analysis of grain size information for possible linkages between geomorphic environments: some case studies from Indian arid zone Z.Geomorp. N. F., V-43(1), pp 27-39.
- Kazmierczak and Pszczolkowski, (1969) Burrows of Enteropneusta in Muschelkalk (middle Triassic) of the Holy Cross mountain, Poland. Acta. Paleo. Polo. V-XIV, 299-315
- Kelly, S.R. and Bromley, R.G (1984) Ichnological nomenclature of Calvate borings. Paleontology, V-27., 793-807.
- Kemp, S. (1915) Fauna of the Chilka lake. Mem. Ind. Mus. V-5, pp 199-325.
- Kemp, S. (1919) Fauna of the Chilka lake. The indian species of Macropathalmus. Rec. Ind. Mus.
- Kennedy, J.F. (1961) Stationary waves and antidunes in alluvial channel. W.M. Keck. Laboratory of hydraulics and water research, California Institute of Technology, Rep. KH-R-2, 146p
- Kindle, E.M. (1935). Dominant factors in formation of firm and soft beaches Jour. Sed. Petrol., v. 6, pp. 16-22.
- King, C.A.M and Williams, W.W. (1949) The formation and movement of sand bars by wave action: Geogr.Jour., V-113; pp70-85.
- Klein, G. De Vries (1985) Intertidal flats and intertidal sand bodies. In Davis, R.A. (Eds)., Coastal sedimentary environments. New York. pp.187-224.
- Kohli, G.R. (1921-'22): Brachyrua of Karachi coast. Proc. Lah.Phil.Soc. iii, pp.81-85.
- Kozloff, E.N. (1996): Seashore life of the northern Pacific Coast- University of Washington Press, 370 (Fourth Printing); Washington.

- Kraeuter, J. (1976) Biodeposition by salt marsh invertebrates, *Mar. Biol.* V-35; 215-223.
- Ksiazkiewicz, M. (1977), Trace fossils in the Flysch of the Polish Carpathians. *Paleont. Polonica.* V-36; 1-208
- Kulkarni, K.G and Ghare, M.A (1989) Stratigraphic distribution of ichnotaxa in the Wagad region of Kutch, India., *Jour. Geol. Soc. Ind.*, V-33(6), pp. 259-267.
- Kulkarni, K.G and Ghare, M.A. (1991) Locomotary traces (Repichnia) from the Jurassic sequence of Kutch, Gujarat., V-37., pp 374-387.
- Kumar, N and Sanders, J.E. 1976 Characteristics of shoreface deposits: Modern and ancient. *Journal of sedimentary petrology*, 46: 145-162.
- Kundu, H.L. (1965) The marine fauna of the Gulf of Kutch, III, Pelecypods. *J. Bom. Nat. Hist. Soc.* V-62, pp83-103
- Kundu, H.L. (1965) The marine fauna of the Gulf of Kutch, III, Pelecypods. *J. Bom. Nat. Hist. Soc.* V-62, pp211-236
- Lemech, H. (1973) Comments on the application concerning Trace fossils. *Bull. Zoo. Nomen.*, V 30, 70p
- Lessertissier, J. (1955) Trace fossiles d'activite animale et leur signification paleobiologique: *Soc. Geol. France, Mem, n.ser.*, V-74, p-1-150.
- Leszczynski, S.; Uchman, A. And Bromley, R. (1996) Trace fossils indicating bottom aeration changes: Folsz limestone, Oligocene, Outer Carpathians, Poland. *Paleo-3*, V-121, pp 79-87.
- Levins, R. (1968) *Evolution in changing environments.* Prince. Univ press, Princeton. 120.
- Leymerie, M.A. (1842) Suite du memoire sur le terrain Cretace du Departement de l'Aube: *Soc. Geol. France., Mem.*, V-5, p, 1-34.
- Levinton, (1972) Stability and trophic structure in deposit feeding and suspension-feeding communities. *Amm. Natur.* V-106; 427-486.
- Livingston, J.S. (1989) Deposit feeding and coastal oceanography. In Lopez, G; Taghon, G., Livingston, J.S. (Eds) *Ecology of Marine depositfeeders. Lecture notes on coastal and estuarine studies*, Springer, New York, pp 1-23.
- Livingston, J.S. and Bambach, R.K. (1975) A comparative study of the Silurian and recent deposit feeding bivalve communities. *Paleobiol.* V-1, 97-124.
- Lundgren, B. (1891) Studier ofver fossilforandelosa block, *Geologiska Foreningens I Stockholm Forhandlingar*, V-13, pp 111-121.

- Lyall, A.K. and Reddy, V.V. (1982) Quaternary geology of the Gulf of Kachchh basin. Pp 203-224 In S.S. Merh (ed) Recent researches in Geology (9) Hindustan Publishing Corp. New Delhi.
- Lyell, C. (1853): Principles of Geology-11th edition Appleton and Co, London.
- MacGinitie, G.E. and MacGinitie, N. (1949) Natural History of Marine animals. McGraw-Hill, New York.
- MacMurdo, J. (1839): Observations of the Sindoo or River Indus. Bombay Geographical Society Proceedings, 5:124p; Mumbai.
- Macsotay, O. (1967) Huellas problematicas y su valor paleoecológico en Venezuela. GEOS (Venezuela), V-16, pp 1-87.
- Malik, J.N. & Sohoni, P.S. & Karanth, R.V. & Merh, S.S. (1999): Historic and Modern Seismicity of the Kachchh peninsula, Western India- Journal of Geological Society of India, 54:545-550 Bangalore.
- Malik, J.N. & Sohoni, P.S. & Merh, S.S. & Karanth, R.V. (2001): Active tectonic control on alluvial fan architecture along Kachchh mainland hill Range, Western India- Zeitschrift Geomorph., V-45, 81-100.
- Magnum, C.P. (1964a) Studies on speciation in Maldanid polychaetes of the North American Atlantic coast. II. Distribution and competitive interaction of five sympatric species, Limnol. Oceanogr. 9; 12-26.
- Magnum, C.P. (1964b) Activity pattern in metabolism and ecology of polychaetes, Comp. Biochem. Physiol. 11; 239-256.
- Mangum, C.P., Santos, S.L., and Thodes, W.R. (1968) Distribution and Feeding in the onuphid polychaete, *Diopatra cuprea* (Bosc), Marine Biology, V-2., pp 33- 40.
- Marshall, N. (1970) Food transfer through the lower trophic levels of the benthic environment. In Steel, J.H. (ed) Marine food chains, 52-66. Univ. Calif. Press. Berkeley.
- Martin, D., Ballesteros, E., Gilli J.M., and Palacin, C. (1993) Small scale structure of infauna polychaete communities in an estuarine environment. Methodological approach- Estuar. Coast. Shelf. Sci. V-36, pp 47-58.
- Martinsson, A (1965) Aspects of a middle Cambrian thanatopora on Oland: Geol. Foren. Stockholm. Forhandl, V-87, 181-230
- Martinsson, A (1970) Toponymy of trace fossils: in trace fossils, Crimes, T.P and Harper, J.C. (eds) Geol. Jour. Sp. Pub. Issue No.3; 323-330.
- Martinsson, A (1972) [Discussion] in W. Hantzschel and Kraus, O, Names based on trace fossils (ichnotaxa): request for a recommendation, Z.N. (s), 1973, Bull. Zoo. Nomencl. V-29, p140.

- McKee, E.D. (1953) Report on studies of stratification in modern sediments and in laboratory experiment. Office of Naval Research, Project, Nonr. 164(00), Nr. 081 123, 61p.
- McPherson, J.G., Shanmugam, G. and Molia, R.J. (1987) Fan deltas and braid deltas: Variety of the coarse grained deltas. *Geol. Soc. Am. Bull.* V-99, pp 331-330
- Meddicott, H.B and Blandford, W.T (1879-'87). *Manual of Geology of India*, 4 Volumes, G.S.I., Govt.Of India press, Calcutta.
- Melville, R.V. (1979) Further proposed amendments to the International Code of Zoological Nomenclature Z.N (G) 182. *Bull.Zoo.Nomen.*, V-36, pp 11- 14
- Menon, P.K.B., Dattagupta, A.K. Dasgupta, D., (1961) The marine fauna of the gulf of Kutch III Gastropoda. *J.Bom.Nat.Hist. Soc.* V-58, pp 475-494.
- Merh, S.S. and Chamyal, L.S. (1993) Quaternary deposits of Gujarat. *Curr. Sci.* V-67, pp. 823-827.
- Merh, S.S. and Patel, P.P. (1988) Quaternary geology and Geomorphology of the Rann of Kutch, In *Proc. Sem. On Recent Quaternary studies in India*, M.S.University, Baroda pp 377- 393.
- Miller, W. III (1998) Complex marine trace fossils. *Lethaia.* V-31, 29-32.
- Miller, D.C.; Bock, M.J.; and Turner, E.J., (1992) Deposit and suspension feeding in Oscillatory flows and sediment fluxes. *Jour. Mar. Res.* V-50; 489-520.
- Miller.D.C 1961 The feeding mechanism of fiddler crabs with ecological consideration of feeding adaption, *Zoologica.N.Y* 46, 89-100.
- Mohan, M. and Bhatt, D.K. (1968) Burdigalian Foraminifers from Kutch, India. *Proc. Nat. Inst. Sci. Ind.*, V-34, 159-190.
- Mohan,M. and Soodan, K.S (1967) *Inordinutosphaera*- a new genus of Globigirina. *Bull. Geol. Soc.Ind.*, V-4., p-22-25.
- Mohan, M and Soodan, K. S. (1970) Middle Eocene planktonic foraminiferal zonation of Kutch, India, *Micropal.*, Vol-18 (1)., pp 37-56, pls 1-2.
- Moore, J.N.; Fritz, W.J.; Futch, R.S.(1984) Occurance of megaripples in a ridge and runnel system, Sapelo island, Georgia: Morphology and Processes. *Jour. Sedi. Petrol.*, V-54; 615-625.
- Muley, N.V and Gupta, A.K. (1982) Study of sediment distribution in the gulf of kachchh from landset imageries pp 42-48, In S.S.Merh (ed) *Recent researches in Geology* (9) Hindustan Publishing Corp. New Delhi.
- Myers, A.C. (1979) Summer and Winter burrows of a Mantis shrimp, *squilla empusa*, in Narragansett Bay, Rhode island (U.S.A); *Estur. Coast. Mar. Sci.* V-8; 87-98.

- Myers, A.C.(1972)Tube-worm sediment relationships of *Diopatra cuprea* (Polychaeta-Onuphidae), *Marine Biology*, V-17, pp 350-356.
- Myrow, P.M. and Southard, J.B. (1991) Combined-flow model for vertical sequence in shallow marine strom deposited beds. *Jour. Sed. Petrol.* V-61(2)., 202-210
- Nair, R.R., Hashmi, N.H., and Rao, P.C. 1982. On possibility of high velocity tidal stream as dynamic barrier to the longshore transport:evidence from the continental shelf off the Gulf of Kutch. *Marine Geology*, V-47, pp. 77-86.
- Nelson, R.E.(1845): Notice of an earthquake and probable subsidence of the land in the district of kutch near the mouth of the Koree or eastern branch of Indus in June 1845-Geographical Society of London Quaternary Journal,103; London.
- Nemec, W. and Steel, R.J. (1984) Alluvial and coastal conglomerates: Their significant features and some comments on gravelly mass-flow deposits; In Koster, E.H. and Steel, R.J. (Eds) *Sedimentology of Gravels and Conglomerates*. Can.Soc.Petrol. Geol. Mem. 10, pp 1-31.
- Nigam, R. and Chaturvedi, S. (1999) Foraminiferal study from Karo creek, Kachchh (Gujarat), north western coast of India. *Ind.Jour. Mar. Sci.* V-29, pp. 133-138.
- Oldham, R.D. (1893) *Manual of the Geology of India*, GSI publications, Calcutta India.
- Osgood, R.G. (1975) Trace fossils from Cincinnati area. *Paleont. Ammer.* V-6, pp. 281-444.
- Otsuka, T., Takahama, N., Nakasto, N.and Nomura, S. (1997) Ancient liquification and their significance in the late quaternary tephra layers in the Karasu river valley, Gunma prefecture, Japan. *Quat. Res.* V-36, pp. 309-327.
- Ouchi,S (1985) response of alluvial rivers to slow active tectonic movement; *Geol.Soc. Amer. Bull.* V- 96, pp. 504-515.
- Owens, E.H And Frobels, D.H. (1977). Ridges and Runnels Systems in Magdalen Island Quebec, *Jour. Sed. Petrol.*, V- 7, pp.191-198.
- Pandey, J (1982) Chronostratigraphic correlation of the Neogene Sedimentaries of Western Indian Shelf, Himalayan and Upper Assam. *Paleontological Society of India, Sp., Publ*,
- Pandey, J (1986) Some recent Paleontological studies and their implications on cenozoic stratigraphy of Indian subcontinent. *Bull. O.N.G.C.*, V-23, pp. 1-44.
- Patel, M.P. and Allhabadi, B.B. (1988) The aeolinites of Kutch with special reference to their diagenesis. *Proc. Nat. Sem. Recent. Quaternary studies in India*, M.S.University, Baroda, pp. 37- 63.

- Patel, S. J. (1990) Study of trace fossils in carbonate rocks of Western Kutch-Gujarat State. Unpublished thesis submitted to The M.S.University of Baroda.
- Patel, S. J. and N. Y. Bhatt (1999) Alingment of larger foraminifers in thalassinoides burrows of the paleogene rocks of Western Kachchh, Gujarat. Bull Ind Geol. Asso, V- 32, pp.27-31.
- Patel, S. J. and Desai, B. G. (2001) The republic day Kachchh Earthquake of 2001: Trauma in *Oratosquilla striata* Jour. Geol. Soc. Ind. V-58, pp. 215-216.
- Patel, S.J. and Desai, B. G. (1999) Animal- sediment relationship in a modern tidal flats environment on Mundra coast, Gulf of Kachchh. Gondwana. Geol Mag., Sp. V-4, pp. 315-320.
- Patel, S.J and Shringarpure, D.M. (1989a) Trace fossil in Lower Miocene carbonate rocks of Ramwada Mandir, Western Kutch, Gujarat State. Mem. Jour. Geol. Soc. Ind.
- Patel, S.J. and Shringarpure, D.M. (1989b) Examples of shell predation by predatory gastropods and parasites in the Middle Eocene rocks of Kutch, Gujarat., Curr. Sci., V-58, pp. 1023- 1024.
- Patel, S.J. and Shringarpure, D.M. (1990). Oligocene-Miocene stages and trace fossils characteristics in Western Kutch, Gujarat. Jour. Geol. Soc. India, Mem.16, p. 96-109.
- Patel, S.J. and Shringarpure, D.M. (1996). Comparative study of the microfauna and trace fossils contents in the Paleogene rocks of Western Kutch , J.M.S.Uni. Baroda, V 52-53 p 1-14
- Patel, S.J. and Shringarpure, D.M. (1998). The intensively bored rocks in Vinjhan (Kutch) indicate Glossifungites ichnofacies Jour. Geol. Soc. Ind., V- 51, pp.87-92.
- Patel, S.J. and Shringarpure, D.M.(1992). Trace fossil *Limulicubichnus* from the Lower Miocene of Kutch. Curr.Sci. V-63, pp.682-984.
- Patel, S. J.; Desai, B. G. and Bhatt, N.Y., (2001) Neotectonic evolution of the coastal landforms between Jakhau and Mundra, Gulf of Kachchh, Western India. Bull. Ind. Geol. Assoc. (In Press)
- Patel, S. J.; Desai, B. G. and Bhatt, N.Y., (2002) Origin of Air Trap structures in Beach-bar complex and their significance- A study from Mandvi Coast, Gulf of Kachchh, Western India. Jour. Geol. Soc. Ind. V-60, 391-400.
- Pearson, T.H & Rossenbureg, R. (1978): Macrobenthic succession in relation to organic enrichment and pollution of the Marine environment-Oceanography, Marine Biology : An Annual Review, 16: 229-311; UCL press.
- Pemberton, S.G. and Frey, R.W. (1982) Trace fossil nomenclature and the Planolites-Paleophycus dilemma. Jour. Paleon. V-56, pp. 843-881.

- Pemberton, S.G. & Frey, R.W. (1985): The *Glossifungites* ichnofacies: Modern examples from the Georgia coast, U.S.A - In Curran, H.A. (ed) Biogenic structures; Their use in interpreting depositional environments, Society of Economic Geologist, paleontologic and mineralogist, Special publication, 35:237-259
- Pemberton, S.G.; MacEachern, J.A.; Frey, R.W. (1992) Trace fossils facies model: environmental and allostratigraphic significance, in Facies Models- Response to sea level change (Eds R.G.Walker and N.P. James), Geological Association of Canada, Ottawa, pp 47-72.
- Pemberton, S.G.; Spila, M.; Pulham, A.J; Saunders, T; MacEachern, J.A.; Robbins, D; and Sinclair, I.K. (2001) Ichnology and sedimentology of shallow marginal marine systems. Short course notes, V-15; Geol. Assoc. Cana. 343p.
- Petersen, C.G.J (1915) On the animal communities of the sea bottom in the skagerak, the Christiania Fjord and the danish Waters, Rep. Dan. Biol. Stn. V23, 3-28.
- Pianka, E.R. (1970) On *r*- and *K*-selection. *Amm. Naturalist*, 104, 592-597.
- Pickerill, R.K. (1990) Nonmarine *Paleodictyon* from the carboniferous Albert Formation of Southern New Brunswick. *Atlantic Geol.* V-26; 157-163
- Pillai, N.K. (1951): Decapod (Brachyura) from Travancore. *Bull. Central. Res. Inst. Univ. of Travancore*, ii, no 1, Ser. C, pp 1-46.
- Platt, L.B. (1962) The Rann of Kutch., *J. Sedi Petrol.*, V-32, pp. 92-98.
- Podar, M.C. (1963) Geology and Oil Possibilites in Kutch in Western India, *Proc. 2nd. Symp. Dev. Pet. Res. Asia and Far East*, No. 18, pp. 226-230.
- Pryor, W.A. (1975) Biogenic sedimentation and alteration of the argillaceous sediments in shallow marine environments. *Bull. Geol. Soc. Amm.*, V-86:1244-1254.
- Pollard, J.E. Goldring, R., Buck, S.G. (1993) Ichnofanbric containing *Ophiomorpha* : Significance in shallow-water facies interpretation. *Jour. Geol. Soc. Lon.* V-150, pp149-164.
- Posey, M.H. (1986) Changes in a benthic community associated with dense beds of a burrowing deposit feeder, *Callianassa californiensis*. *Mar. Ecol. Prog. Ser.* V-31, pp 15-22.
- Powell, E.H.Jr and Gunter, G (1968) Observation on the stone crab *Menippe mercenaria* Say, in the vicinity of Port Aransas, Texas, *Gulf Res. Rep.* V-2, pp 282-299.
- Purdy, E.G. (1964) Sediments as substrates, In Imbrie J. and Newell N. (eds) *Approaches to paleoecology*, John Willey, New York. pp 238-271.
- Rabinowitz, P.D., Coffin, M.F., and Falvey, D., (1983) The seperation of Madagascar and Africa. *Science.* V-220, pp-67-69.

- Rajnath, (1932) A contribution to the stratigraphy of Cutch. Quart. Jour. Geol. Min. Met. Soc. India, V- 4, pp.161-174.
- Rajnath (1942) The Jurassic rocks of Cutch- Their bearing on some problems of Indian Geology. (Presidential Address, Geol. Soc.) Proc. 29th Ind. Sci.Cong.
- Raju, D.S.N (1969) Eocene-Oligocene boundary in Ramania area, Kutch, ONGC. Bull. V-6, 71.
- Raju, D.S.N (1971) Observations on the Eocene, Oligocene, and Miocene foraminiferal biostratigraphy of Kutch, Western India; Publ. Centre Adv. Study Geol., Punjab Univ., Chandigarh, Vol-10., pp 136-155.
- Rao, D.G. (1988) A shallow seismic reflection study of the Gulf of Kachchh- Observation on its structural evolution. Mar. Geol. V-82, pp 277-283.
- Reade, T.M. (1884). Miniature Domes in sand. Geol. Mag., v. 21, pp.20-22.
- Reineck, H.E and Singh, I.B. (1973) Depositional sedimentary environment, 439 p Springer Verlag, Berlin.
- Reineck, H.E And Singh, I.B. (1980). Depositional sedimentary environment, with reference to terrigenous clastics, pp. 66-67. Springer Verlag, Berlin.
- Reman, A.(1933) Wiss Meeresunters.Abt.Kiel,N.F., V-21,pp-161-221.
- Rhoads, D.C. and Young,D.K. (1970) The influence of the deposit feeding organism on sediment stability and community trophic structure Jour. Mar. Res. V-28, pp 150-178.
- Rhoads,D.C. (1967) Biogenic reworking of intertidal and subtidal sediments in Barnstable Harbour and Buzzards bay, Masschusetts., Jour. Geol, 75; 461-476.
- Richter, R. (1924) Flachseebeobachtungen zur palaontologie und geologie, VII-XII: Senkenbergiana, V-26; 119-165.
- Richter, R. (1926) Flachseebeobachtungen zur Palaontologie und Geologie XXI. Bau, Begriff und Palaogeographische Bedeutung von Corophioides luniformis (Blankenhorn, 1971). Senkenbergiana, V-8; 200-291.
- Richter, R. (1936) Marken und Spuren im Hunsruckschiefer. I. GeflieB-Marken. Senkenber. V-18, 215-244.
- Richter, R. (1937) Marken und Spuren aus allen Zeiten. I-II. Senkenbergiana, V-19, 150-169
- Richter, R. (1952) Fluidal-Textur in sediment-Gesteinen und uber sedifluktion Uberhaupt. Notizbl. Hess. Land. Boden. V-4, 67-81.

- Rieth, A. (1932) Neue Funde spongiomorphen fucoiden aus dem jura schwabens- Geol. und. Paleon. Abhandl., Neue Folge, V-19; 257-294.
- Rindsberg, A. (1994) Ichnology of the Upper Mississippian Hartselle sandstone of Alabama, with notes on other Carboniferous Formations., Geol. Surv. Alabama, Bull 158; 107p.
- Rindsberg, A.K. (1990) Ichnologic consequences of the 1985 International Code of Zoological Nomenclature. Ichnos. V-1, pp 59-63.
- Root, R.B. (1967) The Neich exploitation pattern of the blue-grey gnacatcher. Ecol. Monogr. V-37; 317-350.
- Rouse, G.W. and Fauchald, K (1997) Cladistic and Polychaetes. Zool. Scripta. V-26 (2), 139-204.
- Roy, B and Merh, S.S. (1982) The Great rann of Kachchh: An intriguing quaternary terrain. Pp 100-108 In S.S.Merh (ed) Recent researches in Geology (9) Hindustan Publishing Corp. New Delhi.
- Roy, B. and Merh, S.S. (1977) Geomorphology of the Rann of Kutch and Climatic changes. In Agrawal, D.P. and Pande, B.M. (eds) Ecology and Archaeology of Western India, Concept publishing, New Delhi, pp 195-200.
- Samanta, B.K. (1970) Middle Eocene Planktonic foraminifera from Lakhpat, Kutch, Western India. Micropaleont. V-16, (2), 185-215.
- Sanders, H.L. (1958) Benthic studies in Buzzards bay. I Animal sediment relationships, Limnol. Oceanogr. V-3, pp245-258.
- Sanders, H.L. (1968) Marine benthic diversity: A comparative study. Am.Nat. V-102, pp 243-282.
- Saporta, G. De (1887) Nouveaux documents relatifs aux organismes problematique des anciennes mers: Soc Geol France, Bull., Ser. 3, V-15, p-286-302.
- Saporta, G.De. (1884) Les organismes problematiques des Anciennes Mers. Masson, Paris, 102 p.
- Saraswati, P.K. and Banerji, R.K (1984b) Post Trappean sedimentation History of the North-Western Kutch, Proc. Xth, I. C.M.S. pp. 377-390.
- Saraswati. P.K and Banerji, R.K. (1984a) Lithostratigraphic classification of the Tertiary equence of North-Western Kutch, Proc. Xth, I. C. M. S. pp. 369-374.
- Sarjeant, W. A. S. and Kennedy, W.J. (1973) Propasal for a code for the nomenclure of trace fossils. Can. Jour. Ear. Sci. V-10, pp 460-475.
- Sarjent, W.A..S. (1979) Code for trace fossil nomenclature. Paleo-3, V-28, pp. 147-166.

- Savazzi, E. (1982) Burrowing habits and cuticular sculptures in recent sand-dwelling brachyurans decapods from the Northern Adriatic Sea (Mediterranean). *Neus. Jahrbuch. Fur. Geol.und.Paleontologie*, V-163, pp. 369-388.
- Schafer, W (1956) Wirkungen der Benthos-Organismen auf den Jungen Schichtverband. *Senckenbergiana lethaea*, V-38; 183-263.
- Schafer, W. (1972) Ecology and paleoecology of marine environments (G.Y. Craig. Ed-I Oertel transl.), University of Chicago press, Chicago.
- Schaffner, L.C. (1990) Small scale organism distributions and patterns of species diversity: evidence for positive interactions in a benthic community. *Mar. Ecol. Prog. Ser.* V-61; 107-117.
- Schaffner, L.C.; Dellapenna, T.M.; Hinchey, E.K.; Friedrichs, C.T.; Neubauer, M.T.; Smith, M.E; and Kuehl, S.A. (2001) Physical Energy regimes, sea bed dynamics, and Organism-sediment interactions along a Estuarine gradient. In J.Y. Aller; S.A.Woodin and R.C.Aller (eds) *Organism-sediment interactions*, Univ.South. Carolina. Press. 159-179.
- Seilacher, A.(1953) Studien Zur Paläozoologie. I Über die Methoden der Paläozoologie. *Neus Jahrbuch fur Geologie und paläozoologie. Abhandlungen*, 96: 421-425.
- Seilacher, A. (1964) Biogenic sedimentary structures. In *Approaches to paleoecology*. J.Imbrine and Newell, N. (Eds), pp 296-316, New York: Wiley.
- Seilacher, A. (1967) Bathymetry of trace fossils *Marine Geology*, V-5, 413-428.
- Seilacher, A. (1992) Quo vadis, Ichnology? In Maples, C.G. and West, R.R. (eds) *Trace fossils. Short course in Paleontology*, No-5, pp 224-238.
- Sengupta, B.K (1964) Tertiary biostratigraphy of part of Northwestern Kutch. *Jour. Geol. Soc. Ind.*, Vol- 5., pp-138-158.
- Shringarpure, D.M (1984) Mesozoic of Kutch (India): Middle Jurassic to Lower Cretaceous depositional environment as revealed by biosedimentary structures. In Reif, W.E. and Westphal (eds) *3rd symposium on Mesozoic Terrestrial ecosystem*, Tubingen, pp-227-229.
- Shringarpure, D.M (1985) Ichnological studies of the rocks exposed between Chitrod and Manfra, Eastern Kutch-Gujarat state., Unpubl., PhD thesis, M.S.University of Baroda.
- Shringarpure, D.M (1986) Trace Fossils at Omission surface from Mesozoic of Kutch, Gujarat, Western India. *Bull. Geol. Min. Met. Soc. India.*, Vol-54., pp131-148.
- Shringarpure, D.M; Desai, G and Merh, S.S. (1976) Stratigraphic leakage: A problem with the foraminifera assemblages of the Wager Mesozoic sediments of eastern Kutch, *Proc. VIth ind. Col. Micop. & strati*, 245-268, varanasi.

- Shrivastava, P.K. (1971) Recent sediments from Rann of Kutch. Jour. Geol.Soc. Ind. V12, pp 392-395.
- Singh, C.P.S., Jaitly, A.K., and Pandey, D.K. (1982). First Report of some Bajocian-Bathonian (Middle Jurassic ammonoids and the age of the oldest sediments from Kachchh, India. Newsletters on stratigraphy. V-11, pp 37-40, Berlin.
- Singh, I.B. (1978) Microfacies, Petrology and Mineralogy of the Tertiary rocks of Guarnala near Narayan Sarovar, Kutch, India and their paleontologic significance. Jour.Pal. Soc. Ind. V-21,22; pp 78-95.
- Singh, I.B. (1980) The Bijaigarh Shale, Vindhyan System (Precambrian), India- An example of a Lagoonal deposit. Sedimentary Geol. V-25; 83-103.
- Sivewright, R (1907) Cutch and The Ran. The Geographical J. XXIX 518-539. V-5, pp. 168-174.
- Skoog, S.Y.; Venn. C.; Simpson, E.L. (1994) Distribution of *Diopatra cuprea* across Modern Tidal Flats: Implications for *Skolithos*., Palaios, V-9 188-201.
- Snelgrove, P.L. and Butman, C.A. (1994) Animal-sediment relationships revisited: Causes versus effect. Oceano. Mar. Biol. Ann. Rev. 32: 111-177.
- Sohoni, P.S. & Malik, J.N. & Merh, S.S. & Karanth, R.V. (1999): Active tectonic astride Kachchh zone, Kachchh, western India- Journal Geological Society of India., 53:579-586; Bangalore.
- Spat, L.F. (1927-'33) A revision of the Cephalopod Fauna of Kutch (Cutch). Pal. Ind. V-9, 945 p.
- Srivastava, P.S. and John, V.C. (1977) Current regime in the Gulf of Kachchh. Ind. J. Mar. Sci. V-6, pp39-48.
- Stapor, F.W. 1975. Holocene beach ridge plain development, north-west Florida. Z.Geomorphol., N.F. Suppl-Bd., 22:116-144.
- Steiglitz, R.D. And Inden, R.F. (1969). Development of Cavernous sediment in a non beach environment: Journal of Sedimentary Petrology, v.39, pp.342-344.
- Stewart, R.A and Plikey, O.H. (1966) Sediments of the Northern Arabian Sea. Tech Rep., U.S.Nav. Oceanogr.off. pp. 186.
- Strios, S.C. & Laborel, J. & Laborel-Deguen, F & Papageorgious, S. & Evin, J. & Pirazzoli, P.A. (2000): Seismic coastal uplift in a region of subsidence: Holocene raised shorelines of Samos Island, Aegean Sea, Greece-Marine Geology, 170:41-58; Elsevier publication, Amsterdam.
- Sverdrup, H.U; Johnson, M.W.; Fleming, R.H. (1942) The Ocean. 1087pp, Prentice-Hall, New York.

- Swinbanks, D.D. and Murray, J.W. (1981) Biosedimentologic zonation of Boundary Bay tidal flats, Fraser River Delta, British Columbia, *Sedimentology*, V-28, 201-237.
- Taghon, G.L. and Greene, R.R (1992) Utilisation of deposited and suspended particulate matter by benthic interface feeder *Limnol. Oceanogr.* V-37, 1370-1391.
- Taghon, G.L.; Nowell, A.R.M; and Jumars, P.A. (1980) Induction of suspension feeding in spionid polychaetes by high particulates fluxes, *Science*. V-210; 563-564.s
- Taghon, G.L.; Nowell, A.R.M; and Jumars, P.A. (1984) Predicting particle selection by deposit feeders: a model and its implications. *Limnol and Oceanogr.*23; 752-759
- Talwani, P. (1989): Characteristic features of intraplate earthquakes and Models proposed to explain them- In Gregersen, S & Basham, P.W. (eds) *Earthquakes at North-Atlantic passive margins: Neotectonics and postglacial rebound*, 563-579.
- Taylor, A.M. and Goldring, R. (1993) Description and analysis of bioturbation and ichnofabric. *Jour. Geol. Soc. Lon.* V-150, pp 141-148.
- Terwindt, J.H.J and Brouwer, M.J.N (1986) The behaviour of intertidal sand waves during neap-spring tide cycles and the relevance for paleoflow reconstructions. *Sedimentology*, V-33; 1-31.
- Tewari, B.S (1952) The tertiary beds of Vinjan-Miani area, SW Kutch, India. *Curr Sci.* V-21, 217-218.
- Tewari, B.S (1957) Geology and Stratigraphy of the area between Waghopadar and Cheropadi, Kutch, India. *Jour. Pal. Ind. Soc.*, V-4, 6-11.
- Tewari, B.S and Singh, G. (1969) Stratigraphy of Tertiary beds of Dehdapur-Fulra-Phanandro area., N- W Kutch., *Bull. Ind. Geol. Asso.* V-2
- Tewari, B.S; Bharghav, O.S.; Khanna, S.N. (1964) Kutch microfauna, Middle Eocene foraminifera from Waghopadar, South-western kutch, *Jour. Pal. Soc. Ind.* 5-11; 77-82.
- Thakkar, M.G & Maurya, D.M & Raj, R.R & Chamyal, L.S. (1999): Quaternary tectonic history and terrain evolution of the area around Bhuj, Mainland Kachchh- *Journal Geological society of India*, 53:601-610; Bangalore.
- Thayer, C.W. (1975) Sediment-Mediated Biological Disturbances and the evolution of Marine Benthos. In M.J.S. Tevesz and McCall, P.L. (ed) *Biotic interactions in recent and fossil benthic communities*, V-3; 480-626. Plenum press.
- Thorson, G. (1957) Bottom communities (Sublittoral or shallow shelf), p. 461-534. In J.W. Hedgpeth (ed) *Treatise on marine ecology and paleoecology*. V-1, *Geol. Soc. Am. Mem.* 67.

- Tondon, K.K. (1976) Biostratigraphy of the Middle Eocene Rocks of a part of SW Kutch, India. *J. Pal. Soc. Ind.* V-19., pp 71-88.
- Torell, O.M. (1870) *Petrificata suecana Formations Cambricae*: Lunds Univ. Arskr, V-6; 1-14.
- Trask, P.D. (1939) Organic content of recent marine sediments. In Trask P.D. (ed) *Recent marine sediments*, 428-453 *Amm Assoc. Petrol. Geol.*.
- Trivedy, R.K. and Goel, P.K. (1984) *Chemical and Biological methods for Water pollution studies*. Environmental publication. 215p.
- Trueman, E.R. (1968) Burrowing habit and early evolution of the body cavities. *Nature*, V-218, pp96-98.
- Trueman, E.R. and Ansel, A. D (1969) The Mechanisms of burrowing in to soft substrata by marine animals. *Oce. Mar. Biol. Ann. Rev.* V-7, pp 315-366.
- Tweedie, M.W.F: 1950, Notes on grapsoid crabs from Raffles Museum. *Bull. Raffles. Mus.* 23. 310-313.
- Uchman, A.F. (1995) Taxonomy and Paleoecology of Flysch Trace fossils: the Marnoso-arenacea Formation and associated facies (Miocene, Northern Apennines, Italy) *Beringeria*, 1-115.
- Van Straaten, L.M.J.U. (1954) Sedimentology of recent tidal flat deposits and the Psammites du Condroz (Devonian). *Geol. Mijbouw.*, V-16; 25-47.
- Vaugelas, J.De. (1991) Determination et abondance des peuplement de crustacés décapodes thalassinides fouisseurs (*Upogebia* et *Callinassa*) de l'archipel des Lavezzi (Corse). *Travaux scientifiques du Parc Naturel Régional et des Réserves Naturelles de Corse*
- Vermeij, G.J. (1978) *Biogeography and Adaptation: Pattern of marine life*, Harvard University press, Cambridge, Mass.
- Vialov, O.S. (1969) *Vintobraznyy Khod chlenistonogogo iz kryma: paleont. Sbornik.*, Izdatel, Lvov. Univ. V-1, 105-109
- Vialov, O.S. (1971) *Redkie problematiki iz mesozoya pamira i Kavkaza paleont. Sbornik.*, Izdatel, Lvov. Univ V-7, 85-93.
- Vialov, O.S. (1972) The classification of fossils traces of life, 24th inter. Geol Congr. Sec.7, p-639-644.
- Viltisheva, I.F and Karzinkin, G.S. (1970), The use of radioactive carbon ^{14}C in studying the feeding of bottom invertebrates, *Publications of the All-Union Scientific Research Institute of Marine Fisheries and Oceanography (VNIRO)*, 69: 85-95.

- Voigt, E. (1965) Über parasitische polychaeten in kreide-Austern Sowie einige andere in Muschleschalen bohernde Wurmer: Palaon. Zeitsch. V-39, 193-211.
- Voigt, E. (1970) Endolithische Wurm-tunnelbauten (*lapispecus cuniculus* n.g. n. Sp. Und *Dodecaceria*[?] sp.) in Brandungsgerollen der oberen Kreide im nordlichen Harzzorland: gGeol Runds, V-60, 355-380.
- Voigt, E. (1972) Tonrollen als potentielle pseudofossilien: Natur U. Museum, V-102; 401-410.
- Voigt, E. (1975) Tunnelbaue rezenter und fossiler Phoronidea. Paleon. Zeitsc. V-49; 135-147.
- Vredenburg, E.W (1910). A Sumary of Geology of India, IInd Edition.
- Wagen, W. (1871-'73). Jurassic Fauna of Cutch. The Cephalopod. Pal. Indica, (IX), I (I-IV), 247pp.
- Wagle, B.G. (1979) Geomorphology of the Gulf of Kachchh Ind. Mar. Sci. V-8, pp 123-136.
- Wagstaffe and Fiddler The preservation of the Natural history specimen V-1, published by Witherby LTD, London.
- Warne, J. (1975) Traces and significance of marine rock borers. In T.P. Crimes, J.G. Harper (eds.) Trace fossils, Jour. Geol Soc. V-3, 515-526.
- Walker, K.R. and Bambach, R.K. (1974) Feeding by benthic invertebrates: classification and terminologies for paleoecological analysis, Lethaia, V-7; 67-78.
- Warner.G.F: (1977) The Biology of Crabs, Elek Science, London. pp. 1-202.
- Weimar, R.J and Hoyt, J.H (1964) Burrows of *Callianassa major* Say, Geologic indicators of littoral and shallow neretic environments, Jour. Paleon, V-38, 761-767.
- Westergard, A.H. (1931) Diplocraterion, Monocraterion and Scolithus from the Lower Cambrian of Sweden: Sver. Geol. Undersok. Ser. C. Avh, Och, Upps, No 372, 25p 10pls.
- Wetzel, A. (1983) Biogenic structures in a modern upwelling region: Northwest African continental margin, in Coastal Upwelling and its sediment record: Part- B, Sedimentary Records of Ancient Coastal Upwelling (Eds. J. Thiede and E. Suess), Plenum press, New York, pp. 123-144.
- Wieser, W. (1959) The effect of grain size on the distribution of small invertebrates inhabiting the beaches of Puget Sound. Limnol. Ocenogr. V-4, 181-194.
- Williams, A.B. (1965) Marine decapod crustceans of the Carolinas. United States Bureau of Commercial Fisheries, Fish. Bull. V-65, 298p.

- Wilson, W.H.Jr. (1990) Competition and predation in marine soft sediment communities. *Ann. Rev. Ecol. Sys.* V-21, 221-241.
- Woddin, S.A. and Jackson, J.B.C (1979) Interphyletic competition among marine benthos, *Am. Zool.* V-19, pp 1029-1043.
- Woodin, S. A. (1974) Polychaete abundance patterns in a marine soft-sediment environment: the importance of biological interactions. *Ecol. Monogr.* V-44, pp 171-187.
- Woodin, S. A. (1982) Browsing: important in marine sedimentary environments? Spionid polychaete examples. *Jour. Exp. Mar. Biol. Ecol.* V-60, pp 35-45.
- Woodin, S. A. (1983) Biotic interaction in recent marine sedimentary environments. In Tevesz, M.J.S. and McCall, P.L (eds), *Biotic interaction in recent and fossil benthic communities* Plenum press, New York, pp 3-38
- Wright, L.D.; Schaffner, L.C.; and Maa, J.P.Y.; (1997) Biological mediation of bottom boundary layer process and sediment suspension in the lower Chesapeake bay. *Mar. Geol.* V-14; 27- 50.
- Wynne, A.B. (1972) *Memoirs on the Geology of Kutch to accompany the Map compiled by A. B. Wynne and F.Fedden during the seasons of 1867-68 and 1868-69.* *Mem.Geol. Surv. Ind.* V-9, 289 p.
- Zobell, C.E. and Anderson, D.Q. (1936) Vertical distribution of bacteria in marine sediments. *Amm. Petrol. Geol. Bull.* V-20, 258-269.
- Zaigham, Z.G and Malick, K (2000) Prospects of hydrocarbon association with fossil rift structures of the south Indus basin, Pakistan. *AAPG. Bull.* V-84(11), 1833-1843.