List of Figures

Figure No.

Page No.

1.1	(a) Location map of the study area.(b) Major crustal terranes of NW India;(modified after Sinha-Roy, 2000).	2
1.2	Shuttle Radar Topography Mission (SRTM) derived shaded relief map of the Champaner region. Source: USGS Earth-explorer.	4
1.3	Communication map of the study area.	5
2.1	Generalised geological map of Aravalli Mountain belt; (modified after Gupta et al., 1992).	8
2.2	Lithostratigraphic map of Southern Aravalli Mountain belt; (modified after Gupta et al., 1992).Magnetic foliation data presented within Godhra granite is after Mamtani, (2014).	10
3.1	Geological Map of the Study area. Modified after, Jambusaria, 1970; Gupta et. al., 1997.	36
3.2	Field photograph showing	39
	 (a; b) Second order F₁ folds of tight isoclinals variety in micaceous quartzites, loc. E of Jhand; (c) N-S trending meso-scale open warps in grey-quartzites, loc. S of Mota-Raska; (d)Parallel relationship between primary bedding and foliation in vertical quarztites, loc. N of Narukot (dome fringe); (e) Asymmetrical ripple marks and (f) Tabular cross-bedding in quartzites, (e) loc. E of Borkas, (f) loc. W of Wadek. 	
3.3	Field photograph showing:	41
	 (a) Steep F₃ fold axes in quartzites, loc. S of Wadek; (b) S-C fabric defining parallelogram sigmoids in quartzites, loc. S of Borkas; (c) Second order tight F₁ folds and first order open F₂ folds in quartzite-phyllite intercalations, loc. E of Shivrajpur; (d) Mined F₂ core for Mn mineralization loc. W of Bhat; (e) First order higher wavelength F₂ open folds in quartzites, loc. Vadatalav (Shivrajpur to Goghamba bypass); (f) Asymmetrical ripples along limbs of F₂ folds, loc. Vadatalay (Shivrajpur to Goghamba bypass) 	
	ioe. , adamaa (om viajpar to oognamoa oypass).	

3.4	Field photograph showing:	43
	 (a) Meso-scale normal fault in quartzites; (b) Fault gauge development in quartzites (feature seen at top left corner of the photograph); (c) Goethite on upper surface of quartzites; (a; b; c) loc. Vadatalav (Shivrajpur to Goghamba bypass); (d) Planar cross-bedding in quartzites, loc. Koba; (e) Fault breccia development along ENE-WSW trending shear zone, loc. Rustampura; (f) Quartz rods preserved in quartzites, loc. Chethapur; (g; h) Massive to Fragile slaty outcrops covered under thick soil and vegetation, loc. S of Rajgarh. 	
3.5	Field photograph showing:	44
	 (a) Thin band of calc-silicate rock within massive phyllites, loc. Quarry section near Boriya; (b) Interlayer tight folds of shorter wavelength within phyllites, loc W of Navagam; (c; d) F₂ folds preserved in phyllites, loc. Kakalpur; (e) Oblique relationship between secondary S₂ cleavages and primary bedding S₀IIS₁ loc. Bhat; (f) Sigmoid development in phyllites representing top-to-east sense of ductile shear, loc. E of Jaban. 	
3.6	Field photograph showing:	48
	 (a) Parallel laminations preserved in phyllites, loc.Kuivav; (b) L₁ intersection lineations preserved in hornfels, loc S of Kuivav; (c) Chaistolite- Andalusite mineral development over phyllites, loc. Wadek; (d) Protruding cordierite porphyroblast within phyllites, loc. Zarav; (e) Phi shaped porphyroclasts within sheared schists, loc. E of Dhanpur; (f) L₂ lineations preserved within carbonaceous schists, loc. N of Narukot (within dome); (g) Elephant skin weathering preserved in grey dolomitic limestones, loc. NE of Gandhra; (h) Unoriented actinolite and tremolite needles in calc-silicate rock, loc. Vav. 	
3.7	Field photograph from Jothwad showing:	49
	 (a) Enclaves of calc-silicate rock in granite; (b) Caught up folded fragment of calc-silicate rock embedded in granite; (c) Inferred contact between weathered calc-silicate rock and granite at the side face of 5 m deep pit; 	

	(e;d) Tight to isoclinal folds in calc-silicate rock;(f) Mushroom-shaped outcrop geometry in calc-silicate rock.	
3.8	Field photograph showing:	51
	 (a) Calc-silicate Skarn found in vicinity to granite, loc. Jothwad; (b) ESE-WNW trending meso-scale tight F₂ folds within dolomitic limstone, loc E of Sarasuva; (c) Outcrop of massive crystalline limestone an extension of upper dolomitic limestones, loc. W of Shivrajpur; (d) Oligomict meta-conglomerates having light to dark grey colour quartzitic clasts, loc. E of Lambhia; (e) Polymict meta-conglomertes having rounded to sub-rounded clasts of varied assemblages, loc. Jaban; (f) Tensional gashes within polymict meta-conglomerates loc. E of Laban 	
3.9	Field photograph showing:	55
	 (a) Intercalations of meta-subgreywackes and meta-conglomerates, loc. W of Jaban; (b) Panoramic view of granite country, loc N of Dhanpur; (c) Leucocratic equigranular fine grained grey granite, loc. Ranjitnagar; (d) Contact between coarse grained pink granite and fine grained grey granite, loc. SE of Valothi; (e) Intrusive contact between the Champaner meta-sediments and granite, loc. E of Dhanpur; (f) Porphyritic variety of pink granite having oriented feldspar phenocrysts, loc. Singpur. 	
3.10	Field photograph showing:	56
	 (a) Feldspar segregation observed in the coarse grained pink granite, loc. Valothi; (b) Equigranular non-porphyritic pink granite, loc. S of Kevra; Xenoliths of (c) Schists; (d) Phyllites; (e) Calc-silicate rock; (f) Fine grained grey granite within granites of the study area, loc. (c) E of Dhanpur;(d) Ranjitnagar;(e) Zab;(f) Ranjitnagar; (g) Oriented mineral segregations within granites, loc. Kanpur; (h) Orientation of minerals within foliated granites, loc. E of Lambhia. 	
4.1	Landsat data of the Champaner terrain (Source: USGS-Earth Explorer)	58
4.2	Structural Map of the Study area.	60
	Modified after Jambusaria, 1970; Gupta et. al., 1997.	

4.3	Structural Map of Sector 1	61
4.4	Structural analysis of field data from Domain I and Domain II	63
4.5	Structural analysis of field data from Domain IIIa and Domain IIIb	64
4.6	Structural analysis of field data from Domain IVa	66
4.7	Structural analysis of field data from Domain IVb	67
4.8	Structural analysis of field data from Domain V	68
4.9	Structural analysis of field data from Domain VI	70
4.10	Structural Map of Sector 2.	72
4.11	Structural analysis of field data from Domain VII	73
4.12	Structural analysis of field data from Domain VIII	74
4.13	Structural analysis of field data from Domain IX	76
4.14	Structural analysis of field data from Domain X	77
4.15	Structural analysis of field data from Domain XI	77
4.16	Structural analysis of field data from Domain XII	77
4.17	Structural analysis of field data from Domain XIII	78
4.18	Structural Map of Sector 3.	80
4.19	Structural analysis of field data from Domain XIV	81
4.20	Structural analysis of field data from Domain XV	82
4.21	Structural analysis of field data from Domain XVI	83
4.22	SRTM derived shaded relief map of the Narukot Dome.	83
	Respective orientations of the axial planes and	
	fold axis has been demarcated.	
4.23	Structural Map of Sector 4.	85
4.24	Structural analysis of field data from Sector 4	86
4.25	Structural Map of Sector 5.	88
4.26	Structural analysis of field data from Sector 5	88
4.27	Sketch depicting dextral fault in top view, where as top-to-east reverse shear in cross-sectional view. The sketch signifies the transfer of faults from strike slip to reverse in single fault episode.	90
4.28	AMS orientation data plotted as lower hemisphere projections for the study area. Dotted great circles represent the magnetic foliation, whereas continuous great circles represent the S_0 reading recorded from	97

	the field. Filled square, triangle and circle indicate orientations of K1, K2 and K3 respectively.	
4.29	Jelinek plots and Flinn diagram for the quartzites of the study area.	98
5.1	Photomicrograph of quartzite showing:(a) Granoblastic polygonal texture with inequigranular interlobate boundary(b) Granoblastic polygonal texture with seriate interlobate boundary	102
5.2	(a) Schematic sketch showing stages of Bulging recrystallization(BLG), after Passchier and Trouw, 2005; P.42.(b) Photomicrograph of quartzite of study area showing BLG recrystallization.	103
5.3	(a)Schematic sketch showing stages of Sub-grain rotation recrystallization (SGR), after Passchier and Trouw, 2005; P.42.(b) Photomicrograph of quartzite of the study area showing SGR recrystallization.	104
5.4	(a) Schematic sketch showing stages of Grain boundary recrystallization(GBM) after Passchier and Trouw, 2005; P.42.(b) Photomicrograph of quartzite showing GBM recrystallization with left over grains.	105
5.5	 (a) Schematic sketch showing development of Grain Boundary Area Reduction recrystallization (GBAR), after Passchier and Trouw, 2005; P.51. (b) Photomicrograph of quartzite showing GBAR recrystallization, the present feature show anomalous undulose extinction. 	105
5.6	Back-scattered electron image of ilmenite,	108
5.7	 (a) Two varieties of shear band cleavages, after Passchier and Trouw, 2005; p.128; Photomicrographs in XPL of phyllite-mylonite showing (b) anastomosing C-type shear bands; (c) compressional crenulations getting sheared to form extensional crenulations; (d-f) M-domain and Q-domain within the phyllite-mylonite. 	111
5.8	Classification of mantled porphyroclasts having dextral shear senses (after, Passchier and Trouw, 2005; p. 133).	112
5.9	Photomicrographs in XPL with lined sketch of phyllite-mylonite showing (a,b) "Oval shaped" φ-type mantled porphyroclasts with tapering trails at same elevation on both sides; (c,d) "Fish head shaped" φ-type mantled porphyroclasts with swallowed	114

	quartz grain; (e,f) "Amoeboid shaped" φ-type mantled porphyroclasts having irregular boundary.	
5.10	 Photomicrographs in XPL/ PPL with lined sketch of phyllite-mylonite showing (a,b) "Oblong shaped" φ-type mantled porphyroclasts with tapering trails. The upper half of the photomicrograph display prominent S-C fabric; (c,d) "Hatched circular shaped" φ-type mantled porphyroclasts with strain shadows. (e,f) σ-type mantled porphyroclasts of stair stepping variety. 	115
5.11	Photomicrographs in XPL/ PPL with lined sketch of phyllite-mylonite	116
	Showing	110
5 1 2	(a-f) σ type mantled porphyroclasts of stair stepping variety.	117
5.12	Showing	11/
	(a,b) δ -type mantled porphyroclasts of stair stepping variety.	
5.13	Morphology of mica fish, groups 1-6 (after, ten Grotenhuis et al., 2003)	118
5.14	 Photomicrographs in PPL with lined sketch of phyllite-mylonite Showing (a,b) Group 1 opaque fish (R= 2.65; α = 420); (c,d) Group 2 opaque fish (R = 4.33; α = 450); (e,f) Group 3 opaque fish (R = 2.66; α = 220). 	119
5.15	Photomicrographs in PPL/XPL with lined sketch of phyllite-mylonite	120
	Showing, (a,b) Group 4 opaque fish (R= 4.5; α = 420); (c,d) Group 5 opaque fish (R = 2; α = 310); (e,f) Group 6 opaque fish (R = 14; α = 120).	
5.16	Photomicrographs in PPL/XPL with lined sketch of phyllite-mylonite showing oblong shaped opaque porphyroclasts without mantles having top-to-east rotation.	122
5.17	(a) Photomicrograph in XPL with(b) lined sketch of phyllite-mylonite showing pinch and swell microstructure in opaque.	123
5.18	 (a) Photomicrograph in XPL of andalusite hornfels showing presence of quartz and muscovite inclusions along with the recrystallized matrix. (b) Lined sketch of andalusite porphyroblast having Oppositely 	124

	Concave Microfolds (OCMs).	
5.19	Photomicrograph of	128
	 (a) coarse grained granite, loc. Mudhiyari (2.5X, CN); (b) coarse grained porphyritic granite, loc. Wadek (2.5X, CN); (c) fine grained granite, loc. Ranjitnagar (2.5X, CN); (d) granite showing poikilitic texture, orthoclase as oikocryst having presence of quartz chadacryst, loc. SE of Valothi (2.5X, CN); (e) granite showing micro-perthitic texture, loc. S of Wadek (4X, CN); (f) granite showing myrmekitic texture, loc. Dev Dungar, (2.5X, CN). 	
5.20	 Photomicrograph of (a) granite showing flame perthite, loc. Zab village (2.5X, CN); (b) granite showing myrmekitic texture, loc. Sukhi dam, (4X, CN); (c) calc-silicate rock xenolith showing hornfelsic texture, loc.SE of Valothi (2.5X, CN); (d) xenolith of schist showing schistose texture, loc. Sukhi Dam site (2.5X, CN); (e) xenolith of fine grained granite, loc. Ranjitnagar (2.5X, CN); (f) granite having undulose quartz, loc. Nani Khatva (4X, CN). 	129
5.21	 Photomicrograph of granite (a) having undulose biotite, loc.Nani Khatva (2.5X, CN); (b) having kinking in biotite, loc. Nani Khatva(2.5X, CN); (c) having micro faulting in plagioclase feldspar, loc.Kanpur (4X, CN); (d) having fractured quartz grains, loc. Wadek, (2.5X, CN); (e) having micro folds in mica minerals, loc. Lambhia (4X, CN); (f) having chess board extinction in quartz, loc. S of Wadek (2.5X, CN). 	130
5.22	Nine diagnostic Porphyroblast-matrix relationship based on internal $(S_i \sim dotted lines)$ and external $(S_e \sim continuous lines)$ schistosity planes. (after Zwart, 1962)	131
5.23	Schematic representation of pre, inter, syn and post-tectonic porphyroblast growth. The upper part of the diagram shows deformation resulting in a single foliation, while the lower part depicts deformation resulting in crenulation of an older foliation. (after, Passchier and Trouw, 1996)	132
5.24	Photomicrograph of (a) graphite schist from Narukot dome (2.5XCN) showing differentiated crenulations cleavage. S_2 is at high angle with respect to	133

	 S₀ sub parallel S₁; (b) Mn-rich phyllites from Bhat (10XCN) showing continuous crenulation cleavage. (c) phyllite near Borkas (10XPPL) showing discrete crenulation cleavage. (d) phyllite located north of Malabar (10XCN) depicting discontinuous crenulation cleavage. 	
5.25	 Photomicrograph of (a) phyllite located north of Malabar (4XPPL) showing syn-D₂ garnets. (b) andalusite hornfels from Wadek (40XCN) showing development of sillimanite (fibrolite) along andalusite quartz margins; (c) cordierite hornfels (2.5XCN) showing unmatched internal fabric (S_i) with external schistosity plane (S_e). 	135
6.1	Geo-eye image of the Champaner terrain; source: google-earth software. Oval structure along the E margin represents the Narukot Dome. Dotted line across the dome and further W shows locations of 1-32 stations for microtremor measurements.	139
6.2	H/V spectral frequency plot recorded for 1-32 stations from the study area.	140
6.3	Fundamental resonant frequency of 1 to 32 stations along WNW trending profile.	147
6.4	Layered model for the profile along Narukot Dome and to its W.	147
7.1	QAP diagram, after Streckeisen, 1974.	152
7.2	Harker Bivariate plots	153
7.3	REE Chondrite normalized plot (Boynton 1984) suggesting higher LREE content over HREE.	155
7.4	Harker bi-variate plots of SiO ₂ vs trace elements.	157
7.5	The tectonic discriminatory plot by Maniar and Piccoli (1989).	159
7.6	R1-R2 plot by Batchelor and Bowden, (1985) for tectonic discrimination of granites of study area.	160
7.7	Tectonic discrimination plots by Pearce et al., (1984),	161
7.8	 (a) ORG normalized spider plot for ideal syn-collision granites (after, Pearce et al., 1984); (b) ORG normalized spider plot for 10 samples of granites of Study area. 	162
7.9	Chondrite normalised pattern for granites of study area showing Ta	162

enrichment with adjacent Nb anomaly. Thompson, (1982) data set used for normalization purpose.

7.10	Zr-M plots for estimation of zircon saturation temperature.	163
8.1	 (a) Cartoon showing suturing of proto-continents and refolding of earlier folds along with the plunge on account of regional deformation as well as emplacement of syntectonic granite; (b) Closing up of earlier refolded folds orthogonally and development of sinistral/ dextral faults along the pre-existing axial planes due to post intrusive pulse. Granite of post-tectonic nature holds xenoliths of Champaner meta-sediments and earlier syn-tectonic granite. These cartoons have been modified after winter 2012. 	170