LIST OF FIGURES

| Figure No | es Pa | age No. |
|--------------|---|------------|
| 1.1 | Location Map (DEM) of the study area in southern Kachchh (Source ISRO & GSI) KA-Kotda-Roha Area; FA-Fakirwadi Area; GA-Gangeswar Area; VA-Varli Area. | 06 |
| 1.2 | Image-map showing geomorphology of the Great Ranns of Kachchh.(After Merh and Patel, 1988) | 09 |
| 1.3 | Geology and Tectonic Set up and Location Map of Study Area (After Kothyari et.al. 2016c) | 11 |
| | A-Anjar, B-Bhuj, Bh-Bhachau,D-Dholavira, De-Desaplar, F-Fatehgadh,G-Gedi, L-Lakhpat. | |
| | Faults: ABF-Allahbund Fault, GF-Gedi Fault, KHR-Katrol Hill Range,KMF-Kachchh Mainland Fault, IBF-Island Belt Fault, NKHF-North Katrol Hill Fault, NPF-Nagar Parkar Fault, SKHF-South KatrolHill Fault, SWF-South Wagad Fault VF-Vigodi Fault. | |
| | Uplifts: CHR-Central Hill Range, KMU-Kachchh Mainland Uplift, IBI-Island Belt Upland, NHR- Norhten Hill Range, NPU-Nagar Parkar Uplift and WU-Wagad Uplift. | |
| 1.4 | Depositional Domain of Kachchh Rift Basin | .12 |
| | BHG-Banni Half Graben, GOKHG- Gulf of Kachchh Half Graben, IBU-Island Belt Uplift, KU-Kathiyawar Uplift, NPU-Nagar Parkar Uplift, KMU-Kachchh Mainland Uplift, PU-Pancham Uplift, RHG-Rann Half Graben (After Biswas 2016a). | |

| Figuro No. | es | Page No . |
|---------------|--|--------------|
| 3.1 | Topographic-Structure Lineaments and Drainage Map, Kachchh Rectangular area exposures were studied extensively namely Kotda area (KA) Fakirwadi area (FA), Gangeswar area (GA) and Varli area (VA). | 36 |
| | Anjar; 2. Asambia; 3. Bharasar; 4. Bharudia; 5.Bhopa Dunger 237m; 6. Bhuj; 7. Boha; 8. Chakar; 9. Chiyasar; 10. Conical Hill 281m 11. Desaplar (south); 12. Desaplar; 13. Devpur; 14. Dhinordhar Hill 358m; 15. Dhribiyo Dunger 348m; 16. Don; 17. Fakirwadi Hill 258m; 18. Fotdi; 19. Gadhsisa; 20 Gajod; 21. Gangaji; 22. Gangeswar Dome 225m; 23.Gunawari Tributary; 24.Gunawari Southern Plain 25. Kera; 26. Khari Tributary; 27.KMF-Kachchh Mainland Fault; 28. Koday; 29. Kotda; 30. Kotdi; 31. Katrol Hill 349m; 32. Jhura Hill 344m; 33. Layza; 34. Ler Dam; 35. Nagrecha; 36. Naredi; 37 Nakhtrana; 38 Madhapar; 39. Mandvi; 40. Mudra; 41. NKHF-North Katrol Hill Fault; 42. Marutonk Dome 263m; 43. Sumarasara; 44. SKHF-South Katrol Hill Fault; 45. Satpura Dunger-250m; 46.Raydhanjar; 47. Ratadiya; 48.Ramapar Wakra; 49.Ratiya Nana; 50. Roha-Roha fort-Trappear Hill, 266m; 51. Varamsida- Nana Dunger 433m-Trappean Hill; 52. Tapkeswari 255m; 53. Varli, 160m; 54. Poladiya; 55. Zikdi. | 1 |
| 3.2 | Schematic diagram showing tectonic configuration of tilt blockstructure of southern Kachchh Mainland along with Tectonogeomorphic features. (After Das et al., 2016b) | 37 |
| 3.3 | Geomorphology of study area Katrol Hill Range (KHR) Source-1LIS III Data, ISRO, GSI and National Geomorphological and lineament Mapping. | 37 |
| 3.4 | Digital Elevations Model of study area showing depression and domal structures of south dipping Mesozoics, forming East-West trending gorge and valley structures. Also contrast between North of the KHF and rugged topography to the South of the KHF. | 38 |

| Figure No. | es Page No. |
|---------------|--|
| 3.5 | Katrol Hill drainage map |
| 3.6 | Geomorphology of Kotda area |
| 3.7 | Geomorphology of Varli area (VA) |
| 4.1 | Relief and drainage map of southwestern Kachchh46 |
| 4.2 | Three principal types of Miliolite occurrences in Kachchh47 |
| 4.3 | Map showing the locations of field occurrences studied |
| 4.4 | Vertical litho-logs from the Gangeswar area |
| 4.5 | Field photographs 52 |
| | (a) O1a- Obstacle miliolite deposit, KHF scarp (b) O1b- Elephant's skin weathered structure (c) O3- Miliolite overlain south west dipping sandstone (d) O4-Patchy obstacle deposit of miliolitic sand, buff yellowish to reddish in colour along with platy block of sand stone, excavation site southwest. (e) V1-About 20 to 25m thick pile of loose valley fill Miliolitic sand, North to Gunawari river.(f) V3- Loose valley fill miliolitic sand on plain terrain of southern range of KHR. |

| Figuro No. | es | Page No. |
|---------------|--|-------------|
| 4.6 | Field photographs | 53 |
| | (a) O5- Patchy deposit of miliolite on rocky amphitheatre, GD,259m (b) Aeolianite non conformably rests on an erosional surface (c) O11- Valley fill, Khari River Catchment. (d) V5- Loose valley fills Khari catchment. | |
| 4.7 | Field photographs | 56 |
| | (a) F1- Ler Dam Basin-Downstream Gunawari river-Southern flank (b) F2- Gunawari Northern Flank (c) F3-Lithological contact between Miliolite and fossiliferous sst. (d) F7- Gunawari cliff-10m, miliolitic sand, near Gangeswar Mahadev (e) F10- Khari cliff, reworked fluvial sheet miliolite and valley fill miliolite. | |
| 4.8 | Geological set up of Gangeswar area (GA), KHR | 57 |
| 4.9 | Locations of the Fakirwadi area (FA) with reference to Khari river, KH | R59 |
| 4.10 | Vertical Profile FR high land of KHR | 60 |
| | showing successive development of the erosional surfaces of Mesozoic stratum cross section is drawn along XY Lines shown in (Fig.4.9). | |
| 4.11 | Field photographs | 62 |
| | (a) O8-Abandoned Miliolite Quarry, 170 m at Khari catchment, Fakirwadi. (b) O9- Abandoned Miliolite subsurface Quarry,165m at Bhuj-Mandvi Road, Fakirwadi (c) O10- Fakirwadi mound, the rock are showing gently, tabular, planar cross-stratification, width prominent inclination towards North on an average. The thickness of the mound is 3-5m and the width is about 500 m. The mound rests on slightly weathered Jhuran Shale. (d) O10a-Fakirwadi Dome colluviam | |

| Figur No. | es Page No. |
|--------------|---|
| 4.12 | Field photographs |
| | (a) F8- Khari cliff-8m reworked fluvial sheet and valley fill miliolites. (b) F9- Khari cliff-12m, valley fill Miliolite and Jhuran shale. (c) F16a- Fluvial signature. The stratified sandy sheet of impure Miliolite deposits containing gravels and pebbles pointing towards its fluvial deposition are recognized as the fluvial reworked sheet deposits. (d) F11-Khari cliff-12 to 13m, and valley fill miliolite. (e) F16- 3rd order Chock Nadi about 2m cliff of miliolite sheet resting on Mesozoic, SE of Rayadhanjar village |
| 4.13 | Geological set up of Fakirwadi area (FA), KHR66 |
| 4.14 | SRTM DEM based contour map of Kotda-Roha Area (KA) 67 |
| 4.15 | Vertical profile of KA high land of KHR68 |
| | showing successive development of the erosional surfaces of Mesozoic-Deccan stratum From Pos 23° 05'N, 69° 01E to Pos 23° 12'N, 69° 30'E; with stratigraphy exposed and excavated litholog, depicting successive development of miliolite deposits. Cross sections drawn along AB lines (Fig.4.13). |
| 4.16 | South to Varamsida a conspicuous Trap Hill, famous as Nana Dunger, 433m.Obstacle leeward, tabular dune bedding of miliolite deposit as abandoned quarry exposure (O13) |
| 4.17 | Obstacle Miliolite dune deposit, abandoned quarry, Nana Dunger |
| 4.18 | Field photograph showing the lenses of angular basalt fragments in70 miliolite at Varamsida (O13). |
| 4.19 | Roha fort trappean hill, 266m just north edge on Roha village, windward obstacle dune deposits Field photograph (O14) 71 |
| 4.20 | Gravity derived weathered sandstone fragments as well as Deccan basalt angular fragments (O14)72 |

| Figure No. | es F | 'age No. |
|---------------|---|-------------|
| 4.21 | DEM of the Varli Area (VA) showing the position of the Varli Miliolite occurrences and South Katrol Hill Fault (KHF). | 74 |
| 4.22 | An outcrop of the Type-I miliolite deposit showing thin laminations at Varli (O17) | 75 |
| 5.1 | Grain size data of the friable miliolite samples plotted on triangular diagram using the Gradisat software | 81 |
| 5.2 | Relative abundances of Carbonate and Residue constituent of Miliolite Rocks, KHR, Kachchh | 82 |
| 5.3 | Abundance constituents of Miliolite of the study area | 84 |
| 5.4 | Thin section- photomicrographs of Miliolite, KHR, Kachchh | 85 |
| 5.5a | Scanning Electron Microscope (SEM) Hitachi, Germany | .87 |
| 5.5b | Energy Dispersive X-ray Spectroscopy (EDX),Oxford Instruments,UK | 87 |
| 5.6 | Fibrous form of calcite seen as cement in Type-I Miliolite from from Gangeswar area | 88 |
| | It is described as mineral 'Lublinite' by Stoops (1976). | |
| 5.7 | Granular nature of the micritic grains and rhomboidal shape of the calcite cement in Type-II Miliolites. | 89 |
| | Presence of minor amount of Mg, Al and Fe peaks in EDS spectra indicate detrital impurities added to the pure calcareous nature of the Type-I deposits. | |
| 5.8 | Presence clay flaks of minerals in inter granular space in Type-III Miliolites | 89 |
| | As EDS spectra show peaks of Mg, Fe, Na, K, and Al these could be the hydrous silicates representing the Illite-Kaolinite group of clays. | |

| Figure No. | es P | age No. |
|---------------|--|------------|
| 5.9 | X-Ray Fluorescence Spectroscopy (XRF) machine (Spectro Xepos He) | 90 |
| 5.10 | Kappa bridge (MFK-1B) AGICO | .95 |
| 6.1 | Shows bleaching event, subsequently the remnant signal builds up again through exposure to the weak natural flux of nuclear radiation | 98 |
| 6.2 | Flowchart followed in OSL Dating in present study done at the Institute of seismological Research (ISR), Gandhinagar. | 99 |
| 6.3 | The OSL measurements and radial plots of the De values | 102 |
| 7.1 | Principal types of miliolite occurrences in Kachchh | 105 |
| 7.2 | Geochemical Result of four bond elements of KHR, Kachchh Miliolite | 108 |
| 7.3 | The relationship between types and mean MS and linear relations | 109 |
| 7.4 | Summary of the available OSL geochronology of Miliolites in the KHR | 112 |
| 7.5 | Schematic diagram shows overlay of exposed cliff section along the Khari river to the SE to Bharasar showing various lithologies, fault and offsetting in the Quaternary sediments, also optical stimulated luminescence (OSL) dating sample location;(Kundu et al.2010). | .113 |
| 7.6 | A conceptual model of Late Quaternary history of Katrol Hill Range with reference to the Miliolite deposition in Kachchh | 116 |

LIST OF TABLES

| Tables No. | Pa N | ge o. |
|---------------|---|----------|
| 1.1 | Updated and modified lithostratigraphic classification of the Mesozoic rocks of Kachchh Mainland | 5 |
| 1.2 | Kachchh Basin tectono-sedimentary history of the Kachchh basin18 (After Biswas 2016a and b) | |
| 4.1 | Details of the sample codes and its locations in the field | 5-77 |
| | (Gangeswar-GR, Fakirwadi-FR,Kotda-KR and Varli-VR) and its locations in the field (Fig Ref-3.1, 4.3, 4.9 & 4.14). Type identification's-Obstacle dune deposit; V-Valley fill deposit; F- Fluvial reworked sheet deposit. | |
| 5.1 | Results of the granulometric analysis of the samples from Gangeswar Area (GA)79 | 9 |
| 5.2 | Results of the granulometric analysis of the samples from Fakirwadi area (FA) | С |
| 5.3 | Summary of the granulometric study of three types of Miliolite deposits | 1 |
| 5.4 | Relative frequency of detritals, allochems, cement and residues | 5 |
| 5.5 | Geochemical Result of four bond elements XRF | 1 |
| 5.6 (1) | Geochemical Results of bond elements of Miliolite, KHF, Kachchh 92 | 2 |
| 5.6(2) | Geochemical Results of bond elements of Miliolite, KHF, Kachchh 93 | 3 |
| 5.7 | Magnetic Susceptibility values of miliolite samples | 5 |

| Tables No. | P I | 'age No. |
|---------------|---|-------------|
| 6.1 | Summary of the types of Quaternary carbonate deposits and their optically stimulated luminescence age ranges in the study area |)3 |
| 6.2 | U, Th, and K concentrations and optically stimulated luminescence data. Radioactivity, palaeodose, dose rate and luminescence ages 10 (water content = 10 ± 2 , Juyal et al., 2006) | 13. |
| 7.1 | Abrasion test values | 19 |
| 7.2 | Stratigraphy, lithological characteristics, occurrence, ages and depositional events of Late Quaternary sequences in KHR, Kachchh11 | .4 |