

---

## List of Figures

Fig. 1.1: Global geographical prevalence of BB.....	6
Fig. 1.2: Schematic diagram showing host and tissue specificity amongst Xanthomonads .....	8
Fig. 1.3: Typical cells and colony morphology of Xoo .....	8
Fig. 1.4: Typical symptoms of the BB disease observed on the rice plant infected with Xoo.....	9
Fig. 1.5: Schematic diagram showing virulence determinants of Xoo and their interaction with host cell .....	11
Fig. 1.6: Schematic representation of small molecule compounds as antibiotic potentiators and as virulence attenuators .....	19
Fig. 1.7: Schematic diagram depicting Multidrug efflux pumps .....	22
Fig. 1.8: Proposed mechanisms of efflux pump inhibition by EPIs.....	24
Fig.2 1: Inhibition of Xoo BXO43 by the representative isolates.....	39
Fig.2 2: Phylogenetic tree based on 16S rRNA gene sequences of the six isolates and closely related type strains .....	42
Fig.2 3: Effect of isolates on growth of rice plant.....	43
Fig.2 4: Effect of isolates on disease symptoms by detached leaf assay .....	45
Fig.2 5: Recovery of the selected bacterial antagonists from root of the rice plants .....	46
Fig.2 6: Effect of supernatant of <i>B. altitudinis</i> S2 in preventing the disease symptoms .....	47
Fig.2 7: Gel image showing ARDRA band patterns of <i>gyrB</i> amplicon from isolate S2 and type strains <i>B. altitudinis</i> 41KF2b and <i>B. pumilus</i> ATCC 7061.....	48
Fig.2 8: Phylogenetic tree based on <i>gyrB</i> gene sequence of isolate S2 and other related type strains of the <i>Bacillus</i> sp.....	49
Fig. 3.1:Effect of physical parameters on antibiotic S2 production.....	68
Fig. 3.2: Time-course Production of antibiotic S2 .....	69
Fig. 3.3: Kresek disease model showing potential of antibiotic S2 .....	70
Fig. 3.4: TLC-DB of crude extract of <i>B. altitudinis</i> S2 .....	72
Fig. 3.5: Absorption spectrum of extract containing antibiotic S2 .....	73
Fig. 3.6: chromatographs obtained by HPLC and peaks by mass spectrometry of antibiotic S2 .....	74
Fig. 3.7: Gel image showing amplification of pks and pmt genes from <i>B. altitudinis</i> S2 and type strain <i>B. altitudinis</i> 41KF2b .....	75
Fig. 3.8: Micrographs showing effect of exposure to antibiotic S2 using Live-Dead staining kit.....	76
Fig. 3.9: Time-kill assay of antibiotic S2 using Xoo BXO43.....	77

---

---

---

Fig. 3.10: Scanning electron micrograph depicting surface morphology of Xoo BXO43 and <i>S. aureus</i> when exposed to antibiotic S2 .....	78
Fig. 3.11: SDS-PAGE showing effect of antibiotic S2 on total protein of Xoo BXO43 .....	79
Fig. 4.1: Standardization of sub-inhibitory concentration of compounds based on the growth of Xoo BXO43 in PS broth.....	95
Fig. 4. 2: Effect of subinhibitory concentration of compounds on growth of Xoo BXO43 .....	96
Fig. 4.3: Construction of transcriptional reporter vector.....	97
Fig. 4.4: Detection of Cellulase and Pectinase secreted by Xoo BXO43. ....	99
Fig. 4.5: Effect of small molecules on production of exoenzymes.....	99
Fig. 4.6: EPS production by Xoo BXO43 in terms of glucose equivalents. ....	100
Fig. 4.7: Motility assay of Xoo BXO43.....	101
Fig. 4.8: Detached leaf assay to study the effect of selected compounds on pathogenesis by Xoo BXO43.....	102
Fig. 5 1: Checkerboard assay for calculation of FIC index of two compounds..	108
Fig. 5 2: Fluorescence of Xoo BXO43 on PSA plates containing without and with 5 and 10 µg/ml EtBr .....	117
Fig. 5 3: Effect of small molecule compounds on the ethidium bromide efflux from Xoo BXO43 cells .....	117
Fig. 5 4: Effect of caffeic acid on expression of different efflux pump genes of Xoo BXO43. ....	118