## **List of Tables**

Table 2.1: Primers used for the molecular characterization of the antagonistic isolates
Table 2.2: Source of screened isolates and their inhibitory characteristics against Xoo BXO43
Table 2.3: Cz/Cs ratio of selected isolates against the pathogen Xoo BXO43 and the plant and tissue source from which the bacteria were isolated
Table 2.4:Morphological and biochemical characterization of selected six antagonistic bacterial isolates
Table 2.5: Isolates identified by 16S rRNA sequencing and their accession number in NCBI
Table 2.6: Quantitation of effect of isolates on rice plant in terms of root, shoot length and dry weight of root and shoot of rice plant
Table 2.7: Antimicrobial spectrum of <i>B. altitudinis</i> S2
Table 3.1: Primers designed for genes from polyketide synthesis pathway 66
Table 3.2: Screening of media and organic solvents for production and extraction of antibiotic S2 produced by <i>B. altitudinis</i> S2
Table 3.3: Plant parameters to evaluate efficacy of antibiotic S2 in protecting rice plant
Table 3. 4: Stability of antibiotic S2 in different treatments
Table 4.1: Strains, plasmids and oligonucleotides used in construction of transcription-reporter assay system in Xoo BXO43
Table 4.2: Effect of compounds at sub-inhibitory concentration on promoter activity in Xoo BXO43XOPQ harbouring reporter plasmid and grown in XOM2 medium98
Table 4.3: Quantification of swimming motility in terms of diameter covered by Xoo BXO43 in the presence of chemicals
Table 5.1: Antibiotics groups based on their mode of action and chemical nature
Table 5.2: Primers designed for different efflux pump genes for qRT-PCR 113
Table. 5.3: Effect of small molecule compounds on the MIC of antibiotic S2 Without against Xoo BXO43
Table 5.4: Effect of caffeic acid on MIC of other antibiotics against Xoo BXO43