

# List of Figures

Figure 1.1: (a) Plot of activation energies for an ordinary gas phase reaction (blue) and the same reaction with an appropriate catalyst (green). (b) Energy path for atoms (blue) and molecules (green) approaching a surface. (c) Schematic illustration of the formation of a chemical bond between an adsorbate valence level and the s and d states of a transition metal surface. (d) Illustration of the elementary reaction steps on surfaces.. 2

Figure 1.2: (a) Relevance of catalysis for the chemical industry. Over 90% of the world's manufactured consumer goods involve catalysis at one or more stages. Approximately 80% of all catalytic processes require heterogeneous catalysts, 15% homogeneous catalysts, and 5% biocatalysts [7]. (b) The evolution of heterogeneous catalysis from the 19<sup>th</sup> to the 21<sup>st</sup> century [2]. ..... 4

Figure 1.3: A timeline of the development of reactivity descriptors and scaling relationships in heterogeneous catalysis [18]. The grey, blue, and pink boxes represent electronic, structural, and binary descriptors, respectively..... 6

Figure 1.4: Schematic of pure metal clusters. From left to right: icosahedral, decahedral, cuboctahedral, and tetrahedral, with point group symmetries. .... 8

Figure 1.5: The schematic illustration of E-R and L-H mechanism pathway for CO oxidation. .... 13

Figure 1.6: (a) Carbon cycle of CH<sub>3</sub>OH, a platform molecule for the chemical industry [79]. (b) Trend analysis on global CH<sub>3</sub>OH market size and forecast by end-use [80]. .. 15

Figure 2.1: From many-body problem to density functional theory (DFT). Born-Oppenheimer approximation, Hohenberg-Kohn theorem and Kohn-Sham ansatz. .... 27

Figure 2.2: Schematic flowchart for finding self-consistent solutions of the Kohn-Sham equations. .... 31

Figure 2.3: Comparison of a wavefunction in the Coulomb potential of the nucleus (blue) to the one in the pseudopotential (red). The real and the pseudo wavefunction and potentials match above a certain cut-off radius (core radius)  $r_c$ . Sketch inspired by Ref. [17]..... 34

Figure 2.4: Flowchart of Geometry optimization..... 35

Figure 2.5: Schematic illustration of broken symmetry and functional form of the density of states in 1D, 2D, and 3D confined materials [28]. .....	38
Figure 2.6: Schematic illustration of the spin-polarized DOS of the transition metal with the electronic reactivity descriptors. ....	40
Figure 2.7: Density functional theory calculations of the minimum energy path for CH <sub>4</sub> dissociative adsorption on an Ir(111) surface [43]. .....	43
Figure 3.1: Lowest-energy structures (obtained with PBE functional) of the Ni <sub>n+1</sub> and Ni <sub>n</sub> Cu cluster (2 ≤ n ≤ 12). ....	54
Figure 3.2: Skeleton representation of lowest-energy structures of the Ni <sub>n+1</sub> cluster (2 ≤ n ≤ 12) with their corresponding effective coordination number (ECN) and point-group symmetry (structure). ....	55
Figure 3.3: Skeleton representation of lowest-energy structures of the Ni <sub>n</sub> Cu cluster (2 ≤ n ≤ 12) with their corresponding effective coordination number (ECN) and point-group symmetry (structure). ....	56
Figure 3.4: Simulated XANES spectra of (a-b) Ni <sub>n+1</sub> clusters and (c-d) Ni <sub>n</sub> Cu clusters (1 ≤ n ≤ 12). The shaded area in cyan color represents the pre-edge region and its enlarged counterparts shown in the right panel of the corresponding XANES spectra. The calculated E <sub>rising-edge</sub> (blue color) and E <sub>white-line</sub> (red color) peaks corresponding to the (e) Ni <sub>n+1</sub> clusters and (f) Ni <sub>n</sub> Cu clusters (1 ≤ n ≤ 12). ....	58
Figure 3.5: Mixing energy per atom ( $E_{mix}$ ) of Ni <sub>n</sub> Cu clusters (1 ≤ n ≤ 12). The black and light green symbols represent the positive and negative value of ( $E_{mix}$ ), respectively. .	60
Figure 3.6: (a) Schematic orientation for the adsorbate (in x, y, and z axis) onto the adsorbent in four reference positions, i.e., bridge (parallel to the xz plane), hollow (perpendicular to the xy plane and parallel to the z axis), top (aligned parallel to the z direction), and side-on (parallel to the xz plane). The tilted orientation shows the deviation of adsorbate state after complete relaxation in some cases. In some cases, the adsorbate deviates from top to tilted orientation at a certain angle (θ) after complete relaxation. Lowest energy structures of (b) Ni <sub>n+1</sub> CO and (c) Ni <sub>n</sub> CuCO systems.....	62
Figure 3.7: Heatmap of CO adsorption energy ( $E_{CO-ads}$ eV) over (a) Ni <sub>n+1</sub> and (b) Ni <sub>n</sub> Cu clusters with respect to different functionals and their combination with dispersion corrections (D2 and D3).....	66
Figure 3.8: The spin-polarized projected density of states (PDOS) of (a) Ni <sub>9</sub> , (b) Ni <sub>13</sub> , (c) Ni <sub>8</sub> Cu, and (d) Ni <sub>12</sub> Cu clusters. The short dash line in black and red color represents the <i>d-band</i> center of Ni and Cu respectively for spin-up and spin-down state. The long	

dash line at zero indicates the Fermi energy. The filled and unfilled regions represent the bonding and anti-bonding states, respectively..... 69

Figure 3.9: (a) The spin-polarized density of states of CO molecule and labeled orbitals. The spin-polarized projected density of states of (b) Ni<sub>9</sub>CO, (c) Ni<sub>13</sub>CO, (d) Ni<sub>8</sub>CuCO and (e) Ni<sub>12</sub>CuCO systems. The green dash line indicates the HOMO and LUMO state of the CO molecule. The short dash line in black and red color represents the *d-band* center of Ni and Cu respectively. The long dash line at zero indicates the Fermi energy. The filled and unfilled regions for the metals represent the bonding and anti-bonding states, respectively. .... 70

Figure 3.10: The molecular orbital of HOMO and LUMO state (for spin-up and spin-down) of (a) Ni<sub>9</sub>CO and Ni<sub>13</sub>CO systems and (b) Ni<sub>8</sub>CuCO and Ni<sub>12</sub>CuCO systems with an iso-surface value of  $\pm 0.0004 \text{ e}/\text{\AA}^3$ . .... 74

Figure 3.11: Charge density difference for CO adsorbed Ni<sub>9</sub>, Ni<sub>13</sub>, Ni<sub>8</sub>Cu and Ni<sub>12</sub>Cu clusters. The orange (green) color represents the charge accumulation (depletion). The value of iso-surface is set to  $\pm 0.005 \text{ e}/\text{\AA}^3$  for Ni<sub>9</sub>CO and Ni<sub>13</sub>CO systems. The value of iso-surface is set to  $\pm 0.01 \text{ e}/\text{\AA}^3$  for Ni<sub>8</sub>CuCO and Ni<sub>12</sub>CuCO systems. The charge density difference is calculated using the formula  $\Delta\rho = \rho(\text{Cluster} + \text{gas/atom}) - \rho(\text{Cluster}) - \rho(\text{gas/atom})$ . .... 75

Figure 3.12: The change of work function  $\Delta\Phi$  (eV) induced by CO adsorption over Ni<sub>n+1</sub> (left side) and Ni<sub>n</sub>Cu (right side) clusters. The generalized formula to calculate work function for bare system or gas adsorbed system is  $\Phi = E_{\text{vac}} - E_{\text{F}}$ , where,  $E_{\text{vac}}$  is the energy of vacuum level of the system and  $E_{\text{F}}$  is the energy of Fermi level of the system. .... 77

Figure 3.13: The complete L-H reaction pathway and activation barrier for CO oxidation over Ni<sub>9</sub> cluster. IS, IM, TS, and FS indicate initial state, intermediate state, transition state, and final state of reaction, respectively. The orange, black, and red balls represent Ni, C, and O atoms, respectively. The thermodynamically unfavored intermediates are connected with a red line. .... 78

Figure 3.14: The complete L-H reaction pathway and activation barrier for CO oxidation over the Ni<sub>13</sub> cluster. IS, IM, TS, and FS indicate initial state, intermediate state, transition state, and final state of reaction, respectively. The orange, black, and red balls represent Ni, C, and O atoms, respectively. The thermodynamically unfavored intermediates are connected with a red line..... 79

Figure 3.15: The complete L-H reaction pathway and activation barrier for CO oxidation over Ni<sub>8</sub>Cu cluster. IS, IM, TS, and FS indicate initial state, intermediate state,

transition state, and final state of reaction, respectively. The orange, green, black, and red balls represent Ni, Cu, C, and O atoms, respectively.....	80
Figure 3.16: The complete L-H mechanism reaction pathway and activation barrier for CO oxidation onto the Ni <sub>12</sub> Cu cluster. IS, IM, and TS indicate initial state, intermediate state, and final state of reaction respectively. The orange, green, black, and red atoms represent Ni, Cu, C, and O respectively. ....	81
Figure 4.1: The optimized configuration of I <sub>h</sub> Pd <sub>m</sub> Cu <sub>n</sub> (m + n = 13) clusters with different Cu compositions. ....	94
Figure 4.2: Average binding energy per atom ( $E_b$ eV) of I <sub>h</sub> Pd <sub>m</sub> Cu <sub>n</sub> (m + n = 13) clusters with a different numbers of Cu atoms. ....	95
Figure 4.3: Second order energy difference ( $\Delta^2E$ eV) of I <sub>h</sub> Pd <sub>m</sub> Cu <sub>n</sub> (m + n = 13) clusters with a different numbers of Cu atoms. ....	97
Figure 4.4: Average bond length $\bar{R}$ (Å) of I <sub>h</sub> Pd <sub>m</sub> Cu <sub>n</sub> (m + n = 13) clusters with different numbers of Cu atoms. ....	99
Figure 4.5: The PDOS of (a) Pd <sub>13</sub> , (b) Cu <sub>13</sub> , and (c) Pd <sub>5</sub> Cu <sub>8</sub> clusters. The Fermi level is set to zero (black line).....	100
Figure 4.6: (a-n) Raman spectra of I <sub>h</sub> Pd <sub>m</sub> Cu <sub>n</sub> (m + n = 13) clusters.....	103
Figure 4.7: The vibrational mode of I <sub>h</sub> Pd <sub>m</sub> Cu <sub>n</sub> (m + n = 13) clusters corresponding to the high intense peak of their respective Raman spectra.....	104
Figure 4.8: Different adsorption geometries for CO <sub>2</sub> adsorption over I <sub>h</sub> Pd <sub>m</sub> Cu <sub>n</sub> (m + n = 13) clusters.....	108
Figure 4.9: The CO <sub>2</sub> adsorbed I <sub>h</sub> Pd <sub>m</sub> Cu <sub>n</sub> (m + n = 13) clusters. The green, light pink, yellow, and red color sphere indicates Pd, Cu, C, and O atoms, respectively.....	108
Figure 4.10: (a-c) Charge density plots of CO <sub>2</sub> adsorbed Pd <sub>13</sub> , Pd <sub>5</sub> Cu <sub>8</sub> and Cu <sub>13</sub> clusters. The green, light pink, yellow, and red color sphere indicates Pd, Cu, C and O atoms, respectively.....	109
Figure 4.11: The possible reaction pathways of CO <sub>2</sub> hydrogenation to HCOOH, CH <sub>3</sub> OH, and CH <sub>4</sub> on the Pd <sub>5</sub> Cu <sub>8</sub> cluster. The symbol (*) and (g) indicates the adsorbed and gas phase species, respectively.....	114
Figure 4.12: The low-lying adsorption configurations of intermediates in the hydrogenation of CO <sub>2</sub> on the Pd <sub>5</sub> Cu <sub>8</sub> cluster. The white, orange, black, red, and cyan balls represent Pd, Cu, C, O, and H atoms, respectively. ....	116

Figure 4.13: The initial state (IS), transition state (TS), and the final state (FS) of elementary step in CO <sub>2</sub> hydrogenation over the Pd <sub>5</sub> Cu <sub>8</sub> cluster from R1 to R12.....	118
Figure 4.14: The initial state (IS), transition state (TS), and the final state (FS) of elementary step in CO <sub>2</sub> hydrogenation over the Pd <sub>5</sub> Cu <sub>8</sub> cluster from R13 to R18.....	119
Figure 4.15: The optimized pathways for the production of HCOOH, CH <sub>3</sub> OH, and CH <sub>4</sub> over the Pd <sub>5</sub> Cu <sub>8</sub> cluster.....	120
Figure 5.1: Surface sites, bond lengths and angles of Cu <sub>2</sub> O(111). Cu <sub>usa</sub> and Cu <sub>sa</sub> correspond to unsaturated and saturated copper surface sites, respectively. O <sub>up</sub> and O <sub>dn</sub> correspond to unsaturated and saturated oxygen surface sites, respectively. d(O <sub>dn</sub> –Cu <sub>sa</sub> ) and d(O <sub>up</sub> –Cu <sub>sa</sub> ) are the bond length of O <sub>dn</sub> and O <sub>up</sub> to Cu <sub>sa</sub> atoms, respectively. d(Cu <sub>usa</sub> –Cu <sub>sa</sub> ) is the distance between Cu <sub>usa</sub> and Cu <sub>sa</sub> atoms. Color key: white, Cu; and red, O.....	133
Figure 5.2: (a) The density of states of Cu <sub>2</sub> O(111) surface. (b) The charge density plot of valence band maximum (VBM) and conduction band minimum (CBM) of Cu <sub>2</sub> O(111) surface. The iso-surface value set to $\pm 0.0003$ e/Å <sup>3</sup> . The blue and red color represent the depletion and accumulation of charge, respectively. (c-d) The projected density of state (PDOS) of O and Cu atoms, respectively. (e) The average electrostatic potential of the Cu <sub>2</sub> O(111) surface as a function of the distance in the z-direction.....	135
Figure 5.3: Top (upper panel) and side views (middle panel) of the optimized geometries of Pt <sub>4</sub> /Cu <sub>2</sub> O(111), Pt <sub>3</sub> Au/Cu <sub>2</sub> O(111), and Pt <sub>3</sub> Co/Cu <sub>2</sub> O(111) systems. The charge density difference (CDD) of the Pt <sub>4</sub> /Cu <sub>2</sub> O(111), Pt <sub>3</sub> Au/Cu <sub>2</sub> O(111), and Pt <sub>3</sub> Co/Cu <sub>2</sub> O(111) systems (bottom panel). Yellow and light green regions in CDD represent charge depletion and accumulation, respectively. The iso-surface value is set to $\pm 0.2$ e/Å <sup>3</sup> for CDD. Color key: white, Cu; red, O; orange, Pt; yellow, Au; and dark green Co.....	137
Figure 5.4: (a) The total density of states of the Pt <sub>4</sub> /Cu <sub>2</sub> O(111) system with the HOMO and LUMO orbitals where charge depletion indicated by blue region and accumulation of charge indicated by red region. The iso-surface value is set to $\pm 0.0007$ e/Å <sup>3</sup> . (b-e) Atom-wise projected density of states of the Pt <sub>4</sub> /Cu <sub>2</sub> O(111) system. The black and red lines indicate the HOMO and LUMO levels. The Fermi energy is set to zero.....	141
Figure 5.5: (a) The total density of states of the Pt <sub>3</sub> Au/Cu <sub>2</sub> O(111) system with the HOMO and LUMO orbitals where charge depletion indicated by blue region and accumulation of charge indicated by red region. The iso-surface value is set to $\pm 0.004$ e/Å <sup>3</sup> . (b-e) Atom-wise projected density of states of the Pt <sub>3</sub> Au/Cu <sub>2</sub> O(111) system. The black and red lines indicate the HOMO and LUMO levels. The Fermi energy is set to zero.....	142

---

Figure 5.6: (a) The total density of states of the  $\text{Pt}_3\text{Co}/\text{Cu}_2\text{O}(111)$  system along with the molecular orbitals of the HOMO and LUMO states. The iso-surface value is set to  $\pm 0.002 \text{ e}/\text{\AA}^3$ . (b-e) Atom-wise projected density of states of the  $\text{Pt}_3\text{Co}/\text{Cu}_2\text{O}(111)$  system. The black and red lines indicate the HOMO and LUMO levels. The Fermi energy is set to zero. .... 143

Figure 5.7: The optimized geometry and structural parameter (bond length) of CO (upper panel),  $\text{CO} + \text{O}_2$  (middle panel), and  $\text{CO} + \text{O}$  (bottom panel) over  $\text{Pt}_4/\text{Cu}_2\text{O}(111)$  and  $\text{Pt}_3\text{X}/\text{Cu}_2\text{O}(111)$  ( $\text{X} = \text{Au}$  and  $\text{Co}$ ) systems. Color key: white, Cu; red, O; black, C; orange, Pt; yellow, Au; and dark green, Co. .... 145

Figure 5.8: The complete L-H mechanism reaction pathway for CO oxidation onto the  $\text{Pt}_4/\text{Cu}_2\text{O}(111)$  system. Color key: white, Cu; red, O; black, C; and orange, Pt. .... 147

Figure 5.9: The complete L-H mechanism reaction pathway for CO oxidation onto the  $\text{Pt}_3\text{Au}/\text{Cu}_2\text{O}(111)$  system. TS indicates a transition state. Color key: white, Cu; red, O; black, C; orange, Pt; and yellow, Au. .... 147

Figure 5.10: The complete L-H mechanism reaction pathway for CO oxidation onto the  $\text{Pt}_3\text{Co}/\text{Cu}_2\text{O}(111)$  system. TS indicates a transition state. Color key: white, Cu; red, O; black, C; orange, Pt; yellow, Au; and dark green, Co. .... 148