

# List of Tables

|     |                                                                                                                                                                                                                                                                                         |    |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| 1.1 | Quantities of minor actinides produced annually from a conventional pressurized water reactor with a standard burn-up of 33 gigawatt per day. These quantities are quoted for 1 ton and 23 tons of annual consumption of uranium. . . . .                                               | 4  |
| 1.2 | List of medical isotopes used for cross-section measurements in the present work . . . . .                                                                                                                                                                                              | 10 |
| 2.1 | Target details for the n-induced reaction cross-section measurements. . . . .                                                                                                                                                                                                           | 28 |
| 2.2 | Target details for the p-induced reaction cross-section measurements. . . . .                                                                                                                                                                                                           | 29 |
| 2.3 | List of some of the neutron generator reactions with their spectroscopic properties [10,11]. . . . .                                                                                                                                                                                    | 29 |
| 2.4 | List of the commonly used neutron monitor reactions with their spectroscopic properties [11] . . . . .                                                                                                                                                                                  | 29 |
| 2.5 | $\gamma$ -rays along with their absolute intensities (%) for the $^{152}\text{Eu}$ source . . . . .                                                                                                                                                                                     | 33 |
| 5.1 | A summary of the experimental work carried out at BARC FOTIA accelerator facility . . . . .                                                                                                                                                                                             | 72 |
| 5.2 | A summary of the experimental work carried out at BARC-TIFR Pelletron accelerator facility . . . . .                                                                                                                                                                                    | 73 |
| 5.3 | Nuclear spectroscopic data for the isotopes used in the measurement of the $^{232}\text{Th}(n, \gamma)$ , $^{100}\text{Mo}(n, 2n)$ and $^{58}\text{Ni}(n, x)$ reaction cross-sections. The data are taken from the NuDat [40], Q-value calculator and Q-tool [39] data library. . . . . | 76 |
| 5.4 | Tailing corrected cross sections for $^{232}\text{Th}(n, \gamma)^{233}\text{Th}$ reaction [10] . . . . .                                                                                                                                                                                | 84 |
| 5.5 | Tailing corrected cross sections for $^{100}\text{Mo}(n, 2n)^{99}\text{Mo}$ reaction [22] . . . . .                                                                                                                                                                                     | 84 |
| 5.6 | Tailing corrected cross sections for $^{58}\text{Ni}(n, p)^{58}\text{Co}$ reaction [23]                                                                                                                                                                                                 | 85 |
| 5.7 | Tailing corrected cross sections for $^{58}\text{Ni}(n, 2n)^{57}\text{Ni}$ reaction [23]                                                                                                                                                                                                | 85 |

## LIST OF TABLES

|      |                                                                                                                                                                      |     |
|------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 5.8  | Counting and spectroscopic details of the $^{152}\text{Eu}$ source used for the efficiency measurement . . . . .                                                     | 86  |
| 5.9  | Partial uncertainties in the efficiency . . . . .                                                                                                                    | 87  |
| 5.10 | Co-variance matrix ( $V_\varepsilon \times 10^5$ ) for the detector efficiencies . . . . .                                                                           | 88  |
| 5.11 | Correlation matrix for the detector efficiencies . . . . .                                                                                                           | 88  |
| 5.12 | Measured efficiencies with correlation matrix . . . . .                                                                                                              | 88  |
| 5.13 | Fractional uncertainties in various parameters to obtain $^{232}\text{Th}(n, \gamma)$ cross-section . . . . .                                                        | 89  |
| 5.14 | Covariance matrix (%) and corresponding correlation coefficients for the measured $^{232}\text{Th}(n, \gamma)$ cross-sections . . . . .                              | 90  |
| 5.15 | $^{232}\text{Th}(n, \gamma)$ reaction cross-sections at different neutron energies compare with ENDF-B/VII.1, JENDL-4.0 and TALYS-1.8 data . . . . .                 | 90  |
| 5.16 | Default and adjusted parameters used in the TALYS-1.8 calculations . . . . .                                                                                         | 93  |
| 5.17 | $\gamma$ -ray strength function parameters for Giant Dipole Resonance . . . . .                                                                                      | 94  |
| 5.18 | $^{100}\text{Mo}(n, 2n)^{99}\text{Mo}$ reaction cross-sections [22] compare with ENDF-B/VII.1, JENDL-4.0 and TALYS-1.9 data . . . . .                                | 95  |
| 5.19 | Default and adjusted parameters used in the TALYS-1.9 calculations . . . . .                                                                                         | 97  |
| 5.20 | Systematic formulas proposed by different authors for (n, p) and (n,2n) reactions . . . . .                                                                          | 99  |
| 5.21 | Systematic formulas proposed by different authors for (n, $\alpha$ ) reaction . . . . .                                                                              | 100 |
| 5.22 | Calculated (n, p), (n, 2n) and (n, $\alpha$ ) cross-sections using the systematic formulas given in Tables 5.20 and 5.21. . . . .                                    | 101 |
| 5.23 | The Measured cross sections for the $^{58}\text{Ni}(n, p)^{58}\text{Co}$ and $^{58}\text{Ni}(n, 2n)^{57}\text{Ni}$ reactions [23] . . . . .                          | 102 |
| 5.24 | A comparison of the calculated $[\sigma_{(n,p)}]$ results from different systematics with the data from EXFOR library at $\approx 14.5$ MeV neutron energy. . . . .  | 106 |
| 5.25 | A comparison of the calculated $[\sigma_{(n,2n)}]$ results from different systematics with the data from EXFOR library at $\approx 14.5$ MeV neutron energy. . . . . | 106 |
| 6.1  | List of the identified residues in the proton induced $^{nat}\text{Nb}$ , $^{nat}\text{Ag}$ and $^{nat}\text{Ti}$ reactions with their spectroscopic data [34,35].   | 128 |
| 6.2  | The experimentally measured cross-sections for the identified residues in the $p + ^{nat}\text{Nb}$ interaction. . . . .                                             | 131 |
| 6.3  | The experimentally measured cross-sections for the identified reaction residues in the $p + ^{nat}\text{Ag}$ interaction. . . . .                                    | 131 |

## LIST OF TABLES

|     |                                                                                                                                                                                       |     |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 6.4 | The experimentally measured reaction cross-sections for the identified residues in the $p + {}^{nat}Ti$ interaction. . . . .                                                          | 132 |
| 1   | Data related to the ${}^{152}Eu$ source to obtain the detector efficiency                                                                                                             | 164 |
| 2   | Partial uncertainties in the efficiency . . . . .                                                                                                                                     | 165 |
| 3   | Diagonal matrix ( $\times 10^{-3}$ ) for partial uncertainty from the attribute C (r=1) using the values given in the Table 2. . . . .                                                | 165 |
| 4   | Co-variance matrix ( $V_{\epsilon} \times 100$ ) for the detector efficiency . .                                                                                                      | 166 |
| 5   | Measured efficiencies with correlation matrix for the sample and the monitor reaction . . . . .                                                                                       | 171 |
| 6   | Fractional uncertainties in various parameters used to obtain ${}^{100}Mo(n, 2n){}^{99}Mo$ cross-sections . . . . .                                                                   | 172 |
| 7   | Covariance matrix and corresponding correlation coefficients for the measured ${}^{100}Mo(n, 2n){}^{99}Mo$ cross-sections . . . . .                                                   | 173 |
| 8   | Uncertainty (%) present in the measured ${}^{100}Mo(n, 2n){}^{99}Mo$ cross-sections at respective neutron energy . . . . .                                                            | 173 |
| 9   | ${}^{100}Mo(n, 2n){}^{99}Mo$ reaction cross-sections compare with ENDF-B/VII.1, JENDL-4.0 and TALYS-1.9 data . . . . .                                                                | 173 |
| 10  | Co-variance matrix ( $V_{\epsilon} \times 10^6$ ) for the efficiencies of the detector used at FOTIA. . . . .                                                                         | 174 |
| 11  | Co-variance matrix ( $V_{\epsilon} \times 100$ ) for the efficiencies of the detector used at Pelletron. . . . .                                                                      | 175 |
| 12  | Measured efficiencies with correlation matrix for the sample and the monitor reaction measured for set 1 . . . . .                                                                    | 176 |
| 13  | Measured efficiencies with covariance matrix for the sample and the monitor reaction measured for set 2 . . . . .                                                                     | 177 |
| 14  | Measured efficiencies with correlation matrix for the sample and the monitor reaction measured for set 2 . . . . .                                                                    | 177 |
| 15  | Fractional uncertainties in various parameters used to obtain ${}^{58}Ni(n, p){}^{58}Co$ reaction cross-sections . . . . .                                                            | 178 |
| 16  | Fractional uncertainties in various parameters used to obtain ${}^{58}Ni(n, p){}^{58}Co$ reaction cross-sections . . . . .                                                            | 178 |
| 17  | Fractional uncertainties in various parameters used to obtain ${}^{58}Ni(n, 2n){}^{57}Ni$ reaction cross-sections . . . . .                                                           | 178 |
| 18  | Covariance matrix (%) and corresponding correlation coefficients for the measured ${}^{58}Ni(n, p){}^{58}Co$ reaction cross-sections using the FOTIA at BARC . . . . .                | 179 |
| 19  | Covariance matrix (%) and corresponding correlation coefficients for the measured ${}^{58}Ni(n, p){}^{58}Co$ reaction cross-sections using the BARC-TIFR Pelletron facility . . . . . | 179 |

|    |                                                                                                                                                                                                 |     |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 20 | Covariance matrix (%) and corresponding correlation coefficients for the measured $^{58}\text{Ni}(n,2n)^{57}\text{Ni}$ reaction cross-sections using the BARC-TIFR Pelletron facility . . . . . | 179 |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|