## List of Figures

1.1	The pseudoscalar meson nonet in the quark model	5
1.2	Quark line diagrams for (a) OZI allowed decay $\omega \to 3\pi$ and (b) OZI for- bidden decay $\phi \to 3\pi$ .	13
2.1	Feynman diagrams considered for the correlation function $\Pi(q, p)$ . The dotted line stands for the meson.	29
2.2	Plots of our results for $g_{\eta NN}$ as a function of $M^2$ (thick solid line). Also plotted are our results on $g_{\eta NN}$ without contribution from gluonic operator $\alpha_s G\tilde{G}$ (long-dashed line), only due to $\alpha_s G\tilde{G}$ (thin solid line) and due to the part with $f_s$ in $a_\eta$ (short-dashed line). Figs.(a) and (b) are for $\eta$ when $\mathbf{p}^2$ is $-\frac{m_\eta^2}{2}$ and $-\frac{3m_\eta^2}{4}$ respectively	34
2.3	Plots of our results for $g_{\eta'NN}$ as a function of $M^2$ (thick solid line). Also plotted are our results on $g_{\eta'NN}$ without contribution from gluonic operator $\alpha_s G\tilde{G}$ (long-dashed line), only due to $\alpha_s G\tilde{G}$ (thin solid line) and due to the part with $f_s$ in $a_{\eta'}$ (short-dashed line). Figs.(a) and (b) are for $\eta'$ when $\mathbf{p}^2$ is $-\frac{2m_{\eta'}^2}{3}$ and $-\frac{3m_{\eta'}^2}{4}$ respectively	35
2.4	A Feynman diagram with a quark condensate contributing to the correla- tion function $\prod(x,p)$	36
3.1	The Feynman diagram for the $e^+e^- \rightarrow e^+e^-\pi^0$ process $\ldots \ldots \ldots$	40
3.2	Lowest order Feynman diagram for $\gamma^* \gamma \to P, P = \pi^0, \eta, \eta'$	44
3.3	Diagrams contributing to gluonic coefficient function to $\eta\gamma, \eta'\gamma$ TFF	52
3.4	Contribution to the TFF from a bilocal pseudoscalar operator along with a quark condensate.	56
3.5	Feynman diagram contributing to the TFF through a bilocal tensor oper- ator along with a quark condensate.	57
3.6	Feynman diagrams for TFF obtained by expanding quark propagator close to the light-cone in a background gluon field along with a quark condensate.	57

3.7	Feynman diagrams for TFF obtained from light-cone expansion of the prod- uct of two electromagnetic currents.	58
3.8	Feynman diagram for TFF obtained from expansion of a quark propagator near light-cone involving gluon DA and a quark condensate	59
3.9	A Feynman diagram with a gluon condensate contributing to the TFF. Contribution from this diagram vanishes.	59
3.10	Plots of our result for twist-six correction to the TFF $F_{\gamma\gamma^*\to\eta}(Q^2)$ for $h_q=0.0055$ (a), 0.0015 (b) and $-0.0025$ (c).	63
3.11	Comparison of our result for twist-six correction to the TFF $F_{\gamma\gamma^* \to \eta}(Q^2)$ (solid line, right scale) with twist-four correction [98] to the same (dashed line, left scale). Parameters used are the same as used in Table .3.2	64
3.12	Plots of our result for twist-six correction (solid lines) superimposed on the result obtained in Ref.[46] (a)(dashed line, plotted by us) for the TFF $F_{\gamma\gamma^*\to\eta}(Q^2)$ . The shaded area corresponds to the uncertainty in our result due to the variation of the input parameters as given in the text. In (b) the same results are compared with data from Refs. [93, 96]	64
3.13	Plots of our result for twist-six correction (solid lines) superimposed on the result obtained in Ref.[98] (a)(dashed line, plotted by us) for the TFF $F_{\gamma\gamma^* \to \eta}(Q^2)$ . The shaded area corresponds to the uncertainty in our result due to the variation of the input parameters as given in the text. In (b) the same results are compared with the data from Refs. [93, 96]	65
3.14	Comparison of our result for twist-six correction to the TFF $F_{\gamma\gamma^* \to \eta'}(Q^2)$ (solid line, right scale) with twist-four correction [98] to the same (dashed line, left scale). Parameters used are the same as used in Table. 3.2	65
3.15	Plots of our result for twist-six correction to the TFF $F_{\gamma\gamma^*\to\eta'}(Q^2)$ for $h_q=0.0055$ (a), 0.0015 (b) and $-0.0025$ (c).	66
3.16	Plots of our result for twist-six correction (solid lines) superimposed on the result obtained in Ref.[46](a) (dashed line, plotted by us) for the TFF $F_{\gamma\gamma^* \to \eta'}(Q^2)$ . The shaded area corresponds to the uncertainty in our result due to the variation of the input parameters as given in the text. In (b) the same results are compared with the data from Refs. [93, 96]	67
3.17	Plots of our result for twist-six correction (solid lines) superimposed on the result obtained in Ref. [98](a) (dashed line, plotted by us) for the TFF $F_{\gamma\gamma^*\to\eta'}(Q^2)$ . The shaded area corresponds to the uncertainty in our result due to the variation of the input parameters as given in the text. In (b) the same results are compared with the data from Refs. [93, 96]	68
3.18	Four-momenta flowing through the Feynman graphs is conserved at each vertex.	71