

List of Figures

1.1	Pickup reaction	3
1.2	Stripping	3
1.3	Knock-out reaction	3
1.4	Compound Nucleus Evaporation	4
1.5	Cross sections as function of l	6
1.6	The section of the Segre chart	8
1.7	Threshold Anomaly and Breakup Threshold Anomaly	9
1.8	Projectile breakup	11
1.9	Reaction functions, UFF and improved UFF(IMP-UFF)	12
1.10	Fusion barrier distributions-tightly and weakly bound nuclei.	15
1.11	Quarter point angle	17
2.1	Dispersion relation-The linear schematic model for $W(E)$	25
3.1	Schematic picture of FOTIA facility	42
3.2	Schematic picture of scattering chamber	44
3.3	A picture of scattering chamber	45
3.4	Schematic picture of detector telescopes	45
3.5	Block diagram of electronics associated with the detector telescope .	46
3.6	A typical two-dimensional spectrum of ΔE versus E_{total} for ${}^7\text{Li} + {}^{27}\text{Al}$	47

3.7	Elastic scattering angular distributions fitted with the optical model analysis using Woods-Saxon form of potential for the ${}^7\text{Li} + {}^{27}\text{Al}$ system.	48
3.8	Radius of sensitivity for real part of the potential	49
3.9	Radius of sensitivity for imaginary part of the potential	49
3.10	Energy ($E_{c.m.}$) dependence of the (a) real and (b) imaginary parts of Woods-Saxon potential at R_s (~ 8.0 fm) along with the dispersion relation calculation for the ${}^7\text{Li} + {}^{27}\text{Al}$ system	50
3.11	α -particle cross sections for different evaporation channels for ${}^7\text{Li} + {}^{27}\text{Al}$ system from the predictions of PACE code	53
3.12	α -particle energy spectra measured at various energies in ${}^7\text{Li} + {}^{27}\text{Al}$ reaction along with the predictions of PACE code	54
3.13	Fusion cross sections obtained in the present experiment along with the data from the literature and CCFULL predictions for the ${}^7\text{Li} + {}^{27}\text{Al}$ system	56
3.14	Comparison of fusion cross sections in the reaction of ${}^6,{}^7\text{Li}$ and ${}^9\text{Be}$ with ${}^{27}\text{Al}$ target at around the Coulomb barrier energies	56
3.15	Elastic scattering angular distributions along with the CDCC-CRC calculation for the ${}^7\text{Li} + {}^{27}\text{Al}$ system	59
3.16	Breakup and transfer cross sections from the coupled-channels calculations using FRESCO	60
3.17	Inelastic cross sections from the experimental data and the FRESCO calculations	60
3.18	Transfer cross sections from the experimental data and the FRESCO calculations	61
3.19	Comparison of fusion cross sections obtained from the coupled-channels calculations(CDCC-CRC) and present measurement	61

4.1	Schematic picture of BARC-TIFR pelletron facility	65
4.2	A typical two dimensional ΔE - E spectrum for the ${}^7\text{Li} + {}^{159}\text{Tb}$ system	69
4.3	Schematic picture of BARC-TIFR pelletron facility	70
4.4	Plot of radii of sensitivity for ${}^7\text{Li} + {}^{159}\text{Tb}$ system	71
4.5	Energy dependence of the real and imaginary parts of the optical potential obtained for ${}^7\text{Li} + {}^{159}\text{Tb}$ system	73
4.6	Reduced reaction cross sections for the systems that involve ${}^7\text{Li}$ and different target masses	74
4.7	Experimental elastic scattering cross sections normalized to the Rutherford cross sections as a function of $\theta_{c.m.}$ for ${}^7\text{Li} + {}^{159}\text{Tb}$ system	75
4.8	Real and imaginary parts of the DPP as a function of radius R at different energies for ${}^7\text{Li} + {}^{159}\text{Tb}$ system	75
4.9	Real and imaginary parts of the DPP at an average radius $R_s=11.5$ fm as a function of energy for ${}^7\text{Li} + {}^{159}\text{Tb}$ system.	76
5.1	A typical bi parametric spectrum of ΔE versus E_{total} in the reaction of ${}^7\text{Li}$ with ${}^{209}\text{Bi}$	80
5.2	A picture of scattering chamber	81
5.3	Quasi-elastic scattering excitation function and (b) corresponding fusion barrier distributions along with the CC calculations, for ${}^6\text{Li}$ + ${}^{209}\text{Bi}$ system at 160°	83
5.4	Quasi-elastic scattering excitation function and (b) corresponding fusion barrier distributions along with the CC calculations, for ${}^7\text{Li}$ + ${}^{209}\text{Bi}$ system at 160°	84
5.5	Quasi-elastic scattering excitation function for ${}^6\text{Li} + {}^{209}\text{Bi}$ system .	85
5.6	Quasi-elastic scattering excitation function for ${}^7\text{Li} + {}^{209}\text{Bi}$ system .	85
5.7	Fusion barrier distributions for ${}^6\text{Li} + {}^{209}\text{Bi}$ system	87

5.8	Fusion barrier distributions for ${}^7\text{Li} + {}^{209}\text{Bi}$ system	88
5.9	Transmission or capture probability (T_o) in the reaction of ${}^6\text{Li}$ with ${}^{209}\text{Bi}$	89
5.10	Transmission or capture probability (T_o) in the reaction of ${}^7\text{Li}$ with ${}^{209}\text{Bi}$	90
5.11	Quasi-elastic excitation function along with the results of 1n-stripping channel for ${}^6\text{Li} + {}^{209}\text{Bi}$ at $\theta_{lab} = 140^\circ$	95
5.12	Corresponding barrier distribution along with the results of 1n- stripping channel for ${}^6\text{Li} + {}^{209}\text{Bi}$ at $\theta_{lab} = 140^\circ$	95
5.13	Relative contribution of direct breakup and 1n-stripping channel in the reaction of ${}^6\text{Li}$ with ${}^{208}\text{Pb}$, ${}^{209}\text{Bi}$	96

List of Tables

3.1	Woods-Saxon potential parameters and total reaction cross sections obtained from the Optical Model analysis of the elastic scattering data for the ${}^7\text{Li} + {}^{27}\text{Al}$ system	51
3.2	Fusion cross sections for the ${}^7\text{Li} + {}^{27}\text{Al}$ system at various bombarding energies extracted from the measured α -particle spectra.	55
4.1	Optical potential parameters and total reaction cross sections ${}^7\text{Li} + {}^{159}\text{Tb}$ system	71
5.1	Inelastic excitations included in the Coupled Channels (CC) calculations	90
5.2	Total reaction ($\sigma_{react.}^{total}$), integrated breakup ($\sigma_{BU}^{inte.}$) and breakup cross sections at 140° ($\sigma_{BU}^{140^\circ}$) from the CDCC calculation and α cross sections at 140° ($\sigma_\alpha^{140^\circ}$) from the experiment for ${}^6\text{Li} + {}^{209}\text{Bi}$ system. . .	91
5.3	Total reaction ($\sigma_{react.}^{total}$), integrated breakup ($\sigma_{BU}^{inte.}$) and breakup cross sections at 140° ($\sigma_{BU}^{140^\circ}$) from the CDCC calculation and α cross sections at 140° ($\sigma_\alpha^{140^\circ}$) from the experiment for ${}^7\text{Li} + {}^{209}\text{Bi}$ system. . .	92
5.4	The excitations of ${}^{209}\text{Bi}$ included in the calculation	94