List of Figures

| 1.1 | Schematic diagram of fission process | 9 |
|-----|--|------|
| 1.2 | Potential energy diagram showing the double-humped fission barrier | 10 |
| 1.3 | Fission yield as a function of mass number for thermal and 14 MeV | |
| | neutron-induced fission of ²³⁵ U. | . 12 |
| | | |
| 2.1 | A schematic arrangement of the AHWR [1] | . 34 |
| 2.2 | A Schematic diagram of ADS. | 37 |
| 2.3 | A Schematic diagram of ADS for transmutations. | 38 |
| | | |
| 3.1 | The rolling mill at Target lab, BARC-TIFR Pelletron Facility, Mumbai | .46 |
| 3.2 | A schematic diagram BARC-TIFR Pelletron Facility. | 48 |
| 3.3 | Yields of fission products (%) as a function of mass number in the neutron-induce | ed |
| | fission of ²³² Th at average energy of 5.42 MeV along with the literature data | 55 |
| 3.4 | Yields of fission products (%) as a function of mass number in the neutron-induce | ed |
| | fission of ²³² Th at average energy of 7.75 MeV along with the literature data | . 56 |
| | | |
| 4.1 | A Schamatic diagram of Th-U fuel cycle | 61 |
| 4.2 | Six meter Proton Beam Irradiation set-up at BARC-TIFR Pelletron, Mumbai | .65 |
| 4.3 | Schematic diagram showing the arrangements used for neutron irradiation | . 68 |
| 4.4 | High Purity Germanium (HPGe) Detector at RCD lab, TIFR, Mumbai | .69 |
| 4.5 | Gamma ray spectrum of irradiated 232 Th showing the γ -ray energy of 233 Pa | . 71 |
| 4.6 | Gamma ray spectrum of irradiated 232 Th showing the γ -ray energy of 231 Th | .72 |
| 4.7 | Neutron spectrum from ⁷ Li(p,n) reaction irradiated with proton energy of 5.6 MeV | V |

| | calculated using EMPIRE computer code |
|------|--|
| 4.8 | Neutron spectrum from ⁷ Li(p,n) reaction irradiated with proton energy of 12 MeV |
| | calculated using EMPIRE computer code |
| 4.9 | Neutron spectrum from $^7\text{Li}(p,n)$ reaction at E_p =5.6 MeV calculated using the |
| | results of Meadows and Smith [55] |
| 4.10 | Extrapolated neutron spectrum in ⁷ Li(p,n) reaction at E _p =12 MeV obtained from |
| | neutron spectrum at Ep=10 MeV of C. H. Poppe et al. [60] |
| 4.11 | Neutron spectrum from 7 Li(p, n) reaction at E _p =7.8 MeV calculated using the |
| | results C. H. Poppe et al. [60] |
| 4.12 | Neutron spectrum from 7 Li(p, n) reaction at E _p =18 MeV calculated using the |
| | results of C. H. Poppe et al. [60] |
| 4.13 | Gamma ray spectrum of separated ²³³ Pa from neutron irradiated ²³² Th |
| 4.14 | Plot of experimental and evaluated 232 Th(n, γ) reaction cross-section as a function |
| | of neutron energy from 1 keV to 14 MeV. Experimental values from present work |
| | and from ref. [21-37] are in different symbols, whereas the evaluated and |
| | theoretical values from TALYS are in solid line of different colors |
| 4.15 | Plot of experimental and evaluated ²³² Th(n, 2n) reaction cross-section as a |
| | function of neutron energy from neutron energy 5 MeV to 20 MeV. Experimental |
| | values from present work and from refs. [38-46] are in different symbols, whereas |
| | the evaluated and theoretical values from TALYS are in solid lines with |
| | different colors |
| | |
| 5.1 | A schematic diagram of ²³³ U production from ²³² Th |
| 5.2 | Gamma-ray spectrum of radiochemically separated ²³³ Pa showing seven different |

| | γ- rays with their branching intensities with x-axis as Channel numbers112 |
|-----|--|
| 5.3 | Gamma-ray spectrum of radiochemically separated ²³³ Pa showing seven different |
| | γ- rays with their branching intensities with x-axis as Energy (keV)113 |
| 5.4 | Radioactive decay of the ²³³ Pa as a function of time |
| 5.5 | Fission tracks of ca. 15 μ m size with elliptical shapes from 233 Pa(2n _{th} , f), |
| | recorded and developed on the Lexan detector with 500x magnification. Figs. 5a |
| | and 5b correspond to different areas of the same slide and the image |
| | was taken with different microscopes using the same magnification. |
| | The size of the magnified view of the track area of the lexan detector shown in |
| | (a) and (b) are 0.00159 cm ² |
| 5.6 | Plot of experimental and theoretically ²³⁴ Pa(n, f) cross-section. Theoretical fission |
| | sections above 1 keV were calculated using the TALYS computer code version 1.2. |
| | The fission cross-section below 1 keV up to thermal energy was extrapolated by a |
| | normalized fit of the 1/v formula as shown in the figure |
| 5.7 | Plot of experimental and theoretically ²⁴² Am(n, f) cross-section. Theoretical |
| | fission cross - sections above 1 keV were calculated using the TALYS computer |
| | code version 1.2. The fission cross-section below 1 keV up to thermal energy was |
| | extrapolated by a normalized fit of the 1/v formula as shown |
| | in the figure |
| | |
| 6.1 | Typical γ -ray spectrum of irradiated natural Zr metal at thermal neutron energy in |
| | APSARA reactor showing the γ -ray energy of 95 Zr and 97 Zr |
| 6.2 | Photograph of Purnima Neutron Generator (300 KV DC electrostatic accelerator) at |
| | Bhabha Atomic Research Centre (B.A.R.C.), Mumbai |

| 6.3 | Schematic diagram showing the arrangement used for neutron irradiation134 |
|-----|---|
| 6.4 | Plot of experimental and evaluated $^{94}{\rm Zr}(n,\gamma)^{95}{\rm Zr}$ reaction cross-section as a function |
| | of neutron energy 1 MeV to 15 MeV. Experimental values from the present work |
| | and from ref.29 are in different symbols and colors, whereas the evaluated values |
| | from TENDL 2010 is in black solid line |
| 6.5 | Plot of experimental and evaluated $^{90}\text{Zr}(n,\gamma)^{90}\text{Y}^{\text{m}}$ reaction cross-section as a |
| | function of neutron energy 2 MeV to 20 MeV. Experimental values from the |
| | present work and from ref. (30, 31) are in different symbols and colors, |
| | whereas the evaluated values from TENDL 2010 are in black solid line141 |
| | |
| 7.1 | A schematic of regional IAEA-NDS nuclear data mirror site set-up in India147 |
| 7.2 | Creation of EXFOR files using EXFOR-Editor |
| 7.3 | ⁵⁶ Fe(n,p) cross-section experimental data with ENDF/B-VI evaluation153 |