

5. BIBLIOGRAPHY

Abdullah Harun, Chowdhury and Abdullah A Mamun 2006. Physico-chemical conditions and plankton population of two fishponds in Khulna. Univ. J. Zool. Rajshahi Univ. Vol. 25, pp. 41-44 ISSN 1023-6104.

Adoni, A.D. 1975. Studies on microbiology of Sagar Lake, Ph.D. Thesis, Sagar University, Sagar, India. APHA, 1998. Standard methods for examination of water and waste (20th ed.) American Public health Association. Washington D.C..

Ameen, M., Begum, Z.N.T., Ali, S., Rahman, M.M. & Roy, T.K. 1986. A Comparative limnological study of two fish ponds Raipur. Dhaka Uni. Stud. E 1 (1):25-34.

Anene, A., 2005. Condition factors of four cichlid species of a man-made lake in Imo state, Southeast, Nigeria. Turk. J. Fish. Aquat. Sci., 5: 43-47.

APHA 1998. Standard methods for examination of water and waste water (20ed.) American Public Health Association. Washington D.C.

APHA. 2004. Standard Methods for the Examination of Water and Waste Water. 20th Ed. American Public Health Association, Washington D. C.

Arthington A.H. and D.A. Milton. 1986. Reproductive biology, growth and age composition of the introduced *Oreochromis mossambicus* (Cichlidae) in two reservoirs, Brisbane, Australia. Environmental Biology of Fishes 16:257-266.

Bagenal T. B. and Tesch F. W. 1978. Age and growth. In: T.B. Bagenal, (ed) Methods for assessment of fish production in freshwater, 3rd edition. Blackwell Scientific Publication, Oxford, UK. 101–136.

Bakare, O., 1970. Bottom Deposits as Food of Inland Fresh Water Fish. In: Kainji, A Nigerian man-made lake. S.A. Visser, (Ed.), Kanji Lake Studies Vol. 1. Ecology Published for the Nigerian Institute.

Basavaraja, J. Narayana, et, al ; 2014. Fish diversity and abundance in relation to water quality of Anjanapura reservoir, Karnataka, India. International Journal

of Current Microbiology and Applied Sciences. ISSN: 2319-7706, Vol.3 No.3,pp.747-757.

Bhatt M., Yazdani T., Narian K., Yunus M., and Shukla R. 2009. Water quality status of some urban ponds of Lucknow, Uttarpradesh. Journal of Wetlands Ecology.vol-2, pp 67-73.

Bhuiyan, N.I.M.A.S, Nahar, Q. & Islam, M.N. 1997.Physico-chemical condition in relation to meteorological condition of a fish pond in Rajshahi. Univ. j. zool.Rajshahi Univ.16:85-88.

Boschung, H.T., and R.L Mayden. 2004. Mozambique Tilapia: *Oreochromis mossambicus* (Peters). Pp 620. In: Fishes of Alabama. Smithsonian Books. Washington D.C. 960 p.

Bowen S.H. 1979. A nutritional constraint in detritivory by fishes: The stunted population of *Sarotherodonmossambicus* in Lake Sibaya, South Africa. Ecological Monographs 49:17-31.

Boyd, C. E. and C. S. Tucker. 1982. Pond Aquaculture Water Quality Management. Kluwer Academic Publishers, Boston, Massachusett.

Boyd, C.E and Tucker, C.S. 1998. Pond aquaculture water quality management. Kluwer Academic Publishers, London.

Brown W. H. 1961. First Record Of The African Mouthbreeder *Tilapia mossambica* Peters In Texas. Texas Journal of Science 13:352-354.

Bruton M.N., and B.R. Allanson. 1974. The growth of *Tilapia mossambica* Peters (Pisces: Cichlidae) in Lake Sibaya, South Africa. Journal of Fish Biology 6:701-715.

Chowdhury, A.N., Begum, S. & Sultana, S. 1989.Occurrence of seasonal variation of zooplankton in fishpond in relation to some physco- chemical factors.Bangladesh j. zool.17(2):101-106.

Das A. K. 2000. Limmnochemistry of some Andhra Pradesh reservoirs. J. Inland. Fish Soc. India,32:37-44.

Das SK, Chand BK. 2003. Limnology and biodiversity of Ichthyofauna in a pond of Southern Orissa, India, Journal of Ecotoxicology and Environmental monitoring. 13(2): 97-102.

De Silva S.S., Perera M.K., and P. Maitipe. 1984. The composition, nutritional status and digestibility of the diets of *Sarotherodon mossambicus* from nine man-made lakes in Sri Lanka. Environmental Biology of Fishes 11:205-219.

Dial R.S. and S.C. Wainright. 1983. New distributional records for non-native fishes in Florida. Florida Scientist 46:8-15.

Fagade, S.O., 1979. Observation of the biology of two species of Tilapia from the Lagos lagoon Nigeria. Bull. Inst. Fond Afr. Nore (Ser. A), 41: 627-658.

Fakruzzaman, M., Chowdhury, A.H., Naz, S. & Zaman, M. 2001. Zooplankton of some fishponds in Barind Tract in relation to its physico-chemical variables. Univ. j. zool.

Forse, R., Pauly, D. (Eds.) (2011). FishBase. World Wide Web electronic publication. URL: www.fishbase.org <http://www.fishbase.org> [version 08/2011]

Fulton, T. W., The rate of growth of fishes. 22nd Annual Report of the Fishery Board of Scotland (3), 141-241 (1904).

Gupta M.V. and B.O. Acosta. 2004. A review of global tilapia farming practices. World Fish Center P.O. Box 500 GPO, 10670, Penang, Malaysia.

Harshey, D.K., Patil, G. and Singh, D.F. 1982. Limnological studies on a tropical fresh water fish tank of Jabalpur, India. Geobios New Reports 1 (2) : 98-102.

Hecky R.E., and Kilham P. 1988. Nutrient limitation of Phytoplankton in fresh water and marine environments: a review of recent evidence on the effects of enrichments Limnology and Oceanography 33: 796-832.

Hodgkiss I.J., and H.S.H. Man. 1978. Reproductive biology of *Sarotherodon mossambicus* (Cichlidae) In Plover Cove Reservoir, Hong Kong. Environmental Biology of Fish 3:287-292.

Hubbs C., Lucier T., Garrett G.P., Edwards R.J., Dean S.M., Marsh E., and D. Belk. 1978. Survival and abundance of introduced fishes near San Antonio, Texas. *Texas Journal of Science* 30:369-376.

IAAB, Hyderabad publication 31-82.

ICAR 2011. Hand book of fisheries and aquaculture. ICAR Publication New Delhi.

Jain SM, Sharma M and Thakur R 1996. Seasonal variation in physico – chemical parameters of Halali reservoir of Vidisha District, India. *Journal of Ecobiologia*. 8(3) 181 – 188.

Jana, B.B. 1973. Seasonal periodicity of plankton in a fresh water pond in West Bengal, India. *Int. Rev. Ges. Hydrobiol.* 58:127-143.

Jhingran, V.G. 1985. Fish and Fisheries of India. Hindustan Publishing Corporation (India), Delhi.

Keri AlhadiIghwela, Aziz Bin Ahmed and A.B. Abol-Munafi (2011). Condition Factor as an Indicator of Growth and Feeding Intensity of Nile Tilapia Fingerlings (*Oreochromis niloticus*) Feed on Different Levels of Maltose. *American-Eurasian J. Agric. & Environ. Sci.*, 11(4): 559-563, ISSN 1818-6769.

King, R.P. (1996). Length –Weight relationship of Nigeria fresh water fishes. *Naga ICLARM Quarterly*. 19 (4): 53-58.

Kodarkar MS 1995. Conservation of lakes. Indian Association of Aquatic Biologists

Korai, A.L., Sahato, G.A. Lashari, K.H., S.N. 2008. Biodiversity in Relation to Physicochemical Properties of Keenjhar Lake, Thatta District, Sindh, Pakistan. *Turkish Journal Of Fisheries and Aquatic Sciences* 8:259-268.

Kumar N. and Sharma R. C. 2000. Water Quality of river Krishna (Part-2 Biological Characteristics of 7 bioindicators) *J. Nature Conservator*, 14(2): 273-297.

Landau, M. 1992. Introduction to Aquaculture. John Wiley and Sons, Inc. New York.

Le Cren, E. D., 1951. The length weight relationship and seasonal cycle in gonad weight and condition of Perch (*Perca fluviatilis*), J. Anim. Eco., 21, 210-219.

Lee D.S., Gilbert C.R., Hocutt C.H., Jenkins R.E., McAllister D.E., and J.R. Stauffer, Jr. 1980. Atlas of North American Freshwater Fishes. North Carolina State Museum of Natural History, Raleigh, NC. 854 p.

Lett, P. F., and W. G. Doubleday. 1976. The influence of fluctuations in recruitment on fisheries management strategy. International Commission for the Northwest Atlantic Fisheries Selected Papers 1:171–193.

Mallasen M, Barros HP, Traficante DP, Camargo ALS. 2012. Influence of net cage tilapia culture on the water quality of the Nova Avanhandava reservoir, Sao Paulo state, Brazil. Acta Scientiarum Biological Sciences.; 34(3):289-296.

Manimegalai M.1, Karthikeyeni S 2 et al. 2010. Morphometric Analysis – A Tool to Identify the Different Variants in Fish Species *E. Maculatus*. Research article ISSN 0976-4402. 481.

Mauro Jose Cavalcanti, Leandro Rabello Monterio and Paulo Roberto Duarte Lopes 1999. Landmark- based Morphometric Analysis in selected species of Serranid Fishes (Perciformes:Teleostei). Zoological studies 38 (3): 287-294.

Mc Laughlin and Brindle R. L. 2001. A laboratory-scale investigation into the use of powdered iron for in situ: Removal of phosphorus from treated sewage effluent with constructed wetlands. Water Quality. Res. J. Canada. 36(4):759-779.

Mc Rae, B. J., and J. S. Diana. 2005. Factors influencing density of age-0 brown trout and brook trout in the Au Sable River, Michigan. Transactions of the American Fisheries Society 134:132–140.

Mishra, P.C., Dash, M.c., Choudhary, K and Das, R.C. 1988. Pollution studied in river Ib:Microbial analysis of water and sediments Book. “Ecology and Pollu. Of rivers” (Ed.Trivedi, R.K.). Ashish Pub. House, New Delhi, I:293-302.

Morey, G., Moranta, J., Massuti, E., Grau, A., Linde, M., Rivera, F. and Morales-Nin, B.2003.Weight-length relationships of littoral to lower slope fishes from the western Mediterranean. Fisheries Res., 62:89-96.

Murta, G. 2002. “Morphological variation of horse mackerel (*Trachurus trachurus*) in the Iberian and North African Atlantic: implications for stock identification,” J. Mar. Sci., vol. 57, 1240-1248.

Mustapha M. K. 2008. Assessment of water quality of Oyun Reservoir, Offa, Nigeria, and Using 2 selected Physic-chemical Parameters. Turkish Journal of Fisheries and Aquatic Sciences 8:309-319.

Naz, S. 1992. Studies on Physico-chemical Conditions and Plankton of Fishponds in Rajshahi. Unpubl. M.Sc.thesis, Dept. of Botany. Rajshahi University. 167 pp.

Odem, E.P. 1971. Fundamentals of ecology. 3rd Ed. W.B. Saunders Company Philadelphia, 578 pp.

Odum, E.P. 1984. Fundamentals of Ecology. 3rd Ed. W.B. Saunders Company Toronto.

Oni, S.K., J.Y. Olayemi and J.D. Adegboye, 1983. Comparative physiology of three ecologically distinct fresh water fishes, *Alestes nurse* (Ruppell), *Synodontis schall*(Bloch), *S. schneider* and *Tilapia zilli*(Gervais). J. Fish Biol., 22: 105-109.

Parikh, Ankita N. 2011. “Assessment of Urban Ponds with special Reference to Anthropogenic Activities”.

Pathak Neelam B. Parikh Ankita N. and Mankodi Pradeep C. 2013. Morphometric Analysis of Fish Population from two Different Ponds of Vadodara City, Gujarat, India. Research Journal of Animal, Veterinary and Fishery Sciences, Vol. 1(6), 6-9: ISSN 2320 – 6535.

Pervin, M.R. and M.G. Mortuza, 2008. Notes on length-weight relationship and condition factor of fresh water fish, *Labeoboga*(Hamilton) (Cpriniformes: Cyprinidae). Univ. J. Zool. Rajshahi Univ. 27: 97-98.

Radha, and Seenayya, G.2004. Influence of Physico-chemical Parameters on the distribution of heterophilic bacteria in a highly populated and eutrophic freshwater lake. Asian J. Of Microbiol. Biotech. Env. Sc. Vol.6, No.(3), 373-377.

Rath, R.1993. Fresh water Aquaculture. Scientific Publishers, Jodhpur.

Richards, L.A.1954. Diagnosis and improvement of saline and alkali soils. Hand book No.60, Oxford and IBH publication company, Calcutta-16.

Seenayya, G. 1971. Ecological studies in the Plankton of certain fresh water ponds of Hyderabad, India. II, Phytoplankton-I. Hydrobiol. 37: 35-88.

Shafland P.L. and J.M. Pestrak. 1982. Lower lethal temperatures for fourteen non-native fishes in Florida. Environmental Biology of Fishes 7:139-156.

Shastri, Y .and Pandse, D.C. 2001. Hydrobiological study of Dahikhuta reservoir J. Environment Biol. 22:67-70.

Singh M. 1965. Phytoplankton periodicity in small lake near Delhi (I) Seasonal fluctuations of the physico-chemical characteristics of the water. Phykos. 4:61-68.

Soni R. N and Bhatt S. A. 2008. Periodical Ecological Study of an urban pond

Stergiou, K.I. and D.K. Moutopoulos, 2001. A review of length-weight relationships of fishes from Greek marine waters. Naga, ICLARM Quart, 24(1-2): 23-39.

Swaranlatha, N. and Rao, A.N., 1998. Ecological studies of Banjara Lake with reference to water pollution J. Environment Biol. 19:179-186.

Trewevas E. 1983. Tilapiine Fishes Of The Genera *Sarotherodon*, *Oreochromis* And *Danakilia*. British Museum Of Natural History, Publication Number 878. Comstock Publishing Associates. Ithaca, New York. 583 p.

Tripathi, A.K. and Pandey. S. N. 1989. Studies of algae of populated ponds of Kanpur II. Qualitative, quantitative and periodical occurrence of the blue green algae of Chandori pond. Res. J Pl. Environ. 5(1): 89-93.

Unni K. S. 1985. Comparative limnology of several reservoirs of central India. Int. Review of Hydriobiol. 70(6):845-856.

Vivekananda, E. 2005. Stock assessment of tropical marine fishes, Indian Council of Agricultural Research, New Delhi, India, 115.

Weatherley , A. H., 1972. Growth and ecology of fish populations. Academic Press., London, 293p.

Welcome, R.L., 1979. Fisheries Ecology of Flood Plain Rivers. Longman Press, London, pp: 317.

Wells, J. V., and M. E. Richmond. 1995. Populations, metapopulations, and species populations: what are they and who should care? Wildlife Society Bulletin 23:458–462.

Wetch, P.S.1952. Limnology. Mc. Graw. Hill Book. Co., New York. 538pp.

Wetzel, R.G.!983. Limnology. Second edition Saunders college Publishing, USA. 767 PP.

Yadav, Y.S., Singh, R. K., Choudhary, M. and Kolekar, V.(1987). Limnology and productivity in Dighalibeel (Assam). Trop. Ecol. 28: 137-146.

Yazdandoost, M.Y. and Katdare, M.S.2000. Water quality index of major rivers in Pune. J. Industrial Pollution Control, 16: 231-238.