## List of Tables

| Table No. | Title   | Page No. |
|-----------|---|----------|
| Ι         | Pyrazosulfuron-Ethyl (PE) compound details  | 12       |
| 1.1       | The relation between concentration of PE and mortality rate of <i>O. mossambicus</i>  | 24       |
| 1.2       | LC50 values (mg/L) with their fiducial limits used in acute toxicity tests and for <i>O. mossambicus</i>  | 24       |
| 1.3       | Behavioural response of <i>O. mossambicus</i> in different concentration of PE up to 96 h.  | 26       |
| 1.4       | Haematological Parameter, Blood Glucose and Total Protein in <i>O. mossambicus</i> affected by acute exposure of PE:  | 27       |
| 2.1       | Changes in Haematological parameters during exposure and recovery treatment in <i>O. mossambicus</i> as a function of sublethal levels of PE  | 57       |
| 2.2       | Changes in Biochemical parameters during exposure and recovery treatment in <i>O. mossambicus</i> as a function of sublethal levels of PE   | 59       |
| 3.1       | Condition factor, HSI, SSI and GSI of <i>O. mossambicus</i> subjected to sub-acute concentrations of PE.  | 84       |
| 3.2       | Bonferroni comparisons for HSI, SSI of <i>O. mossambicus</i> subjected to sub-acute concentrations of PE.   | 85       |
| 3.3       | Bonferroni comparisons for Male GSI and Female GSI of <i>O. mossambicus</i> subjected to sub-acute concentrations of PE.  | 85       |
| 4.1       | Summary of the reaction index (Iorg rp), Organ index (I org), Total reaction index (I rp) and the total index (Tot-I) of the selected organ as calculated for the groups 1/20 <sup>th</sup> LC <sub>50</sub> , 1/10 <sup>th</sup> LC <sub>50</sub> and 1/5 <sup>th</sup> LC <sub>50</sub> for the 7 <sup>th</sup> days exposure of PE. (CD-circulatory disturbance, RC-regressive changes, PC-progressive changes, I- Inflammation) | 110      |
| 4.2       | Summary of the reaction index (Iorg rp), Organ index (I org), Total reaction index (I rp) and the total index (Tot-I) of the selected organ as calculated for the groups 1/20 <sup>th</sup> LC <sub>50</sub> , 1/10 <sup>th</sup> LC <sub>50</sub> and1/5 <sup>th</sup> LC <sub>50</sub> for the 14 <sup>th</sup> days exposure of PE. (CD-circulatory disturbance, RC-regressive changes, PC-progressive changes, I- Inflammation) | 110      |
| 4.3       | Gill tissue Histological assessment sheet of <i>O. mossambicus</i> on exposure of PE 7 <sup>th</sup> day  | 111      |
| 4.4       | Gill tissue Histological assessment sheet of <i>O. mossambicus</i> on exposure of PE 14 <sup>th</sup> day   | 112      |
| 4.5       | Liver tissue Histological assessment sheet of O. mossambicus on exposure of PE 7 <sup>th</sup> day  | 113      |
| 4.6       | Liver tissue Histological assessment sheet of <i>O. mossambicus</i> on exposure of PE 14 <sup>th</sup> day  | 115      |
| 4.7       | Kidney tissue Histological assessment sheet of <i>O. mossambicus</i> on exposure of PE 7 <sup>th</sup> day  | 117      |
| 4.8       | Kidney tissue Histological assessment sheet of <i>O. mossambicus</i> on exposure of PE 14 <sup>th</sup> day   | 119      |
| 5.1       | Frequency of Micronuclei and nuclear abnormalities counted in the species <i>O. mossambicus</i> exposed to PE at 7 <sup>th</sup> days   | 145      |
| 5.2       | Frequency of Micronuclei and nuclear abnormalities counted in the species <i>O. mossambicus</i> exposed to PE at 14 <sup>th</sup> days  | 145      |