

References

- Abele, L. G., 1974. Species diversity of decapod crustaceans in marine habitats. *Ecology*, 55: 156-161.
- Abida B., HariKrishna S. and Khan I., 2009. Analysis of heavy metals in water, sediments and fish samples of Madivala lakes of Bangalore, Karnataka. *Int. J. chem. tech. res.*, 1: 245-249.
- Adebisi, A. A., 1988. The physico-chemical hydrology of a tropical seasonal river Upper Ogun River. *Hydrologia*, 79: 159-165.
- Adekola, F. A. and O. A. A. Eletta, 2007. A study of heavy metal pollution of Asa River, Ilorin, Nigeria; Trace metal monitoring and geochemistry. *Environ Monit Assess*, 125: 157–163.
- Ajao, E. A. and S. O. Fagade, 1990. A study of sediment communities in Lagos Lagoon, Nigeria. *J. Oil Chem. Pollut.*, 7: 85-105.
- Allen, C. J., 2010. Ecology of the intertidal crab *Dotilla intermedia* from tsunami-impacted beaches in Thailand. Ph.D. Thesis, University of Southampton.
- Almeida, M. J., Flores, A. and Queiroga, H., 2008. Effect of crab size and habitat type on the locomotory activity of juvenile shore crabs *Carcinus maenas*. *Est. Coast. Shelf Sci.*, 80: 509–516.
- Andem, A. B., U. Udofia, K. A. Okorafor, J. A. Okete and A. A. A. Ugwumba, 2012. A study on some physical and chemical characteristics of Ona River Apata Ibadan South-west, Oyo State Nigeria. *Euro J Zool Res*, 1(2):37-46.
- Andersen L. E. and Melzer A., 2002. Intertidal crabs as potential bio monitors in Port Curtis, Centre for environmental management. Central Queensland University, Technical report 12.
- Anila Kumari, K. S., Abdul Azis, P. K., Natarajan. P., 2007. Water quality of the Adimalathura estuary, Southwest coast of India. *J Mar biol Ass India*, 49(1):1-6.
- Ansell, D. A., 1988. Migration or shelter? Behavioural options for deposit feeding crabs on tropical sandy shores. In: Chelazzi, G., Vannini. M.

- (Eds) Behavioural adaptations for life. Plenum Press, New York, p 15–26.
- Armonies, W., 1988a. Active emergence of meiofauna from intertidal sediment. *Mar Ecol Prog Ser*, 43: 151–159.
- Armonies, W., 1988b. Physical factors influencing active emergence of meiofauna from boreal intertidal sediment. *Mar Ecol Prog Ser*, 49: 277–286.
- Arya S., J.N. Trivedi and K.D. Vachhrajani, 2014. Brachyuran Crabs as a Biomonitoring tool: A Conceptual Framework for Chemical Pollution Assessment. *Int Res J Env Sci*, 3(1):49-57.
- Asprey W. P., 1971. Inter-species sexual discrimination and approach avoidance conflict in two species of fiddler crabs, *Uca pugnax* and *Uca pugilator*. *Animal Behaviour*, 19: 669-676.
- Attrill, M. J. and Rundle, S. D., 2002. Ecotone or ecocline: ecological boundaries in estuaries. *Estuar. Coast. Shelf Sci.*, 55: 929-936
- Avnimelech, Y., G. Ritvo, L. E. Meijer and M. Kochba, 2001. Water content, organic carbon and dry bulk density in flooded sediments. *Aquacultural Engineering*, 25: 25–33.
- Backwell, P. R. Y. and Jennions, M. D., 2004. Coalition among male fiddler crabs. *Nature*, 430: 417 pp.
- Backwell, P. R. Y. and Passmore, N. I., 1996. Time constraints and multiple choice criteria in the sampling behaviour and mate choice of the fiddler crab, *Uca annulipes*. *Behavioral Ecology and Sociobiology*, 38: 407-416.
- Backwell, P. R. Y., Jennions, M. D., Christy, J. H. and Passmore, N. I., 1998. Synchronized courtship in fiddler crabs. *Nature*, 391: 31-32.
- Bacon, M. R., 1971. Distribution and ecology of the crabs *Cyclograpsus lavauxi* and *C. insularum* in Northern New Zealand. *N. Z. J. mar. Freshwater Res.*, 5: 415–426.
- Balls, P. W., 1994. Nutrient inputs to estuaries from nine Scottish east coast rivers; influence of estuarine processes on inputs to the North Sea. *Estuar. Coast. Shelf Sci.*, 39: 329–352.

- Barnes, R. S. K., 1967. The osmotic behaviour of a number of grapsid crabs with respect to their differential penetration of an estuarine system. *J. exp. Biol.*, 47: 535–551.
- Barnes, R.D. and S. Hughes, 1988. *An Introduction to Marine Ecology*. 2nd Edn., Blackwell Scientific Publications, UK., pp: 351.
- Bartolini, F., Penha-Lopes, G., Limbu, S., Paula, J., and Cannicci, S., 2009. Behavioral Responses of the mangrove fiddler crabs (*Uca annulipes* and *U. inversa*) to urban sewage loadings: Result of a mesocosm approach. *Mar. Pollut. Bull.*, 58: 1860-1867.
- Beleem, I. B., Yogesh Kumar, J. S., Satyanarayana, C., Venkatraman, K., Kamboj, R. D., 2014. Distribution of Marine Crabs from the Marine National Park, Gulf of Kachchh. *Scholars Academic Journal of Biosciences*, 2(7): 419 – 427.
- Bellucci, L. G., El Moumni, B., Collavini, F., Frignani, M. and Al bertazzi, S., 2003. Heavy metals in Morocco Lagoon and river sediments. *Journal de Physique*, 107, 139–142.
- Bertini, G and Fransozo, A., 1999. Spatial and seasonal distribution of *Petrochirus diogenes* (Anomura, Diogenidae) in the Ubatuba Bay, São Paulo, Brazil. *Iheringia*, 86:145–50.
- Bertini, G., Fransozo, A. and Costa, R.C., 2001. Ecological distribution of three species of *Persephona* (Brachyura, Leucosiidae) in the Ubatuba region, São Paulo, Brazil. *Nauplius*, 9:31–42.
- Bertness, M. D., Leonard, G. H., 1997. The role of positive interactions in communities: lessons from intertidal habitat. *Ecology*, 78: 1976–1989.
- Betrolootto, R. M., Tortarolo, B., Frignani, M., Bellucci, L. G. and Albanese, S. C., 2003. Heavy metals in coastal sediments of the Ligurian Sea off Vado Ligure. *Journal de Physique*, 107: 159–162.
- Beukema, J. J., Honkoop, P. J. C. and Dekker, R., 1998. Recruitment in *Macoma balthica* after mild and cold winters and its control by egg production and shrimp predation. *Hydrobiologia*, 375/376: 23-34.
- Bezerra L. E. A., C. B. Dias, G. X. Santana and H. Cascon, 2006. Spatial distribution of fiddler crabs (genus *Uca*) in a tropical mangrove of northeast Brazil. *Scientia marina*, 70(4): 759-766.

- Billen, G. and M. Somville, 1982. An ecophysiological model of nitrification in the Scheldt estuary. *Mathematical Modelling*, 3: 523–533.
- Blackmore, G., 1996. Biomonitoring of heavy metal pollution in Hong Kong coastal waters, using barnacles. *Asi.Mar. Biol.*, 13: 1-13.
- Blackmore, G., B. Morton and Z. G. Huang, 1998. Heavy Metals in *Balanus amphitrite* and *Tetraclita squamosa* (Crustacea: Cirripedia) Collected from the Coastal Waters of Xiamen, China. *Mar. Pollut. Bull.*, 36(1): 32-40.
- Bolam, S. G., 2003. Spatial patterns of estuarine macro benthic assemblages: relationship with hydrodynamic regime. Estuarine Process Research Project, CEFAS, March 2003 Report.
- Bolam, S. G., and Whomersley, P., 2005. Development of Macrofaunal communities on dredged material used for mudflat enhancement: a comparison of three beneficial use schemes after one year. *Mar. Pollut. Bull.*, 50: 40–47.
- Botto, F., Iribarne, O., Gutierrez, J., Bava, J., Gagliardini, A., Valiela, I., 2006. Ecological importance of passive deposition of organic matter into burrows of the SW Atlantic crab *Chasmagnathus granulatus*. *Mar. Ecol. Prog. Ser.*, 312: 201–210.
- Brenchley, G. A. and Carlton, J.T., 1983. Competitive displacement of native mud snails by introduced periwinkle in the New England intertidal zone. *Biol. Bull.*, 165: 543– 558.
- Bromley R.G., 1996. Trace fossils, biology, taphonomy and application. Second Edition. Chapman and Hall.
- Brown, A. C. and A. McLachlan, 1990. Ecology of Sandy Shores. Elsevier Science Publishers, Amsterdam, The Netherlands.
- Bryan, G. W. and Langston, W. J., 1992. Bio-availability, accumulation and effects of heavy metals in sediments with special reference to United Kingdom estuaries: A review. *Environ. Pollut.*, 76(2): 89–131.
- Bryan, G. W. and L. G. Hummerstone, 1973. Adaptation of the polychaete *Nereis diversicolor* to estuarine sediments containing high concentrations of heavy metals: General observations and adaptations to copper. *J. Mar. Biol. Assn.*, 51: 845-863.

- Bu-Olayan, A. H. and B. V. Thomas, 2005. Validating species diversity of benthic organisms to trace metal pollution in Kuwait bay, off the Arabian Gulf. *Applied Ecology and Environmental Research*, 3(2): 93-100.
- Butler, S. and F.L. Bird, 2007. Estimating density of intertidal ghost shrimps using counts of burrow openings. Is the method reliable? *Hydrobiologia*, 589: 303-314.
- Caddy, J. F. and O. Defeo, 2003. Enhancing or restoring the productivity of natural populations of shellfish and other marine invertebrate resources. *FAO Fish Tech*, pp 448, FAO, Rome.
- Cadee, G.C., 2001. Sediment dynamics by bioturbating organisms. In: Reise, K. (Ed.), *Ecological Comparisons of Sedimentary Shores*. Springer, pp. 127–148.
- Cameron, A. M., 1966. Some aspects of the behaviour of the soldier crab *Mictyris longicarpus*. *Pac. Sci.*, 20: 224-234.
- Campbell, P.G.C. and A. Tessier, 1989. Geochemistry and bioavailability of trace metals in sediments, In, *Aquatic Ecotoxicology: Fundamental Concepts and Methodologies*, Volume I, CRC Press, Boca Raton, Florida.
- Cannicci, S., J. Paula and M. Vannini, 1999. Activity pattern and spatial strategy in *Pachygrapsus marmoratus* (Decapoda: Grapsidae) from Mediterranean and Atlantic shores. *Mar. Biol.*, 133: 429–435.
- Cardoso, I., J. P. Granadeiro and H. Cabral, 2010. Benthic macroinvertebrates' vertical distribution in the Tagus estuary (Portugal): The influence of tidal cycle. *Est. Coast. Shelf Sci.* 86: 580–586.
- Cardoso, P., Pardal, M. A., Lilleb, A. I., Ferreira, S. M., Rafaelli, D., and Marques, J. C., 2004. Dynamic changes in sea grass assemblages under eutrophication and implications for recovery. *J. Exp.Mar. Biol Ecol.*, 302: 233–248.
- Carr, S.D., Tankersley, R.A., Hensch, J.L., Forward, R.B., Luettich, R.A., 2004. Movement patterns and trajectories of Oviparous blue crabs *Callinectes sapidus* during the spawning migration. *Estuar.Coast. Shelf Sci.*, 60: 567–579.

- Casariego, A.M., J. Alberti, T. Luppi, P.Daleo, O. Iribarne, 2011. Habitat shifts and spatial distribution of the intertidal crab *Neohelice (Chasmagnathus) granulata* Dana. *Journal of Sea Research*, 66: 87–94.
- Chakrabarti, A., Chakrabarti, R. and Hertweck, G., 2006. Surface traces and bioturbate textures from bubbler crabs: an indicator of subtropical to tropical tidal flat environments. *Senckenbergiana Maritime*, 36: 19-27.
- Chandy, M., 1973. New records of brachyuran decapods from the Gulf of Kutch, J. Bombay Nat. Hist. Soc. 70 (2): 401-402.
- Chandy, M., 1969. *Thalamita poissonii* (Audouin & Savigny) de Man, (Crustacea: Brachyura). A new record to Indian coasts, J. Bombay Nat. Hist. Soc. 66 (3): 635-637.
- Chapman, M. G. and A. J. Underwood, 1996. Influences of tidal conditions, temperature and desiccation on patterns of aggregation of the high-shore periwinkle, *Littorina unifasciata*, in New South Wales, Australia. *J. Exp. Mar. Biol. Ecol.*, 196: 213–237.
- Charmantier, G., Giménez, L., Charmantier-Daures, M., Anger, K., 2002. Ontogeny of osmoregulation, physiological plasticity and larval export strategy in the grapsid crab *Chasmagnathus granulata* (Crustacea, Decapoda). *Mar. Ecol. Prog. Ser.*, 229: 185–194.
- Chelazzi, G. and M. Vannini, 1988. *Behavioral Adaptation to Intertidal Life*. Plenum Press, New York.
- Chelazzi, G., J.L. Deneubourg and S. Focardi, 1984. Cooperative interactions and environmental control in the intertidal clustering of *Nerita rextilis* (Gastropoda: Prosobranchia). *Behaviour*, 90: 151-166.
- Chhapgar B. F., 1957. On the marine crabs (Decapoda – Brachyura) of Bombay state. Part I. *J. Bombay Nat. Hist. Soc.*, 54: 399-439.
- Chhapgar B. F., 1958. More additions to the crab fauna of Bombay state. *J. Bombay Nat. Hist. Soc.*, 65(3): 608-617.
- Chhapgar, B. F. and Mundkur T., 1995. Occurrence of the crab *Euxanthus exsculptus* (Herbst) In Gujarat. *Journal Bombay Natural. Hist. Society*, 92: 286.
- Christy, J. H. 1988a. Pillar function in the fiddler crab *Uca beebei* (L): effects on male spacing and aggression. *Ethology*, 78: 53-71. .

- Christy, J. H. 1988b. Pillar function in the fiddler crab *Uca beebei* (II): competitive courtship signalling. *Ethology*, 78: 113-128.
- Christy, J.H. and M. Salmon. 1984. Ecology and evolution of mating systems of fiddler crabs (genus *Uca*). *Biological Reviews*, 59: 483-509.
- Christy, J.H., 1978. Adaptive significance of reproductive cycles in the fiddler crab *Uca pugilator*: a hypothesis. *Science*, 199:453–455.
- Christy, J.H., 1986. Timing of larval release by intertidal crabs on an exposed shore. *Bull Mar Sci*, 39:176–191.
- Christy, J.H., 1987. Competitive mating, mate choice and mating associations of brachyuran crabs. *Bull mar sci.*, 41 (2): 177-191.
- Clayton, D. A. and A. Al-Kindi, 1998. Population structure and dynamics of two scopimerine sand crabs *Scopimera crabriacauda* Alcock 1900 and *Dotilla sulcata* (Forskall 1775) in an estuarine habitat in Oman. *Tropical Zoology*, 11: 197-215.
- Clayton, D.A., 1988. Hood construction as a spacing mechanism in *Cleistostoma kuwaitense* (Crustacea: Ocypodidae). *Mar. Biol.*, 99: 57-61.
- Clutton-Brock, T. H. and S. D. Albon, 1979. The roaring of red deer and the evolution of honest advertisement. *Behaviour*, 69: 145-170.
- Connell, J.H., 1983. On the prevalence and relative importance of interspecific competition: evidence from field experiments. *Am. Nat.*, 122: 661–696.
- Connell, J. H., 1961. Effects of competition, predation by *Thais lapillus*, and other factors on natural populations of the barnacle *Balanus balanoides*. *Ecological Monographs*, 31:61–104.
- Cook, A., 1981. A comparative study of aggregation in *pulmonate* slugs (Genus *Limax*). *J. Anim. Ecol.*, 50: 703-714.
- Cordell, J., 1978. Swamp dwellers of Bahia. *Natural History*, 78:62-74
- Cordell, J., 1989. Social marginality and sea tenure in Bahia. A sea of small boats. Edited by: Cordell J. Cambridge: Cultural Survival Inc.; pp. 125-151.
- Costa, R. C., Fransozo, A. and Castilho, A. L. and Freire, F. A. M., 2005. Annual, seasonal and spatial variation of abundance of the shrimp *Artemesia longinaris* (Decapoda: Penaeoidea) in south-eastern Brazil.

- Journal of the Marine Biological Association of the United Kingdom, 85:107–12.
- Costanza, R., R. d'Arge, R. de Groot, S. Farber, M. Grasso, B. Hannon, K. Limburg, S. Naeem, R. V. O'Neill, J. Paruelo, R. G. Raskin, P. Sutton and M. van den Belt, 1997. The value of the world's ecosystem services and natural capital. *Nature* 387: 253–260.
- Costanza, R., W. M. Kemp and W. R. Boynton, 1993. Predictability, scale, and biodiversity in coastal and estuarine ecosystems: implications for management. *Ambio*, 22: 88–96.
- Crabtree, R. E., 1995. Relationship between lunar phase and spawning activity of Tarpon, *Megalops atlanticus*, with notes on the distribution of larvae. *Bull Mar Sci*, 56:895–899.
- Crane, J., 1975. Fiddler crabs of the world. Ocypodidae: Genus *Uca*. Princeton University Press, Princeton.
- Crothers J. H., 1968. The biology of the shore crab *Carcinus maenas* (L). II. The life of the adult crab. *Field Stud.*, 2: 579-614.
- Cunningham, P. A and Myers, L. E., 1987. Effects of Diflubenzurin (Dimilin) on Survival, Molting, and Behavior of Juvenile Fiddler Crabs, *Uca pugilator*. *Archives of Environmental Contamination and Toxicology*, 16: 745-752.
- Daborn, G. R., Amos, C. L., Brylinsky, M., Cristian, H., Drapeau, G., Faas, R. W., Grant, J., Long, B., Paterson, D. M., Perillo, G. M. E., Piccolo, M. C., 1993. An ecological cascade effect: migratory birds affect stability of intertidal sediments. *Limnology and Oceanography*, 38: 225–231.
- Dauer, D. M., 1993. Biological criteria, environmental health and estuarine macrobenthic community structure. *Mar. Pollut. Bull.*, 26: 249-257.
- Dauvin, J.C., 2007. Paradox of estuarine quality: benthic indicators indices, consensus or debate for the future. *Mar. Pollut. Bull.*, 55: 271–281.
- Davies, N. B., and T. R. Halliday, 1978. Deep croaks and fighting assessment in toads *Bufo bufo*. *Nature*, London, 274: 683-685.
- Day, J. W., C. A. S. Hall, W. M. Kemp and A. Yanez-Arancibia, 1989. *Estuarine Ecology*. John Wiley & Sons, New York, NY. 558 pp.

- Dayton, P. K., 1971. Competition, disturbance and community organization: the provision and subsequent utilization of space in a rocky intertidal community. *Ecol. Monogr.*, 41: 351– 389.
- deBruyn, A. M. H. and J. J. Meeuwig, 2001. Detecting lunar cycles in marine ecology: periodic regression versus categorical ANOVA. *Mar Ecol Prog Ser*, 214: 307–310.
- Defeo, O. and McLachlan, A., 2005. Patterns, processes and regulatory mechanisms in sandy beach macrofauna: a multi-scale analysis. *Mar Ecol Prog Ser*, 295: 143-153.
- deRivera, C. E., 2005. Long searches for male-defended breeding burrows allow female fiddler crabs, *Uca crenulata*, to release larvae on time. *Animal Behaviour*, 70: 289-297.
- Depledge, M. H., 1984. Disruption of circulatory and respiratory activity in shore crabs *Carcinus maenas* (L.) exposed to heavy metal pollution. *Comp. Biochem. Physiol.*, 78C: 445-459.
- Deshkar, S., Lakhapurkar, J. and Gawali, D., 2012. State of three estuaries of Gulf of Khambhat. *Indian Journal of GeoMarine Science*, 41(1): 70-75.
- Dev Roy, M. K., 2013. Diversity and distribution of marine brachyuran crab communities inhabiting west coast of India: in *Ecology and Conservation of Tropical Marine Faunal Communities Part 1*. (edt. K. Venkatraman, C. Sivaperuman and C. Raghunathan) Springer Berlin Heidelberg, pp 147 – 169.
- Dexter, A. R., 2004. Soil physical quality Part I. Theory, effects of soil texture, density and organic matter and effects on root growth. *Geoderma*, 120: 201-214.
- Edgar, G. J. and Barrett, N. S., 2000. Effects of catchment activities on macrofaunal assemblages in Tasmanian estuaries. *Estuar Coast Shelf Sci*, 50: 639-54.
- Elliott, M. and P. F. Kingston, 1987. The sublittoral benthic fauna of the estuary and Firth of Forth, Scotland. *Proceedings of the Royal Society of Edinburgh*, 93B: 449-465.

- Elliott, M., Whitfield, A.K., Potter, I.C., Blaber, S.J., Cyrus, D.P., Nordlie, N.G., Harrison, T.D., 2007. The guild approach to categorizing estuarine fish assemblages: a global review. *Fish and Fisheries*, 8: 241–268.
- Emery, K. O. and Stevenson, R. E., 1957. Estuaries and lagoons. I: Physical and chemical characteristics. In *Treatise on Marine Ecology and Paleoecology*, Vol. 1, (J. W. Hedgpeth, ed.). The Geological Society of America. pp.673-750.
- Engle, V. D., Summers, J. K. and Gaston, G. R., 1994. A benthic index of environmental condition of Gulf of Mexico estuaries. *Estuaries*, 17(2): 372–384.
- Ens, B. J., M. Klaassen and L. Zwarts, 1993. Flocking and feeding in the fiddler crab (*Uca tangeri*): prey availability as risk-taking behaviour. *Netherlands J. Sea Res.*, 31: 477-494.
- Escapa, M., Iribarne, O. and Navarro, D., 2004. Indirect effect of intertidal burrowing crabs on infaunal zonation patterns, tidal behavior and risk of mortality. *Estuaries*, 27: 132 – 146.
- Farrington, J. W., E. D. Goldberg, R. W. Risebrough, J. H. Martin and V. T. Bowen, 1983. US Mussel Watch 1976-1978; an overview of the trace-metal, DDE, PCB, hydrocarbon, and artificial radionuclide data. *Env. Sci. Technol.*, 17: 490-496.
- Feare, C.J., 1971. The adaptive significance of aggregation behaviour in the dog whelk *Nucella lapillus* (L.). *Oecologia*, 7: 117-126.
- Fielder D.R., 1970. The feeding behaviour of the sand crab *Scopimera inflata* (Decapoda, Ocypodidae). *J. Zoo. Lond*, 160: 35-49.
- Flemming, B. W. and Delafontaine, M.T., 2000. Mass physical properties of muddy intertidal sediments: some applications, misapplications and non-applications. *Cont. Shelf Res.*, 20: 1179 – 1197.
- Flores, A. A. V, Abrantes, K. G. and Paula, J., 2005. Estimating abundance and spatial distribution patterns of the bubble crab *Dotilla fenestrata* (Crustacea: Brachyura). *Aus Ecol*, 30: 14-23.
- Flores, A. A. V. and J. Paula, 2001. Intertidal distribution and species composition of brachyuran crabs at two rocky shores in Central Portugal. *Hydrobiologia*, 449: 171–177.

- Folk, R.L. and W.C. Ward, 1957. Brazos River bar: a study in the significance of grain size parameters. *Journal of Sedimentary Petrology*, 27: 3–26.
- Fujii, T., 2007. Spatial patterns of benthic macrofauna in relation to environmental variables in an intertidal habitat in the Humber estuary, UK: Developing a tool for estuarine shoreline management *Est. Coast. Shelf Sci.*, 75: 101-119.
- Gadhavi, M. K. and K. D. Vachhrajani, 2015. Macrobenthos interactions and their distribution on lower estuarine mudflats of Mahi River, Gujarat, India. *Electronic Journal of Environmental Sciences*, 8: 49-56.
- Gallep, G.W., J.F. Kitchell and S.M. Bartell, 1978. Phosphorus release from Lake Sediments as affected by chironomid. *Ver inter vere for Limnologic*, 20: 458-465.
- George, A. D. I., J. F. N. Abowei and E. R. Daka, 2009. Benthic macro invertebrate fauna and physico-chemical parameters in Okpoka Creek sediments, Niger Delta, Nigeria. *International Journal of Animal and Veterinary Advances*, 1: 59–65.
- George, S. G., Pirie, B. J. S., Cheyne, A. R., Coombs, T. L. and P. T. Grant, 1978. Detoxification of metals by marine bivalves: ultrastructural study of the compartmentation of copper and zinc in the oyster, *Ostrea edulis*. *Mar. Biol.*, 45: 147-156.
- Gherardi, F., Russo, S., Lazzara, L., 2002. The function of wandering in the sand-bubbler crab, *Dotilla fenestrata*. *J Crust Bio.*, 22: 521-531.
- Gibson, G.R., M.L. Bowman, J. Gerritsen and B. D. Snyder, 2000. Estuarine and Coastal Marine Waters: Bioassessment and Biocriteria Technical Guidance. U.S. Environmental Protection Agency, Office of Water, Washington, DC.
- Gimenez, L., 2004. Marine community ecology: importance of trait-mediated effects propagating through complex life cycles. *Mar. Ecol. Prog. Ser.* 283, 303–310.
- Gohil, M. N. and P. C. Mankodi, 2013. Diversity of Fish Fauna from Downstream Zone of River Mahisagar, Gujarat State, India. *Res. J. Animal, Veterinary and Fishery Sci.*, 1(3): 14-15.

- Goldberg, E. D., M. Koide, V. Hodge, A. R. Flegal and J. Martin, 1983. US Mussel Watch: 1977-1978 results on trace metals and radionuclides. *Estuar. Coastal Shelf Sci.* 16, 69-93.
- Goldberg, E. D., V. T. Bowen, J. W. Farrington, G. Harvey, J. H. Martin, P. L. Parker, R. W. Risebrough, W. Robertson, E. Schneider and E. Gamble, 1978. The mussel watch. *Environ. Conserv.*, 5:101-125.
- Gor A. and A. Shah, 2014. Water quality index of Mahi River, Vadodara, Gujarat. *Inter. J. Engg. Devel. Res.*, 2(3): 3214-3219.
- Griffin, D. J. G., 1971. The ecological distribution of grapsid and ocypodid shore crabs (Crustacea: Brachyura) in Tasmania. *J. anim. Ecol.*, 40: 597–621.
- Griffis, R.B. and Suchanek, T.H. 1991: A model of burrow architecture and trophic modes in *thalassinidean* shrimp (Decapoda: Thalassinidea). *Mar. Ecol. Prog. Ser.*, 39: 171-183.
- Gross, W. J., 1964. Trends in water and salt regulation among aquatic and amphibious crabs. *Biol. Bull.*, 127: 447–466.
- Hakansson, I. and J. Lipiec, 2000. A review of the usefulness of relative bulk density values in studies of soil structure and compaction. *Soil Tillage Res.*, 53 (2000): 71–85.
- Halperin, J., Ansaldo, M., Pellerano, G.N., Luquet, C.M., 2000. Bimodal breathing in the estuarine crab *Chasmagnathus granulatus* Dana 1851. Physiological and morphological studies. *Comp. Biochem. Physiol.* 126: 341–349.
- Hammerstein, P. and G. A. Parker, 1982. The asymmetric war of attrition. *J. theor. Biol.*, 96: 647-682.
- Hartnoll, R. G., 1975. The Grapsidae and Ocypodidae (Decapoda: Brachyura) of Tanzania. *J. zool. Lond.*, 177: 305–328.
- Hartnoll, R. G., 1973. Factors affecting the distribution and behaviour of the crab *Dotilla fenestrata* on East African shores. *Est Coast Mar Sci*, 1: 137-152
- Hayes, C. L., I. Booksmythe, M. D. Jennions and P. R. Y. Backwell, 2013. Does male reproductive effort increase with age? Courtship in fiddler crabs. *Biol Lett*, 9: 20121078.

- Hilmy, A. M., N. E. A. El-Hamid and K. S. Ghazaly, 1988. Biochemical and physiological changes in the tissues and serum of both sexes in *Portunus pelagicus* (L.) following acute exposures to zinc and copper. *Folia Morphologica*, 36:79-94.
- Hines, A.H., Lipcius, R.N., Haddon, A.M., 1987. Population dynamics and habitat partitioning by size, sex and molt stage of blue crabs *Callinectes sapidus* in a subestuary of central Chesapeake Bay. *Mar. Ecol. Prog. Ser.* 36, 55–64.
- Hiroki, K. A. N., Fransozo, A., Costa, R. C., Castilho, A. L., Shimizu, R. M. and Almeida, A. C., 2011. Bathymetric distribution of the shrimp *Rimapenaeus constrictus* (Stimpson, 1874) (Decapoda, Penaeidae) in two locations off the south eastern Brazilian coast. *Mar. Biol. Res.*, 7:176–85.
- How, M. J., Zeil, J. and J. M. Hemmi, 2007. Differences in context and function of two distinct waving displays in the fiddler crab, *Uca perplexa* (Decapoda: Ocypodidae). *Behav Ecol Sociobiol*, 62:137-148.
- Hsiao-Chien Y., Chen I. M., Chen P. and Wang W. H., 2009. Heavy metal concentrations of the soldier crab (*Mictyris brevidactylus*) along the inshore area of Changhua, Taiwan. *Environ. Monit. Assess.*, 153: 103-109.
- Hunt H. and Scheibling, R., 1997. Role of post settlement mortality in recruitment of benthic marine invertebrates. *Mar. Ecol. Prog. Ser.*, 155: 269–301.
- Huntingford, F. A. and A. K. Turner, 1987. *Animal conflict*: 1-441. Chapman and Hall, London.
- Hyatt, G. W. and M. Salmon, 1978. Combat in the fiddler crabs *Uca pugilator* and *U. pugnax*: a quantitative analysis. *Behaviour*, 65: 182-211.
- Idowu, E.O. and A. A. A. Ugwumba, 2005. Physical, chemical and benthic faunal characteristics of a Southern Nigeria Reservoir. *The Zoologist*, 3: 15-25.
- Ikomi, R.B., F.O. Arimoro and O. K. Odihirin, 2005. Composition, distribution and abundance of macroinvertebrates of the Upper Reaches of River Ethiopie Delta State, Nigeria. *The Zoologist*, 3: 68-81.

- Iribarne, O., Martinetto, P., Schwindt, E., Botto, F., Bortolus, A. and B. P. Garcia, 2003. Evidences of habitat displacement between two common soft-bottom SW Atlantic intertidal crabs. *J Exp Mar Biol Ecol*, 296:167–182.
- Jain, C. K., Singhal, D. C. and Sharma, U. K., 2005. Metal pollution assessment of sediment and water in the river Hindon, India. *Environmental Monitoring and Assessment*, 105: 193–207.
- Jaroensutasinee, M. and P. Tantichodok, 2002. Effects of Size and Residency on Fighting Outcomes in the Fiddler Crab, *Uca vocans hesperiae* (Decapoda, Brachyura, Ocypodidae). *Crustaceana*, 75(9): 1107-1117
- Jennions, M. D. and Backwell, P. R. Y., 1998. Variation in courtship rate in the fiddler crab *Uca annulipes*: is it related to male attractiveness? *Behavioral Ecology*, 9: 605-611.
- Jennions, M.D. and Backwell, P.R.Y., 1996. Residency and size affect fight duration and outcome in the fiddler crab *Uca annulipes*. *Biological Journal of the Linnean Society*, 57: 293-306.
- Johannesson, L.T., Stevens, R.L. and Alexanderson J.H., 2000. Sediment Character in a Micro-Tidal, Harbor-Estuary Environment, Goteborg, Sweden, *Estuaries*, 23(3): 400-410.
- Jones, G., 1994. Global warming, sea level change and the impact on estuaries. *Mar. Pollut. Bull*, 28: 7-14.
- Jones, M. B., 1976. Limiting factors in the distribution of intertidal crabs (Crustacea: Decapoda) in the Avon-Heathcote estuary, Christchurch. *N. Z. J. mar. Freshwater Res.*, 10: 577–87.
- Jones, S.E., Jago, C.F., 1993. In situ assessment of modification of sediment properties by burrowing invertebrates. *Mar. Biol.*, 115: 133–142.
- Karbassi, A. R, Monavari, S. M., Nabi Bidhendi, G. R., Nouri, J. and Nematpour, K., 2008. Metal pollution assessment of sediment and water in the Shur River. *Environ. Monit. Assess. journal*, 147: 107–116.
- Karbassi, A. R., Nouri, J., and Ayaz, G. O., 2007. Flocculation of Cu, Zn, Pb and Ni during mixing of Talar river water with the Caspian seawater. *International Journal of Environmental Research*, 1: 66–73.

- Kellmeyer, K. and Salmon, M., 2001. Hatching rhythms of *Uca thayeri* Rathbun: timing in semidiurnal and mixed tidal regimes. *J Exp Mar Biol Ecol*, 260:169–183.
- Kennish, M. J., 2002. Environmental threats and environmental future of estuaries. *Environmental Conservation*, 29: 78–107.
- Ketpadung, R. and N. Tangkrock-olan, 2006. Changes in oxygen consumption and heart rate of the blue swimming crab, *Portunus pelagicus* (Linnaeus, 1766) following exposure to sublethal concentrations of copper. *Journal of Environmental Biology*, 27(1): 7-12.
- Key, R.S., 1983. Ecology of the infauna of Spurn Bight mudflats: an area proposed for reclamation. Ph.D. Thesis, University of Hull.
- Khan, A. S., Raffi, S. M. and Lyla, P. S., 2005. Brachyuran crab diversity in natural (Pitchavaram) and artificially developed mangroves (Vellar estuary). *Curr. Sci.*, 88: 1316-1324.
- Klerks, P. L. and J. S. Weis, 1987. Genetic adaption to heavy metals in aquatic organisms: A review. *Environ. Pollut.*, 45: 173- 190.
- Kim, T. W and Choe, J. C., 2003. The effect of food availability on the semilunar courtship rhythm in the fiddler crab *Uca lactea* (de Haan) (Brachyura: Ocypodidae). *Behavioral Ecology and Sociobiology*, 54: 210-217.
- King, R. S., Hines, A. H., Craige, F. D., and Grap, S. 2005. Regional, watershed and local correlates of blue crab and bivalve abundances in subestuaries of Chesapeake Bay, USA. *J. Exp. Mar. Biol. Ecol.*, 319: 101–116.
- Klause, M. D., 1986. Mucous secretions of the acoel *turbellarian* *Convoluta Orsted*: an ecological and functional approach. *Journal of Experimental Mar. Biol. Ecol.*, 97: 123–133.
- Klavins, M., Briede, A., Parele, A., Rodinov, V. and Klavina, I., 1998. Metal accumulation in sediments and benthic invertebrates in lakes of Latvia. *Chemosphere*, 36 (15): 3043-3053.
- Klumpp, D. W. and C. Burdon-Jones, 1982. Investigations of the potential of bivalve molluscs as indicators of heavy metal levels in tropical marine waters. *Aust. J. Mar. Freshwat. Res.*, 33: 285-300.

- Koga, T., 1995. Movements between microhabitats depending on reproduction and life history in the sand-bubbler crab *Scopimera globosa*. Mar. Ecol. Prog. Ser., 117:64-75.
- Korringa, P., 1947. Relations between the moon and the periodicity in the breeding of marine animals. Ecol Monogr, 17:349–381.
- Kosuge T, Poovachiranon S, Murai M, 1994. Male courtship cycles in three species of tropical Ilyoplax crabs (Decapoda, Brachyura, Ocypodidae). Hydrobiologia, 285: 93-100.
- Krichman, D., Mitchell, R., 1982. Contribution of particle-bound bacteria to micro heterotrophic activity in five coastal ponds and two marshes. Appl. Environ. Microbiol. 43, 200–209.
- Lauenstein, G. G., A. Robertsou and T. P. O'Connor, 1990. Comparison of trace metal data in mussels and oysters from a mussel watch programme of the 1970s with those from a 1980s programme. Mar. Pollut. Bull., 21: 440-447.
- Leimar, O. and M. Enquist, 1984. The effect of asymmetries in owner-intruder interactions. Journ. Theor. Biol., 111: 475-491.
- Levin, L.A., 1981. Dispersion, feeding behavior and competition in two spionid polychaetes. J. Mar. Res., 39: 99– 117.
- Lim, S. S. L., 2004. A comparative study of some mouthpart adaptations of *Uca annulipes* (H. Milne Edwards, 1837) and *U. vocans* (Linnaeus, 1758) (Brachyura, Ocypodidae) in relation to their habitats. Crustaceana, 77(10): 1245-1251.
- Lima, P. A., V. Fransozo, L. S. Andrade, A. C. Almeida, M. Furlan and A. Fransozo, 2014. Distribution and population structure of the flecked box crab *Hepatus pudibundus* (Decapoda, Brachyura) in the western South Atlantic, Mar. Biol. Res., 10:6: 589-600, DOI: 10.1080/17451000.2013.841940.
- Lipcius, R., Rochelle, R. D., Seebo, M. S., Colón-Carrión, D., 2005. Density, abundance and survival of the blue crab in sea grass and unstructured salt marsh nurseries of Chesapeake Bay. J. Exp. Mar. Biol. Ecol., 319: 69–80.

- Little, C. and L. Smith, 1980. Vertical zonation on rocky shores in the Severn Estuary. *Estuar. coast mar. Sci.*, 11: 651–669.
- Luoma, S. N., 1996. The developing framework of marine ecotoxicology: Pollutant as a variable in marine ecosystems? *J. Exp Mar Biol Ecol*, 200, 29-55.
- Luoma, S. N., 1989. Can we determine the biological availability of sediment-bound trace metals? *Hydrobiologia*, 176/177: 379–396.
- Luoma, S. N., D. J. Cain, K. Ho and A. Hutchinson, 1983. Variable tolerance to copper in two species from San Francisco Bay. *Mar. Environ. Res.*, 10: 209-222.
- MacIntosh, D. J., 1989. The ecology and physiology of decapods of mangrove swamps. *Proceedings of the Zoological Society of London*, 59: 315–341.
- McLusky, D. S., Bryant, V., Campbell, R., 1986. The effects of temperature and salinity on the toxicity of heavy metals to marine and estuarine invertebrates. *Oceanography and Biology, an Annual Review*, 24: 481–520.
- Malins, D.C. and G. K. Osdrander, 1994. *Aquatic Toxicology, Molecular, Biochemical and Cellular Perspectives*, Lewis Publishers, Boca Raton, Florida.
- Mantelatto, F. L. M. and Fransozo, A., 1999. Characterization of the physical and chemical parameters of Ubatuba Bay, northern coast of São Paulo State, Brazil. *Revista Brasileira de Biologia*, 59:23–31.
- Marden, J. H. and J. K. Waage, 1990. Escalated damselfly territorial contests are energetic wars of attrition. *Anim. Behav.*, 39: 954-959.
- Marden, J. H. and R. A. Rollins, 1994. Assessment of energy reserves by damselflies engaged in aerial contests for mating territories. *Anim. Behav.*, 48: 1023-1030.
- Martin, G. D., Vijay, V. J., Laluraj, C. M., Madhu, N. V., Joseph, T., Nair, M., 2008. Fresh water influence on nutrient stiochiometry in a tropical estuary, South West coast of India. *Applied Ecology and Environmental Research*, 6(1): 57–64.

- Martinetto, P., M. Valinas, Palomo, G. and Iribarne, O.O., 2007. Negative interactions between two SW Atlantic intertidal crabs in soft-bottom habitats. *Mar. Biol.*, 151: 1479-1490.
- Matsumasa, M. and Murai, M., 2005. Changes in blood glucose and lactate levels of male fiddler crabs: effects of aggression and claw waving. *Animal Behaviour*, 69: 569- 577.
- Maynard Smith, J. and G. Parker, 1976. The logic of asymmetric contests. *Anim. Behav.*, 24: 159-175.
- McIntyre, A.D., 1970. The range of biomass in intertidal sand, with special reference to the bivalve *Tellina tenuis*. *Journal of Biological Association of the UK*, 50: 561-575.
- McKillup, S. C., 1983. A behavioural polymorphism in the marine snail *Nassarius pauperatus*: geographic variation correlated with food availability and differences in competitive ability between morphs. *Oecologia*, 56: 58-66.
- McLay, C. L. and D. J. McQueen, 1995. Intertidal zonation of *Cyclograpsus lavauxi* H. Milne Edwards, 1853 (Brachyura: Grapsidae) along the coast of the South Island of New Zealand. *Crust. Res.*, 24: 49–64.
- McLusky, D. S., V. Bryant and R. Campbell, 1986. The effects of temperature and salinity on the toxicity of heavy metals to marine and estuarine invertebrates. *Oceanography and Biology, an Annual Review*, 24: 481–520.
- Meire, P., Ysebaert, T., Damme, S.V., Bergh, E.V.D., Maris, T. and Struyf, E., 2005. The Scheldt estuary: a description of a changing ecosystem. *Hydrobiologia*, 540:1–11.
- Meire, P.M., Seys, J.J., Ysebaert, T. J. and Coosen, J., 1991. A comparison of the macrobenthic distribution and community structure between two estuaries in SW Netherlands. In: Elliott, M., Ducrotoy, J.P. (Eds.), *Estuaries and Coasts: Spatial and Temporal Intercomparisons*. Olsen and Olsen, Fredensborg, pp. 221-230.
- Menendez, R. J., 1987. Vertical zonation of the Xanthidae mud crabs *Panopeus obesus* and *Panopeus simpsoni* on oyster reefs. *Bull. mar. Sci.*, 40: 73–77.

- Menge, B.A., 1976. Organization of the New England rocky intertidal community: role of predation, competition and environmental heterogeneity. *Ecol. Monogr.*, 46: 335– 393.
- Miller, D. C, 1961. The feeding mechanism of fiddler crabs, with ecological considerations of feeding adaptations. *Zoologica*, 46: 89-101.
- Milner, R.N.C., Booksmythe, I., Jennions, M.D., Backwell, P.R.Y., 2010. The battle of the sexes? Territory acquisition and defence in male and female fiddler crabs. *Anim Behav*, 79:735–738.
- Mizushima, N., Nakashima, Y. and Kuwamura, T., 2000. Semilunar spawning cycle of the humbug damselfish *Dascyllus aruanus*. *J Ethol*, 18:105–108.
- Moksnes, P.O., 2004. Self-regulating mechanisms in cannibalistic populations of juvenile shore crabs *Carcinus maenas*. *Ecology* 85, 1343–1354.
- Moran, M. J., 1985a. Distribution and dispersion of the predatory intertidal gastropod *Morula marginalha* Blainville. *Mar. Ecol. Prog. Ser.*, 22: 41-52.
- Moran, M. J., 1985b. The timing and significance of sheltering and foraging behaviour of the predatory gastropod *Morula marginalha* Blainville (Muricidae). *J. Exp. Mar. Biol. Ecol.*, 93: 103-114.
- Morgan, S.G. and Christy, J.H., 1994. Plasticity, constraint, and optimality in reproductive timing. *Ecology*, 75:2185–2203.
- Morgan, S.G. and Christy, J.H., 1995. Adaptive significance of the timing of larval release by crabs. *Am Nat*, 145:457–479.
- Moulton, J.S., 1962. Intertidal clustering of an intertidal gastropod. *Biol. Bull. Mar. Biol.*, 123: 170-178.
- Moura, G., Vilarinho, L. and Machado, J., 2000. The action of Cd, Cu, Cr, Zn, and Pb on fluid composition of *Anodonta cygnea*(L.): organic components. *Comparative biochemistry and physiology part b: biochemistry and molecular biology*, 127: 105–112.
- Munsiri, P., C. E. Boyd and B. F. Hajek, 1995. Physical and Chemical Characteristics of Bottom Soil Profiles in Ponds at Auburn, Alabama, USA and a Proposed System for Describing Pond Soil Horizons. *Journal of the World Aquaculture Society*, 26 (4):346 - 377.
- Murai, M. 1992. Courtship activity of wandering and burrow-holding male *Uca arcuata*. *Ethology* 92: 124-134.

- Murai, M. and P. R. Y., Backwell, 2005. More signalling for earlier mating: conspicuous male claw waving in the fiddler crab, *Uca perplexa*. *Animal Behaviour*, 2005, 70, 1093–1097.
- Murai, M., Goshima, S. and Kakasone, Y., 1982. Some behavioural characteristics related to food supply and soil texture of burrowing habitats of *Uca vocans* and *U. lactea perplexa*. *Mar Biol* 66: 191-197.
- Murai, M., Goshima, S. and Kakasone, Y., 1983. Adaptive droving behaviour observed in the fiddler crab *Uca vocans vocans*. *Mar Biol* 76: 159-164.
- Murai, M., S. Goshima, and Y. Henmi, 1987. Analysis of the mating system of the fiddler crab, *Uca lactea*. *Animal Behaviour*, 35: 1334-1342.
- Murray, J.M.H., Meadows, A., Meadows, P.S., 2002. Biogeomorphological implications of microscale interactions between sediment geotechnics and marine benthos: a review. *Geomorphology* 47, 15–30.
- Nakasone, Y. 1977. Crab zonation in the Yuhi River, Okinawa Island. *Jap. J. Ecol.*, 27: 61-70.
- Nakasone, Y., H. Akamine and K. Asato. 1983. Ecology of the fiddler crab *Uca vocans vocans* (Linnaeus) (Decapoda: Ocypodidae). Relation between the mating system and the drove. *Galaxea*, 2: 119-133.
- Naylor E., 1958. Tidal and diurnal rhythms of locomotor activity in *Carcinus maenas*. *J. exp. Biol.*, 35: 602-610.
- Needham, J.G. and Needham, P.R., 1962. A guide to fresh water biology. Holdendy Inc. San Francisco.
- Newell, R.C., 1979. Biology of intertidal animals. Marine Ecological Surveys. Elsevier, New York, New York, USA.
- Ng, P.L.K., D. Guinot and P.J. F. Davie, 2008. Systema brachyuraorum: Part I. An annotated checklist for extant brahcyurna crabs of the world. *The Raffles Bulletin of Zoology*, 17: 1-286.
- Nirmal Kumar, J. I., George, B., Kumar, R. N., Sajish, P. R., Viyol, S., 2009. Assessment of spatial and temporal fluctuations in water quality of a tropical permanent estuarine system, Tapi, west coast of India. *Appl. Ecol. Environ. Res.*, 7: 267–276.
- Nishida, A. K., N. Nordi and R. R. N. Alves, 2006. The lunar-tide cycle viewed by crustacean and mollusc gatherers in the State of Paraiba,

- Northeast Brazil and their influence in collection attitudes. *Journal of Ethnobiology and Ethnomedicine*, 2:1-12.
- Odum, E. P., 1971. *Fundamentals of ecology*. 3rd W.B. Saunders Company. Philadelphia, pp. 574.
- Omori, K., 1995. The adaptive significance of a lunar or semilunar reproductive cycle in marine animals. *Ecol Model*, 82:41–49.
- Ono, Y., 1965. On the ecological distribution of ocypodid crabs in the estuary. *Mem. Fac. Sci. Kyushu Univ., Ser. E (Bioi.)*, 4: 1-60.
- Otokunefor, T. V. and Obiukwu, C., 2005. Impact of Rifinery Effluent on physico -chemical Properties of a water body in Niger Delta. *Applied Ecology and Environ. Research*, 3(1):61-72.
- Paine, R.T., 1974. Intertidal community structure: experimental studies on the relationship between a dominant competitor and its principal predator. *Oecologia*, 15: 93–120.
- Pajevic, S., Vuckovic, M., Stankovic, Z., Krstic, B., Kevresan, Z., Radulovic, S. (2002). The content of some macronutrient and heavy metals in aquatic macrophytes of three ecosystems connected to the Danube in Yugoslavia. *Arch. for Hydrobio. Suppl.*, 141: 73-83.
- Pala, A., K. Y. Gina, A. Y. Linb and M. Reinhard, 2010. Impacts of emerging organic contaminants on freshwater resources: Review of recent occurrences, sources, fate and effects. *Science of the Total Environment*, 408(24): 6062–6069.
- Palmer, M. A. and Brandt, R.R., 1981. Tidal variation in sediment densities of marine benthic copepods. *Mar. Ecol. Prog. Ser.* 4, 207–212.
- Panda, U. C., Sundaray, S. K., Ratha, P., Nayak, B. B., Bhatta, D., 2006. Application of factor and cluster analysis for characterization of river and estuarine water systems – A case study: Mahanadi River (India). *Journal of Hydrology*, 331: 434-445.
- Pandya P. J. and K. D. Vachhrajani, 2013. On the diversity of brachyuran crabs from lower estuarine mud flats of Mahi River with new record of two species (Decapoda: Brachyura) from Gujarat, India. *Arthropods*, 2(4), 242-250.

- Pandya P. J., 2011. Benthic community structure of Mahi River estuary with special reference to animal-sediment relationship. Ph. D. Thesis. The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India.
- Pandya, P. J. and Vachhrajani, K. D., 2011. Life under ecological stress: An estuarine case study. In: Animal Diversity, Natural History and Conservation (Gupta VK, Verma AK, Eds). Daya Publication House, New Delhi, India, 1: 427- 436.
- Pandya, P. J. and Vachhrajani, K. D., 2012. Malacological study of Mahi river, Gujarat (India) with reference to estuarine gradient. Elect. J. Env. Sci., 2012, 5: 41-47.
- Pardo, M., Palma, A., Prieto, C., Sepúlveda, P., Valdivia, I., Ojeda, P., 2007. Processes regulating early post-settlement habitat use in a sub tidal assemblage of brachyuran decapods. J. Exp. Mar. Biol. Ecol., 344: 10–22.
- Parrish, J. K., 1999. Using behavior and ecology to exploit schooling fishes. Environ Biol Fishes, 55:157–181.
- Patel V. And P. Punita, 2013. Assessment of seasonal variation in water quality of River Mini, at Sindhrot, Gujarat. Inter. J. Envi. Sci., 3(5): 1424-1436.
- Patel, B. K. and K. D. Vachhrajani, 2013. Pollution Status in Mangrove Ecosystem of Mahi and Dadhar River Estuaries. National Conference on Biodiversity: Status and Challenges in Conservation, pp163-172.
- Paterson, D.M., 1997. Biological mediation of sediment erodibility: ecology and physical dynamics. In: Burt, T.N., Parker, W.R., Watts, J. (Eds.), Cohesive Sediments. Wiley, Chichester, pp. 215–229.
- Perillo, G.M.E., Minkoff, D.R. and Piccolo, M.C., 2005. Novel mechanism of stream formation in coastal wetlands by crab-fish-groundwater interaction. Geo-Marine Letters, 25: 214–220.
- Perkins, E.J., 1974. The biology of estuaries and coastal waters. New York Academic Press.
- Peter M.J.H., J.J. Middelburg and C.H.R. Heip, 2001. Benthic community structure and sediment processes on an intertidal flat: results from the ECOFLAT project. Continental Shelf Research, 21: 2055–2071.

- Peterson, C. H. and Andre, S.V., 1980. An experimental analysis of interspecific competition among marine filter feeders in a soft-sediment environment. *Ecology*, 61: 129– 139.
- Peterson, C.H., 1977. Competitive organization of the soft bottom macrobenthic communities of southern California lagoon. *Mar. Biol.*, 43: 343–359.
- Phillips, D. J. H. 1985. Organochlorines and trace metals in greenlipped mussels, *Perna viridis*, from Hong Kong waters: a test of indicator ability. *Mar. Ecol. Prog. Ser.*, 21: 251-258.
- Phillips, D. J. H. and K. Muttarasin, 1985. Trace metals in bivalve molluscs from Thailand. *Mar. Environ. Res.*, 15: 215-234.
- Phillips, D. J. H. and P. S. Rainbow, 1988. Barnacles and mussels as biomonitors of trace elements: a comparative study. *Mar. Ecol. Prog. Ser.*, 49: 83-93.
- Phillips, D. J. H. and P. S. Rainbow, 1993. *Biomonitoring of Aquatic Trace Contaminants*. Chapman and Hall, London.
- Phillips, D. J. H. and W. W. S. Yim, 1981. A comparative evaluation of oysters, mussels and sediments as indicators of trace metals in Hong Kong waters. *Mar. Ecol. Prog. Ser.*, 6: 285-293.
- Pope, D. S., 2000. Testing function of fiddler crab claw waving by manipulating social context. *Behavioral Ecology and Sociobiology*, 47: 432-437.
- Potter, I. C., Chuwen, B.M., Hoeksema, S.D. Elliott, M., 2010. The concept of an estuary: A definition that incorporates systems which can become closed to the ocean and hypersaline. *Est. Coast. Shelf Sci.*, 87: 497–500.
- Pritchard, D.W., 1967. "What is an estuary: physical viewpoint". In Lauf, G. H., *Estuaries*. A.A.A.S. Publ. 83. Washington, DC. pp. 3–5.
- Rainbow, P. S. and B. D. Smith, 1992. Biomonitoring of Hong Kong coastal trace metals by barnacles, 1986-1989. (Eds.) B. Morton, *In the Marine Flora and Fauna of Hong Kong and Southern China III. Proceedings of the Fourth International Marine Biological Workshop*, pp. 585-597.

- Rainbow, P. S., Z. G. Huang, Y. Songkai and B. D. Smith, 1993. Barnacles as biomonitors of trace metals in the coastal waters near Xiamen, China. *Asi. Mar. Biol.*, 10: 109-121.
- Rainbow, P. S., 1995. Biomonitoring of Heavy Metal Availability in the Marine Environment. *Marine Pollution Bulletin*, 31 (4-12): 183-192.
- Rajan, K. P., Kharouf, H. H. and Lockwood, A. P. M., 1979. Rhythmic cycles of blood sugar concentrations in the crab *Carcinus maenas*. In *Cyclic Phenomena in Marine Plants and Animals* (Edited by Naylor E. and Hartnoll R. G.). pp. 451-458.
- Ravaniah, G., Georgeena Kumari, P., morthy, C. V. M., 2010. Water quality analysis of the Pennar estuary, Nellore. *J. Curr. Sci.*, 15(2): 321-334.
- Reading, K. and Backwell, P.R.Y., 2007. Can beggars be choosers? Male mate choice in a fiddler crab. *Anim. Behav.*, 74: 867–872.
- Reaney, L.T., Backwell, P.R.Y., 2007. Risk-taking behavior predicts aggression and mating success in a fiddler crab. *Behav. Ecol.*, 18: 521–525.
- Reichert, S. E., 1984. Games spiders play III: cues underlying context-associated changes in agonistic behaviour. *Anim. Behav.*, 32: 1-15.
- Reid D. G. and Naylor E., 1989. Are there separate circatidal and circadian clocks in the shore crab *Carcinus maenas*. *Mar. Ecol. Prog. Ser.*, 52: 1-6.
- Reid D. G. and Naylor E., 1993. Different free-running periods in split components of the circatidal rhythm in the shore crab *Carcinus maenas*. *Mar. Ecol. Prog. Ser.*, 102: 295-302.
- Renshun, Z., 1992. Suspended sediment transport processes on tidal mud flat in Jiangsu Province, China. *Estuar. Coast. Shelf Sci.*, 35: 225–233.
- Rhoads, D. C. and Young, D. K., 1970. The influence of deposit feeding organisms on sediment stability and community trophic structure. *J. mar. Res.*, 28: 150-178.
- Rhode, K. and R. Sandland, 1975. Factors influencing clustering in the intertidal snail *Cerithium monoliferum*. *Mar. Biol.*, 30: 203-216.
- Robertson, D. R., Petersen, C. W., Brawn, J. D., 1990. Lunar reproductive cycles of benthic-brooding reef-fishes: reflections of larval biology or adult biology? *Ecol Monogr*, 60:311–329.

- Robertson, D. R., Swearer, S. E., Kaufmann, K. and Brothers, E. B., 1999. Settlement vs. environmental dynamics in a pelagic pawning reef fish at Caribbean Panama. *Ecol Monogr*, 69:195–218.
- Rombouts, G. Beaugrand, L.F. Artigas, J. C. Dauvin, F. Gevaert, E. Goberville, D. Kopp, S. Lefebvre, C. Luczak, N. Spilmont, M. Travers-Trolet, M.C. Villanueva and R.R. Kirby., 2013. Evaluating marine ecosystem health: Case studies of indicators using direct observations and modelling methods. *Ecological Indicators*, 24: 353-365.
- Rosberg, R.K., Levitan, D., 1992. For adults only? Supply-side ecology and history of larval biology. *Trends Ecol. Evol.*, 7: 130– 133.
- Rosenberg, M.S., 2001. The systematics and taxonomy of fiddler crabs: A phylogeny of the genus *Uca*. *J Crustac Biol*, 21:839-869.
- Rosenberg, R. H. and M. Enquist, 1991. Contest behaviour in Weidmeyer's admiral butterfly *Limenitis weidemeyerii* (Nymphalida): the effect of size and residency. *Anim. Behav.*, 42: 805 811.
- Rtal, A., L. Nonnotte and J. P. Truchot, 1996. Detoxification of exogenous copper by binding to hemolymph proteins in the shore crab, *Carcinus maenas*. *Aquat Toxic.*, 36, 235-252 (1996).
- Ruddle, K., 1994. Local knowledge in the folk management of fisheries and coastal marine environments. In *Folk Management in the World's Fisheries* Edited by: Dyer, C.L. and McGoodwin, J.R. University Press of Colorado, Niwot, Colorado.
- Russo, S., Gherardi, F., Vannini, M., 1998. Anti-predatory role of the droves composed by the tropical crab, *Dotilla fenestrata*. *Ins Soc Life*, 2: 191-194.
- Rybak, A., Messyasz, B. and Lecka, B., 2013. The accumulation of metal (Co, Cr, Cu, Mn and Zn) in freshwater *Ulva* (Chlorophyta) and its habitat. *Ecotoxicology*, 22: 558–573.
- Rygg, B., 1985. Effect of sediment copper on benthic fauna. *Marine Ecology Progress Series* 25, 83–89.
- Rzymiski, P., Niedzielski, P., Klimaszyk, P. and Poniedzialek, B., 2014. Bioaccumulation of selected metals in bivalves (Unionidae) and

- Phragmites australis* inhabiting a municipal water reservoir. Environ. Monit. Assess., 185(5): 3199–3212.
- Safahieh, A., Abdolapour, M. F. and Savari, A., 2011. Heavy metal contamination in sediment and Sole Fish (*Euryglossa orientalis*) from Musa Estuary (Persian Gulf). World J Fish Mar Sci, 3: 290-297.
- Saher, N. U. and N. A. Qureshi, 2010. Zonal distribution and population biology of *Ilyoplax frater* (Brachyura: Ocypodoidea: Dotillidae) in a coastal mudflat of Pakistan. Current Zoology, 56 (2): 244–251.
- Salahshur, S., Zeinab Y. and A. R. Bakhtiari, 2014. Bioaccumulation of Cd, Pb and Zn in the Oyster *Saccostrea cucullata* and Surface Sediments of Hendourabi Island-Persian Gulf, Iran. J Mar Biol Oceanogr, 3:2.
- Salmon, M, 1987. On the reproductive behavior of the fiddler crab *Uca Thayeri*, with comparisons to *U. pugilator* and *U. vocans*: evidence for behavioral convergence. J Crustac Biol, 7:25–44
- Santschi, P., P. Hohener, G. Benoit and M. B. Brink, 1990. Chemical processes at the sediment-water interface. Marine Chemistry, 30: 269-315.
- Satpathy, K. K., Mohanty, A. K., Natesan, U., Prasad, M. V. R., Sarkar, S. K., 2009. Seasonal variation in physicochemical properties of coastal waters of Kalpakkam, east coast of India with special emphasis on nutrients. Environ. Monit. Assess. 164: 153–171.
- Schubart, C. D. and R. Diesel, 1998. Osmoregulatory capacities and penetration into terrestrial habitats: a comparative study of Jamaican crabs of the genus *Armases Abele*, 1992 (Brachyura: Grapsidae: Sesarminae). Bull. mar. Sci., 63: 743–752.
- Scott D.A, 1989. A Directory of Asian Wetlands, IUCN, the World Conservation Union, Gland, Switzerland, and Cambridge, UK. pp.118.
- Scott, J. K. and M. S. Redmond, 1989. The effects of a contaminated dredged material on laboratory populations of the tubicolous amphipod *Ampelisca abdita*. In, Aquatic Toxicology and Hazard Assessment edited by U.M. Cowgill and L.R. Williams. ASTM, Philadelphia, pp. 289-301.
- Scott, J.A., 1974. Mate-location behaviour of butterflies. Am. Midl. Nat., 91: 103-117.

- Shih, H.T., Mok, H.K. and Chang, H.W., 2005. Chimney building by male *Uca formosensis* Rathbun 1921 (Crustacea: Decapoda: Ocypodidae) after pairing: a new hypothesis for chimney function. *Zool. Stud.*, 44: 242–251.
- Shukla, M. L., B. K. Patel, J. N. Trivedi and K. D. Vachhrajani, 2013. Brachyuran crabs diversity in mangrove habitat of Gulf of Khambhat, Gujarat, India, *Res. J.Mar. Sci.*, 1(2): 8-11.
- Sibly, R. M. and P. Calow, 1989. A life cycle theory of response to stress. *Biol. J. Linn. SOL.*, 37: 101-116.
- Singh, K. P., Malik A. and Sinha S., 2005. Water quality assessment and apportionment of pollution sources of Gomti river (India) using multivariate statistical techniques—a case study. *Analytica chimica Acta*, 538: 14 – 27.
- Singh, G., Ramanathan, A. L., Prasad, M. B. K., 2005. Nutrient cycling in Mangrove ecosystem: a brief overview. *J. Ecol. Environ. Sci.*, 30(3):231–244.
- Site, A. D., 2001. Factors Affecting Sorption of Organic Compounds in Natural Sorbento Water Systems and Sorption Coefficients for Selected Pollutants- A Review. *J. Phys. Chem. Ref. Data*, 30(1): 187-439.
- Skov, M. W. and R. G. Hartnoll, 2001. Comparative suitability of binocular observation, burrow counting and excavation for the quantification of the mangrove fiddler crab *Uca annulipes* (H. Milne Edwards). *Hydrobiologia* 449, 201–212.
- Slatyer, R.A., E.S.Y. Fok, R. Hocking and P.R.Y. Backwell, 2008. Why do fiddler crabs build chimneys? *Biol. Lett.*, 4: 616-618.
- Snelgrove, P.V.R. and Butman, C.A., 1994. Animal/sediment relationships revisited: cause versus effect. *Mar. Biol. Annu. Rev.*, 32: 111-177.
- Sousa, W.P., 1980. The response of a community to disturbance: the importance of successional age and species life histories. *Oecologia*, 45: 72–81.
- Spencer, K. L. and MacLeod, C. L., 2002. Distribution and partitioning of heavy metals in estuarine sediment cores and implications for the use of sediment quality standards. *Hydrol. Earth Syst. Sci.*, 6: 989-998.

- Spicer, J. I. and R. E. Weber, 1992. Respiratory impairment by water-borne copper and zinc in the edible crab *Cancer pagurus* (Crustacea: Decapoda) during hypoxic exposure. *Mar. Biol.*, 112: 429-435.
- Stankovic, Z., Pajevic, S., Vuckovic, M. and Stojanovic, S., 2000. Concentrations of trace metals in dominant aquatic plants of Lake Provala (Vojvodina, Yugoslavia). *Biolo. Plant.*, 43: 583-585.
- Stanton, M. L., 1983. Spatial patterns in the plant community and their effects upon insect search. In, *Herbivorous insects. Host-seeking behaviour and mechanisms*, edited by S. Ahmad, Academic Press, New York, pp. 125-151.
- Staples, C. A., Dorn, P. B., Klecka, G. M., O'Block, S. T. and Harris, L. R., 1998. A review of the environmental fate, effects and exposures of bisphenol A. *Chemosphere* 36: 2149–2173.
- Strbac, S., Sajnovic, A., Kasanin G. M., Vasic, N. Dojcinovic, B., Simonovic, P. and Jovancicevic, B., 2014. Metals in sediment and *Phragmites Australis* (common reed) from Tisza River, Serbia. *Applied Ecology and Environ. Research*, 12(1): 105-122.
- Sugiyama, Y., 1961. The social structure of a sand-crab, *Scopimera globosa* De Haan, with special reference to its population. *Physiol. Ecol. Japan*, 10: 10-17.
- Sunda, W. G. and R. R. Guillard, 1976. The relationship between cupric ion activity and the toxicity of copper to phytoplankton. *J. Mar. Res.* 34: 411-422.
- Sunda, W. G., D. W. Engel and R.M. Thuotte, 1978. Effect of chemical speciation on toxicity of cadmium to grass shrimp, *Palaemonetes pugio*: Importance of free cadmium ion. *Environ. Sci. Technol.*, 12: 409-413.
- Takagi K. K., P. Cherdskajai, I. Mimura, Y. Yano, K. Adulyanukosol and M. Tsuchiya, 2010. Soldier crab (*Dotilla myctiroides*) distribution, food resources and subsequent role in organic matter fate in Ao Tang Khen, Phuket, Thailand. *Est. Coast. Shelf Sci.*, 87: 611-617.
- Tall, L., Methot, G., Armellin, A. and Pinel-Alloul, B., 2008. Bioassessment of Benthic Macro invertebrates in Wetland Habitats of Lake Saint-Pierre (St. Lawrence River). *Journal of Great Lakes Research*, 34: 599-614.

- Talley, T.S., Crooks, J.A., Levin, L.A., 2001. Habitat utilization and alteration by the invasive burrowing isopod, *Sphaeroma quoyanum*, in California saltmarshes. *Mar. Biol.*, 138; 561–573.
- Taylor, M.H., 1984. Lunar synchronization of fish reproduction. *Trans Am Fish Soc*, 113:484–493.
- Thornhill, R., 1984. Fighting and assessment in *Harpobittacus scorpionflies*. *Evolution*, 38: 204-214.
- Thurman, C. L. 1984. Ecological notes on fiddler crabs of south Texas, with special reference to *Uca subcylindrica*. *J. Crust. Biol.*, 4: 665–681.
- Thurman, H.V., 1997. *Introductory Oceanography*. Prentice Hall, New Jersey:
- Todd, C.D. and Laverack, M.S. 1991. *Coastal marine zooplankton*. Cambridge University Press.
- Tripathi, V. M., Tewari, D. D., Tewari, H. D., Tewari, S. and Uppadhyay, M. P., 2008. Physico-chemical characteristics of river Rapti nearby industrial area Balrampur of UP, India. *Natl. Environ. pollu. Technol.*, 7(1): 73-77.
- Trivedi, D. J., Trivedi, J. N., Soni, G. M., Purohit, B. D. and Vachhrajani, K. D., 2015. Crustacean fauna of Gujarat state of India: a review. *Elect. J. Env. Sci*, 8: 23-31.
- Trivedi, J. N. and K. D. Vachhrajani, 2013. First record of *Cryptopodia angulata* H. Milne Edwards and Lucas, 1841 from Saurashtra coast, Gujarat, India (Decapoda: Brachyura: Parthenopidae). *Check List*, 9 (4): 897-898.
- Trivedi, J. N. and K. D. Vachhrajani, 2013a. First record of two porcellanid crabs from Gujarat state, India (Crustacea: Decapoda: Porcellanidae). *J. Mar. Biol. Ass. India*, 55 (1): 55-58.
- Trivedi, J. N. and K. D. Vachhrajani, 2013b. Taxonomic account of genus *Scylla* (de Haan, 1833) from Gujarat State, India with two new records of species. *Arthropods*, 2 (4): 159-171.
- Trivedi, J. N. and K. D. Vachhrajani, 2015. First record of brachyuran crab *Leptodius affinis* (De Haan, 1835) (Crustacea: Decapoda: Xanthidae) from the western coast of India. *Marine Biodiversity Records*, 8: e2 doi:10.1017/S175526721400133X.

- Trivedi, J. N. and Vachhrajani, K. D. 2013. First record of *Cryptopodia angulata* H. Milne Edwards and Lucas, 1841 from Saurashtra coast, Gujarat, India (Decapoda: Brachyura: Parthenopidae). Checklist, 9(4): 897–898.
- Trivedi, J. N., G. M. Soni and K. D. Vachhrajani, 2014. First record of brachyuran crab *Heteropanope glabra* Stimpson, 1858 (Crustacea, Decapoda, Pilumnidae) from India. Marine Biodiversity records, 8: e112 doi:10.1017/S1755267215000883.
- Trivedi, J. N., G. M. Soni, D. J. Trivedi and K. D. Vachhrajani, 2015. A new species of *Ilyoplax* (Decapoda, Brachyura, Dotillidae) from Gujarat, India. Journal of Asia-Pacific Biodiversity, 8: 173-177.
- Trivedi, J. N., G. M. Soni, S. Arya and K. D. Vachhrajani, 2014. First record of *Macrophthalmus laevis* A. Milne Edwards, 1867 (Decapoda: Brachyura: Macrophthalmidae) from India. J. Mar. Biol. Ass. India, 56 (2): 84-87.
- Trivedi, J. N., Gadhavi, M. K., Vachhrajani, K. D., 2012. Diversity and habitat preference of brachyuran crabs in Gulf of Kutch, Gujarat, India. Arthropods, 1(1): 13-23.
- Trivedi, J.N., Vachhrajani, K.D., 2012. Distribution and diversity of brachyuran crabs along the coastal region of Junagadh district, Gujarat. Proceedings of the Biodiversity and Conservation of Coastal and Marine Ecosystems of India, 8-14.
- Turchin, P. and P. Kareiva, 1989. Aggregation in *Aphis varians*: an effective strategy for reducing predation risk. Ecology, 70: 1008-1016.
- Turchin, P. and P. Kareiva, 1989. Aggregation in *Aphis varians*: an effective strategy for reducing predation risk. Ecology, 70: 1008-1016.
- Tweedie, M. W. F., 1950. Notes on grapsid crabs from the Raffles Museum. Bull Raffles Mus, 23: 310-324.
- Uncles, R. J. and J. A. Stephens, 1996. Salt intrusion in the Tweed Estuary. Estuar. Coast. Shelf Sci., 43: 271–293.
- Underwood, A.J., Denley, E. J. and Moran, M.J., 1983. Experimental analysis of the structure and dynamics of midshore rocky intertidal communities in New South Wales. Oecologia, 56: 202– 219.

- Uzoekwe, S. A. and F. A. Oghosanine, 2011. The effect of refinery and petrochemical effluent on water quality of Ubeji creek Warri, Southern Nigeria. *Ethiopian Journal of Environmental Studies and Management*, 4(2): 2011.
- Vachhrajani K. D. and Pandya P.J ., 2011. Life under ecological stress: An estuarine case study. In: *Animal Diversity, Natural History and Conservation* (Gupta, V.K., Verma,A.K.). Daya Publication House, New Delhi, India, 1: 427- 436.
- Van Damme, S., E. Struyf, T. Maris, T. Ysebaert, F. Dehairs, M. Tackx, C. Heip and P. Meire, 2005. Spatial and temporal patterns of water quality along the estuarine salinity gradient of the Scheldt estuary (Belgium and The Netherlands): results of an integrated monitoring approach. *Hydrobiologia*, 540: 29–45
- Van Holde, K. E., Miller, K. I. and Decker, H., 2001. Hemocyanin and invertebrate evolution. *Journal of Biological Chemistry*, 276: 15563–15566.
- Varadharajan, D., Soundarapandian, P. and Pushparajan, N., 2013. Effect of Physico-Chemical Parameters on Crabs Biodiversity. *J Marine Sci Res Dev*, 3:116.
- Verrell, P. A., 1986. Wrestling in the red-spotted newts (*Notophthalmus viridescens*): resource value and contestant asymmetry determine contest duration and outcome. *Anim. Behav.*, 34: 398-402.
- Vogel, F., 1984. Comparative and functional morphology of the spoon-tipped setae on the second maxillipeds in *Dotilla Stimpson*, 1858 (Decapoda, Brachyura, Ocypodidae). *Crustaceana* 47: 225–234.
- Waage, J. K., 1974. Reproductive behaviour and its relation to territoriality in *Calopteryx maculata* (Beauvois) (Odonata: Calopterygidae). *Behaviour*, 51: 240-256.
- Wada K, 1981. Growth, breeding and recruitment in *Scopimera globosa* and *Ilyoplax pusillus* (Crustacea: Ocypodidae) in the estuary of Waka River. *Middle Japan Publ. Seto. Mar Biol Lab*, 20: 243-259.

- Wada K, 1984. Pair formation in the two forms of *Macrophthalmus japonicus* De haan (Crustacean: Brachyura) at a co-occurring area. J. Ethol, 2: 7-10.
- Wada K, 1993. Territorial behaviour and size of home range and territory in relation to sex and body size in *Ilyoplax pusilla* (Crustacea: Brachyura; Ocypodidae). Mar Biol., 115; 47-52.
- Wada, K. and Murata, I. 2000. Chimney building in the fiddler crab *Uca arcuata*. J. Crust. Biol., 20: 505–509.
- Wada, K., 1978. Two forms of *Macrophthalmus japonicus* De Haan (Crustacea: Brachyura). Pub Seto Mar Biol Lab, 24: 327-340.
- Wada, K., 1983. Spatial distributions and population structures in *Scopimera globosa* and *Ilyoplax pusillus* (Decapoda: Ocypodidae). Publ. Seto Mar. Biol. Lab., 27: 281–291.
- Wada, K., 1984. Barricade building in *Ilyoplax pusillus* (De Haan) (Crustacea: Brachyura). Journ. exp. mar. Biol. Ecol., 83: 73-83.
- Wada, K., S. Yum, and J.K. Park, 1994. Mound building in *Ilyoplax pingi* (Crustacea: Brachyura: Ocypodidae). Mar. Biol. 121: 61-65.
- Walkly, A. and I. A. Black., 1934. An Examination of Degtjareff Method for Determining Soil Organic Matter and a Proposed Modification of the Chromic Acid Titration Method. Soil Sci., 37: 29-37.
- Wang, C., G. Han, Y. Jia, X. Feng, P. Guo and X. Tian, 2011. Response of litter decomposition and related soil enzyme activities to different forms of nitrogen fertilization in a subtropical forest. Ecol. Res., 26 (3): 505-513.
- Watling, L., 1991. The sedimentary milieu and its consequences for resident organisms. American Zoologist, 31:789–796.
- Weis, J. S. 1980. Effect of zinc on regeneration in the fiddler crab *Uca pugilator* and its interactions with methylmercury and cadmium. Mar. Env. Res., 3:249-255.
- Weisberg, S. B., Ranasinghe, J. A., Dauer, D. M., Schaffner, L. C., Diaz, R. J., Frithsen, J. B., 1997. An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. Estuaries, 20(1): 149–158.

- Widdows, J. and Brinsley, M., 2002. Impact of biotic and abiotic processes on sediment dynamics and the consequences to the structure and functioning of the intertidal zone. *J. Sea Res.*, 48(2): 143-156.
- Widdows, J., Brinsley, M.D., Bowley, N., Barrett, C., 1998. A benthic annular flume for in situ measurement of suspension feeding/ biodeposition rates and erosion potential of intertidal cohesive sediments *Est. Coast. Shelf Sci.*, 46: 27–38.
- Williams J. A., 1985. An endogenous tidal cycle of blood sugar concentrations in the shore crab, *Careinus maenas*(L.). *Comp. Biochem. Physiol.*, 81: 627-631.
- Wilson, W.H., 1980. A laboratory investigation of the effect of a terebellid polychaete on the survivorship of Nereid polychaete larvae. *J. Exp. Mar. Biol. Ecol.* 46, 73– 80.
- Wilson, W.H., 1991. Competition and predation in marine soft-sediment communities. *Ann. Rev. Ecol. Syst.*, 21: 221–241.
- Wolanski, E., 2007. *Estuarine Ecohydrology*. Elsevier, Amsterdam, 157 pp.
- Wood, R., Widdows, J., 2002. A model of sediment transport over an intertidal transect, comparing the influences of biological and physical factors. *Limnology and Oceanography*, 47: 848–855.
- Woodin, A., 1974. Polychaete abundance patterns in a marine soft-sediment environment: the importance of biological interactions. *Ecol. Monogr.*, 44: 171– 187.
- Woodin, A., 1976. Adult– larval interactions in dense infaunal assemblages: patterns of abundance. *J. Mar. Res.*, 34: 25– 41.
- Yamaguchi, T. and S. Tabata, 2005. Territory usage and defense of the fiddler crab, *Uca lactea* (Decapoda, Brachyura, Ocypodidae). *Crustaceana*, 77: 1055-1080.
- Yamaguchi, T. and Tanaka, M., 1974. Studies on the ecology of a sand bubbler crab, *Scopimera globosa* De Haan (Decapoda, Ocypodidae). I. Seasonal variation of population structure. *Jap. J. Eco.*, 24: 165-174.
- Yamaguchi, T., 1971. Courtship behavior of a fiddler crab, *Uca lactea*. *Kumamoto Journal of Science* 10: 13-37.

- Yamaguchi, T., 1998. Evidence of actual copulation in the burrow in the fiddler crab, *Uca lactea* (De Haan, 1835) (Decapoda, Brachyura, Ocypodidae). *Crustaceana*, 71: 565-570.
- Yamaguchi, T., 2001a. The breeding period of the fiddler crab, *Uca lactea* (Decapoda, Brachyura, Ocypodidae) in Japan. *Crustaceana*, 74:285-293.
- Yamaguchi, T., 2001b. The mating system of the fiddler crab, *Uca lactea* (Decapoda, Brachyura, Ocypodidae) in Japan. *Crustaceana*, 74:389-399.
- Yamaguchi, T., 2001c. Daytime larval release of the fiddler crab, *Uca lactea* (Decapoda, Brachyura, Ocypodidae). *Crustaceana*, 74:545-555.
- Yang, H. and N. L. Rose, 2003. Distribution of Hg in the lake sediments across the UK. *Science of the Total Environment*, 304: 391-404.
- Yin, K., P. J. Harrison, S. Pond and R. J. Beamish, 1995a. Entrainment of nitrate in the Fraser river estuary and its biological implications. II. Effects of spring vs. neap tide and river discharge. *Estuar.Coast. Shelf Sci.*, 40: 529-544.
- Yin, K., P. J. Harrison, S. Pond and R. J. Beamish, 1995b. Entrainment of nitrate in the Fraser river estuary and its biological implications. III. Effects of winds. *Estuar.Coast. Shelf Sci.*, 40, 545-558.
- Ysebaert, T. and Herman, P.M.J., 2002. Spatial and temporal variation in benthic macrofauna and relationship with environmental variables in an estuarine, intertidal soft-sediment environment. *Mar. Ecol. Prog. Ser.*, 244: 105-124.
- Ysebaert, T., Herman, P.M.J., Meire, P., Craeymeersch, J., Verbeek, H. and Heip C.H.R., 2003. Large-scale spatial pattern in estuaries: estuarine macrobenthic communities in the Schelde estuary, NW Europe. *Est. Coast. Shelf Sci.* 57: 335-355.
- Zajac, R. N. and R. B. Whitlatch, 1989. Natural and disturbance induced demographic variation in an infaunal polychaete, *Nephtys incisu*, *Mar. Ecol. Prog. Ser.*, 57: 89-102.

- Zhang, H., B. Cui, R. Xiao and H. Zhao, 2010. Heavy metals in water, soils and plants in riparian wetlands in the Pearl River Estuary, South China. *Procedia Environmental Sciences*, 2: 1344–1354.
- Zimmer-Faust, R. K., 1987. Substrate selection and use by a deposit-feeding crab. *Ecology*, 68: 955–970.
- Zucker N, 1988. Preliminary observations of chelipeds use during social activities in sentinel crabs (Brachyura, family: Ocypodidae, genus: Macrophthalmus) from northern Queensland, Australia. *Bull Mar Sci.*, 43: 98-102.
- Zucker, N., 1974. Shelter building as a means of reducing territory size in the fiddler crab, *Uca terpsichores* (Crustacea: Ocypodidae). *The American Midland Naturalist* 91: 224-236.
- Zucker, N., 1978. Monthly reproductive cycles in three sympatric hood building tropical fiddler crabs (genus *Uca*). *Biol Bull*, 155:410–424.
- Zucker, N., 1981. The role of hood-building in defining territories and limiting combat in fiddler crabs. *Animal Behaviour*, 29: 387-395.