

FIGURES

	Page
1. Effect of angiotensin (10 μ g) on heart rate and blood pressure under the influence of morphine and chloralose (same anaesthetic agent used throughout).	29
2. Effects of equipressor doses of angiotensin (2.5 μ g) and Pitressin (2 units) on heart rate of the same dog (above, angiotensin used; below, Pitressin).	31
3. Effects on heart rate of angiotensin (10 μ g) and Pitressin (3 units) as influenced by mechanical buffering of the rise in blood pressure.	40
4. Effects of angiotensin (10 μ g) on heart rate before and during buffering of pressor response.	41
5. Changes in heart rate and blood pressure during slow infusion of angiotensin (1.25 μ g/min.).	43
6. Effects on heart rate and blood pressure produced by progressively increasing the rate of injection of angiotensin. I - Angiotensin 0.63 μ g. per minute, II - Angiotensin 1.25 μ g. per minute, III- Angiotensin 2.50 μ g. per minute; IV - Angiotensin 5.00 μ g. per minute.	45
7. Effects of angiotensin (10 μ g) on heart rate and blood pressure before and after sinoaortic denervation.	49
8. Effect of vagal stimulation and angiotensin (10 μ g) on heart rate and blood pressure after the ganglia blocking by pentolinium (kymograph tracing).	56

- | | | |
|-----|--|----|
| 9. | Effect of angiotensin (10 μ g) on heart rate before and after tying the adrenal veins. | 59 |
| 10. | Effect of angiotensin (10 μ g) on heart rate after inderal (5 mg/kg body weight). | 61 |
| 11. | Effect of angiotensin (10 μ g) on heart rate after vagotomy and after vagotomy plus inderal (5 mg/kg body weight). | 63 |
| 12. | Effect of angiotensin (10 μ g) on heart rate after reserpine (2-3 mg/kg body weight). | 66 |
| 13. | A comparative study of effect of angiotensin (10 μ g) on heart rate after-
a) tying adrenal vein, b) injection of inderal, and c) injection of reserpine. | 68 |
-