
References

- 1. Andrew at el (1980) Tuberculosis in patients with end stage of renal disease. Am. J. Med. 1980; 68: 59-65.
- Aneja KS at el. (1980) Feasibility of involvement of multipurpose health worker in District Tuberculosis Programme. Ind J Tub, 1980; 27: 158.
- 3. Aneja KS at el. (1980) Influence of initial motivation on treatment of tuberculosis patients. Ind J Tub, 1980; 27: 123.
- 4. Aneja KS at el. (1984) Active case finding in tuberculosis as a component of primary health care. Ind J Tub, 1984; 31: 65.
- 5. Aneja KS, Krishnamoorthy VV. (1982) Influence of training variation of case finding at peripheral health institutions. In District Tuberculosis Programme. NTI Newsletter, 1982; 19: 22.
- 6. Arora BC (1976) Case finding and treatment in National Tuberculosis Programme. Ind J Pub Health, 1976; 13: 35.
- 7. Arora BC at el. (1980) Problems of organization and administration of District Tuberculosis Programme in Southern Region. NTI Newsletter, 1980; 17: 42.
- 8. Arya VK. (2000) An evaluative study of National Tuberculosis Programme in Anand district, Gujarat. A study report on supervised field training for DPH. All India institute of hygiene and public health Calcutta. April 2000.
- Baily GVJ at el. (1967) Potential yield of pulmonary tuberculosis cases by direct microscopy of sputum in a district of south India. Bull WHO, 1967; 37: 875.

- 10. Baily GVJ at el. (1974) A concurrent comparison of an unsupervised twice-weekly regimen of chemotherapy in a routine outpatient treatment programme. Ind J Tub, 1974; 21: 152.
- 11. Baily GVJ at el. (1983) Tuberculosis control in India current problems and possible solutions. Ind J Tub, 1983; 30(2): 45.
- Banerji D. (1967) Behavior of tuberculosis patients towards a treatment organization offering limited supervision. Ind J Tub, 1967; 14(3): 156.
- Banerji D. (1969) Effect of treatment default on results of treatment in routine practice in India. Proceeding of the 20th International TB Conference, Paris :IUAT, 1969.
- Banerji D. (1983) The approach of primary health care and India's National Tuberculosis Programme. Proceeding of annual Conference of the Madhya-Pradesh TB Association, Rewa (MP) 1983.
- 15. Banerji D. (1989) Rural social transformation and changes in health behavior. Eco & pol weekly, 1989; 1474.
- 16. Banerji D. (1993) A social science approach to strengthening India's National Tuberculosis programme. Ind J Tub, 1993; 40: 61.
- Banerji D. Anderson S. (1963) A sociological study of awareness of symptoms among persons with pulmonary tuberculosis. Bull World Health Organ 1963; 29:665-683.
- Barnhoorn F, Adriaanse H (1992) In search of factors responsible for non-compliance among Tuberculosis patients in Wardha district in India. Social Sci. and Med. 1992; 34(3): 291-306.

- Bergner L, Yerby AS (1968) Low income and barrier to use of health services. N. Engl. J. Med. 1968; 287: 541-546.
- 20. Bhatnagar D, Almeida J. (1997) Factor influencing the perceived priority of tuberculosis in India.1997.WHO/97/234.
- 21. Boucot KR at el (1952) Tuberculosis among diabetics. The Philadelphia survey. Am. Rev. Tuberc. 1952; 65: 1-50.
- CDC (1997) Center for disease control and prevention. Tuberculosis morbidity- United States 1996. Morb. Mortal. Wkly. Repo. 1997; 46: 695-700.
- 23. Chakraborty A.K. et al (1995) Prevalence of tuberculosis in rural area by an alternative surveys method with out prior radiographic screening. Tuber. And Lungs Disease. 1995; 76: 1.
- Chakraborty AK (1997) Prevalence and incidence of tuberculosis infection and disease in India: A comprehensive review. 1997. WHO/TB/97.231.
- 25. Chakraborty AK. (1981) Tuberculosis case finding among symptomatic in the community A reappraisal. Ind J Tub, 1981;
 28: 12.
- 26. Chandrashekhar P. (1988) Eradication of tuberculosis a myths or reality? Ind J Tub, 1988; 35: 61.
- 27 Chandrashekhar P. (1985) Concept of surveillance and monitoring in tuberculosis. NTI Newsletter, 1985; 21(3): 54.
- 28 Chaulet P. (1986) Treatment of Tuberculosis: Case holding until cure. WHO, 1986; WHO/TB/83.141 Rev.1.

- 29. Chaulk CP et al. (1995) Eleven years of community- based directly observed therapy for tuberculosis. JAMA 1995; 274: 945-951
- 30. China Tuberculosis Control Collaboration. (1996) Results of directly observed short-course chemotherapy by 112,842 Chinese patients with smear-positive tuberculosis. Lancet 1996; 347:358-362.
- Comstock G. W. at el (1974) The prognosis of a positive tuberculin reaction in child hood and adolescence. Am. J. Epidemiol. 1974; 99: 131-138.
- 32. Cowie RL (1994) The epidemiology of tuberculosis in gold minors with silicosis. Am. J. Respir. Crit. Care Med. 1994; 156: 1460-1462.
- Dai ZC, Duanmu HJ. (1997) Directly Observed Treatment Shortcourse (DOTS) in china [abstract] 19th Eastern Region Conference of IUAT&LD, Singapore, 5-8 September 1997.
- Datta M. at el (2000) Tuberculosis in North arcot district of Tamil Nadu- a sample surveys. Ind. J. Tub. 2000; 147: 147-154.
- 35. Datta M. at el. (1993) Critical assessment of smears positive pulmonary tuberculosis patients after chemotherapy under the District Tuberculosis Programme. Tubercle, 1993; 74: 180.
- 36. Dholakia R. (1996) The potential economic benefits of DOTS strategy against TB in India.1996.WHO/TB/96.218.
- 37. DTC Anand (2000) NTP Action Plan 2000: 2.
- 38. DTC Mehsana (2000) RNTCP Action Plan year 2000; 5.

- 39. Dye C. at el (1999) Global burden of Tuberculosis. JAMA 1999; 282(7): 677-686.
- Edwards JH (1957) Contribution of smoking to respiratory disease.
 Br. J. Pre. Soc. Med. 1957; 11: 10-12.
- 41. Enavson DA at el (1957) The incidence of active tuberculosis in a large urban area. Am. J. Epidemi. 1989; 129: 1268-1276.
- 42. Fox W. (1961) Self- administration of medicaments. A review of published work and a study of the problems. Bull. Int Union tuberc 1961; 31:307-331.
- 43. Fox W. (1983) Compliance of patients and physicians: Experience and lessons from tuberculosis – I. Br Med J, 1983; 287: 1.
- 44. Fox W. (1983) Compliance of patients and physicians: Experience and lessons from tuberculosis – II. Br Med J, 1983; 287: 101.
- 45. Fox W. (1984) Short course chemotherapy for pulmonary tuberculosis and some problems of its programme application with particular reference to India. Lung India, 1984; 2(2): 161.
- 46. Fox W. (1988) tuberculosis case finding and treatment programme in the developing countries. Br Med Bull, 1988; 44: 44(4).
- 47. Fox W. (1990) Tuberculosis in India: past, present and future. Ind J Tub, 1990; 37: 175.
- 48. Friedman LN at el (1987) Tuberculosis screening in alcoholics and drugs addicts. Am. Res. Respi. Dis. 1987; 136: 1188-1292.
- 49. Frimodt Moller J. (1960) A community-wide tuberculosis study in a south Indian rural population. Bull WHO, 1960: 22: 61.

Reference[.]

- 50. Gehani at el. (1984) A study to determine the reliability of assessing the regularity of self-administration of drugs at home by patient's attendance at the clinic. Ind J Tub, 1984; 31: 74.
- 51. Gonzalez E, Armas L, Alonso A (1994) Tuberculosis in the Republic of Cuba: its possible elimination. Tuberc Lung Dis 1994:75:188-194.
- , 52. Gopi P.G. at el (1979) A tuberculosis prevalence surveys based on symptoms questioning and sputum examination. Ind. J. Tub. 1997; 44: 171.
- 53. Gothi G.D. at el (1976) Estimation of Prevalence of Bacillary -tuberculosis on the basis of X-ray and chest symptomatic screening. Ind. J. Med. Res. 1976; 64: 1150.
- 54. Gothi G.D. at el. (1979) Prevalence of Tuberculosis in south Indian district twelve year after initial surveys. Ind. J. Tub. 1979; 26: 121
- 55. Gothi GD at el. (1970) Case of pulmonary tuberculosis among the outpatients attending general health institutions in an Indian city. Bull WHO, 1970; 43: 35.
- 56. Gothi GD at el. (1971) Collection and consumption of selfadministered anti-tuberculosis drugs under programme conditions. Ind J Tub, 1971; 18(4): 107.
- 57. Gothi GD at el. (1978) Prevalence and incidence of sputum negative active pulmonary tuberculosis and fate of pulmonary radiological abnormalities found in a rural population. Inf J Tub, 1978; 25: 122.
- 58. Gothi GD, Baily GVJ. (1965) Problems of treatment of TB patients in rural area. Ind J Tub, 1965; 12: 22.

59. Govind P at el. (1976) An appraisal of different procedures of home visiting for reducing drugs default – an interim report. Ind J Tub, 1976; 23: 107.

,**`**

- 60. Govt. of India (1973) Report of committee of multipurpose worker under health and family planning programme. Ministry of Health and Family welfare, Govt. of India. 1973.
- 61. Govt. of India (1986) Job responsibilities of staff of the primary health centre. Rural health division, Ministry of Health and Family welfare, Govt. of India New Delhi 1986.
- 62. Govt. of India (1995) Background documents-workshop on revised National Tuberculosis Control Programme of India (18th-19th Feb. 1995. TB division DGHS New Delhi 1995; 6-7.
- 63. Govt. of India (2000) National Health Programme Annual Report 1999-2000. Ministry of Health and Family welfare, Govt. of India 2000; 174-177.
- 64. Goyal S.S., Mathur G.P., Pamra S.P. (1978) Tuberculosis trends in urban community. . Ind. J. Tub. 1978; 25: 77.
- 65. Groth P.E. at el (1959) Epidemiological basis of tuberculosis eradication in an advanced country. Bull. World. Health Organ. 1959; 21: 5-49.
- 66. Groth PE at el (1957) Epidemiological basis of tuberculosis eradication in an advanced country. Bull. World Health Organ. 1957; 21: 5-49.
- 67. Gupta SP. (1985) National Tuberculosis Programme: Its development concepts, monitoring and evaluation aspects. J Comm Dis, June 1985; 17(2): 146.

- 68. Holm J. (1994) TB control in developing world: it is time for a: change. World Health Forum, 1994; 5: 103.
- 69. Holmes C B, Hausler H, Nunn P (1998) A review of sex difference in the epidemiology of tuberculosis. Int J. Tuber. Lung Dis. 1998; 2: 96-104
- 70. ICMR (1959) Tuberculosis in India- A Sample Survey 1955-1958. Special report series No.34
- 71. ICORCI. (1988) In depth study of National Tuberculosis Programme – An overview. Institute of Communication, Operations Research and Community Development, Bangalore, India. November 1988.
- 72. Jagota P at el. (1989) Study of camps for examining sputum of chest symptomatic attending outpatients of peripheral health institutions. Ind J Tub, 1989; 36(1): 27.
- 73. Jagota P at el. (1994) Fate of smear positive patients of pulmonary tuberculosis at an urban District Tuberculosis Centre, five years after treatment. Ind J Tub, 1994; 41: 223.
- 74. Jagota P, Aneja KS. (1978) Follow-up of adequately treated cases of pulmonary tuberculosis a futile effort. NTI Newsletter, 1978; 15: 45.
- 75. Jayaswal R (1998) Tuberculosis and AIDS. 49th TB Seal Campaign 1998 The TB association of India 1998.
- 76. Khanna BK at el. (1977) Drugs default in tuberculosis. Ind J Tub, 1977; 24(3) 31.

77. Khatri GR. (1999) The revised national Tuberculosis Control Programme: A status reports on first 1,00,000 patients. Ind J tub 1999; 64: 157-166.

Υ.

- 78. Kilpatrick. (1987) Compliance in relation to tuberculosis Supplement to Tubercle, 1987; 68: 31.
- 79. Kochi A. (1991) The global tuberculosis situation and new control strategy of the World Health Organization. Tubercle, 1991; 72: 1.
- Krishna swami KV at el. (1977) A sociological study of awareness of symptoms and action taken by the patients to seek relief. Ind J Tub, 1977; 24(1):-4-5.
- 81. Krishna swami KV at el. (1979) An investigation of accuracy of home address given by patients in an urban community in south India. Tubercle, 1979; 60(1).
- 82. Krishnaswamy KV, Rahmin MA, Parthsarthy R (1977) A sociological study of awareness of symptoms of pulmonary tuberculosis and action taken by the patients to seek relief. Ind. J. Tub. 1977; 24: 15-20.
- 83. Kumaresan JA, Ali AK, Parkkli LM. (1998) Tuberculosis control in Bangladesh: success of the DOTS strategy. Int J Tuberc Lung Dis 1998; 2: 992-998.
- Lalit Kant (2000) An estimation of burden of TB in India (Editorial) Ind. J. Tub. 2000; 47(3): 127-128.
- 85. Leroy S et al (1997) Progression of HIV infection in patients with tuberculosis disease. Am. J. Epidem. 1997; 145: 293- 300.
- Lowe CR (1956) An association between smoking and respiratory tuberculosis. Br. J. Prev Med. 1956; 2: 1081-1086

- Luelmo F. (1994) India Revised National Tuberculosis Programme. May 1994; WHO/TUB.
- Mayurnath at el. (1991) Prevalence study of tuberculosis infection over 15 years in a rural population in Chingaleput District (South India). Ind J Med Res, 1993: 93.
- Melotra ML at el. (1971) Defaults and their effects on pulmonary tuberculosis patients on intermittent chemotherapy. Ind J Tub, 1971; 18: 114.
- 90. Mohanty KC at el (1998) Changing trend of HIV infection in patients with respiratory disease in Bombay since 1988. Ind. J. Tub. 1998; 41: 147-150.
- 91. Mohanty KC, Nair S, Sharabudhe T. (1994) Changing trend of HIV infection in patients with respiratory disease in Bombay since 1998. Ind J Tub 1994; 41: 147-150.
- 92. Moller F. at el (1981) A control study of the effect of domiciliary tuberculosis chemotherapy programme in a rural community in south India. Ind. J. Med. Res. 1981; 73: 1.
- 93. Murray CJL at el. (1991) Cost effectiveness of chemotherapy for pulmonary tuberculosis in three sub-Saharan African countries. Lancet 1991; 338: 1305-1308
- 94. Murthy N (1982) Reluctant patients-The women of India. World Health Forum 1982; 3(3): 315-316.
- 95. Nagpal DR at el. (1977) Prevalence of symptoms in a rural community of South India and the utilization of area health center. Ind J Med Res, 1977; 66: 635.

- 96. Nagpal DR. (1967) District Tuberculosis Programme in concept and outline. Ind J Tub, 1967; 14: 186.
- 97. Nagpal DR. (1987) Sociological aspect of tuberculosis for programme assessment. Int J Tub, 1987; 34: 101.
- Nagpal DR. (1989) India's National Tuberculosis Programme An overview. Ind J Tub, 1989; 36: 205.
- 99. Nagpal GR. (1982) Why integrated tuberculosis programme have not succeeded as per expectations in many developing countries: A collection of observations. Ind J Tub, 1982; 29(3): 149.
- 100. Nair SS et al (1998) Behavior pattern of chest symptomatic (unpublished) 1998.
- 101. Nair SS, Gothi GD. (1973) place of contact examination in tuberculosis programme. Ind J Tub, 1973; 20(4): 164.
- 102. Nair SS. (1974) Significance of patients with X-ray evidence of active tuberculosis not bacteriologically confirmed. Ind J Tub, 1974; 21: 3.
- 103. Nair SS. (1977) A simple model for planning and assessment of programme for tuberculosis control. Ind J Pub Health, 1977; 21: 111.
- 104. Nair SS. (1977) Determination of the appropriate index and time for assessing the effectiveness of a Tuberculosis Control Programme. Ind J Tub, 1977; 24(2): 58.
- 105. Narang P. at el (1992) Smear and culture positive cases of tuberculosis found among the symptomatic in Wardha district. Ind. J. Tub. 1992; 39: 159

Reference.

- 106. Narayan R at el (1979) A sociological study of awareness of symptoms and action taking of person with Pulmonary TB (A resurvey) Ind. J. Tub. 1979; 26: 136-146.
- 107. Narayan R et al (1976) Prevalence of chest symptoms and action taken by symptomatic in a rural community. Ind. J. Tub. 1976; 23: 160-168.
- 108. NTI (1994) District Tuberculosis Programme Introduction to DTP
 4th edition. Govt. of India, National TB Institute Bangalore 1994:
 1-12.
- 109. NTI (1997) Performance of National TB Control programme of India for year 1996. NTI Bulletin 1997; 33(1/2): 19-20.
- 110. NTI Bangalore. (1986) District Tuberculosis Programme Introduction manual, NTI, January 1986.
- 111. O' Rourke J. (1965) Evaluating control programme. Ind J Tub, 1965; 12: 87.
- 112. Pamra S.P., Goyal S.S., Mathur G.P. (1973) Change in Prevalence and incidence of Pulmonary Tuberculosis in recent years. Ind. J. Tub. 1973; 20: 57.
- 113. **Pamra SP. (1987)** Tuberculosis Control in India, 1987; WHO/TB. 149.
- 114. Pathania V, Almedia J. (1997) TB patients and private non-profit health care providers in India. 1997.WHO/TB/97.233.
- 115. Paul R (1961) Silicosis in Northern Rhodesia copper minors. Arch. Environ. Health 1961; 2: 96-109.

_*

- 116. Prajapati PB (1994) A study of District Tuberculosis Programme with reference to treatment defaults and effectiveness of retrieval actions. A MD Thesis submitted to MS University of Baroda Year 1994.
- 117. Purohit SD et al (1988) Awareness about TB among general population- a pilot study. Ind. J. Tub. 1988; 35: 183-187.
- 118. Radha Narayan at el. (1976) Prevalence of chest symptoms and action taken by symptomatic in a rural community. Ind J Tub, 1976; 23: 160.
- 119. Radha Narayan at el. (1982) Illness perception and medical relief in rural communities. Ind J Tub, 1982; 29: 98.
- 120. Radha Narayan at el. (1983) A study of tuberculosis services a component of primary health care. Ind J Tub, 1983; 30: 69.
- 121. Radha Narayan at el. (1987) Study of utilization of general health and tuberculosis services by a rural community. NTI Newsletter, 1987;23(4): 91.
- 122. Radha Narayan, Srikantaramu N. (1981) Symptoms awareness and action taking of persons with pulmonary tuberculosis in rural communities surveyed repeatedly to determine the epidemiology of the disease. Ind J Tub, 1981; 28: 125.
- 123. Radha Narayan, Srikantaramu N. (1987) Significance of some social factors in the treatment behavior of tuberculosis patients. NTI Newsletter, 1987; 23(4): 76
- 124: Radha Nrayan at el. (1979) A sociological study of awareness of symptoms and action taking of persons with pulmonary tuberculosis. Ind J Tub, 1979; 26: 136.

125. Radhakrishna S (1983) National TB Programme: Relative merits of enhancing the operational efficiency of different component of the treatment programme. Ind. J. Tub. 1983; 30: 3-8.

~

- 126. Rajeswari R. at el (1999) Socio- economic impact of tuberculosis on patients and family in India. Int J Tuberc Lung Dis 1999; 3: 869-877.
- 127. Ramana GNY, Naidu BMCS, Murthy KJR (1997) Mapping of TB treatment providers at selected sites in Andhra Pradesh State, India. 1997 .WHO/TB/97.233.
- 128. Rao K.N. (1981) Textbook of Tuberculosis (2nd Edition) The TB association of India New Delhi. 1981; 50-52.
- 129. Rao KN (1981) Tuberculosis in relation to Diabetes. Text Book of Tuberculosis, 2nd edition TB association of India 1981: 252-257.
- 130. Rieder H.L. (1999) Epidemiological basis of Tuberculosis Control. (First Edition) IUAT and LT 1999; 72-74.
- 131. Saiyad HN (1987) Study of Respiratory morbidity in agate worker. NIOH annual report 1987-88.
- 132. Sarin R, Day LBS. (1995) Indian National Tuberculosis Programme: Revised Strategy. Ind J Tub, 1995; 42(2): 95.
- 133. Saxena P, Mathur GP, Singh MM. (1987) Treatment taken before reporting at a tuberculosis clinic. Ind J Tub, 1987, 34: 104.
- 134. Seetha MA. (1976) Changing role of paramedical worker in the fight against tuberculosis. NTI Newsletter, 1976; 13: 40.

135. Selwyn PA at el (1989) A prospective study of the risk of TB among intravenous drugs user with HIV infection. N. Engl. J. Med. 1989; 320: 545-550.

*

- 136. Seetha MA, Aneja KS. (1977) Defaulter retrieval and cast of defaulter actions at Peripheral Health Institutions. NTI Newsletter, 1977; 14: 87.
- 137. Saxena P et al (1987) Treatment taken before reporting TB clinic. Ind. J. Tub. 1987; 34: 104-107.
- 138. Shivram S. (1981) Tuberculosis in India- The prospect. Ind. J. Tub. 1981, 29(2): 79-86.
- 139. Snider DE (1978) The relationship between tuberculosis and silicosis (Editorial) Am. Rev. Respir. Dis. 1978; 118: 455-560.
- 140. Stead W.W., Lafgreen J.P. (1983) Does risk of tuberculosis increases in old age? J.Infect. Dis. 1983; 147: 951-955.
- 141. Styblo K. (1984) Epidemiology of tuberculosis. VEB Gustav Fisher Verlag Jena: 1984.
- 142. Subramanion T. (1990) Sample survey of awareness of symptoms and utilization of health factionaries by chest symptomatics. Ind. J. Tub. 1990; 37: 67-71.
- 143. Tejzak EE (1997) Tuberculosis and Human immunodeficiency virus infection. Med. Clin. North. Am. 1997; 81: 345-360.
- 144. Thomas B.E. at el (1999) Women and tuberculosis Ind. J. Tub. 2000; 137: 177-198.

Reference.

- 145. TRC Madras (1980) Tuberculosis prevention trial of BCG vaccine in south India for tuberculosis prevention. Ind. J. Med. Res. 1980; 72: 72 (supplement) 1.
- 146. Tuberculosis Chemotherapy Centre, Madras. (1964) A concurrent comparison of intermittent (twice- weekly) Isoniazid plus streptomycin and daily Isoniazid plus PAS in the domiciliary treatment of pulmonary tuberculosis. Bull World Health Organ 1964; 31:247-271.
- 147. Upleker M, Rangan S. (1996) TB: The search for solutions. Bombay The foundation for research in community health .1996.
- 148. Whalen C at el (1995) Accelerated course of HIV infection after tuberculosis. Am. J. Resp. Crit. Care. Med. 1995; 151: 29-35.
- 149. WHO (1992) Tuberculosis Programme Review: India, September 1992.
- 150. WHO (1997) Report on the Tuberculosis epidemic 1997; WHO/TB/ 97; 224.
- 151. WHO (1999) World Health Report 1999; WHO/99/12368.
- 152. WHO (2000) Global TB Report 2000, WHO/CDS/TB 2000; 275. 23
- 153. WHO (2000) Joint Tuberculosis Programme Review, India.February 2000 WHO (SEA) /TB/224/ 9-13.
- 154. WHO (2000) Research for Action Understanding and Controlling Tuberculosis in India? WHO (SEA) New Delhi 2000; 12-20.
- 155. World Bank (1995) India: Policy and Finance strategies for strengthening primary health care services. World Bank Report No. 13042-In. 1996.

- 156. World Health Organization. (1996) Group at risk: report on the tuberculosis epidemic 1996. WHO/TB/96.
- 157. World Health Organization. (1997) Report on tuberculosis epidemic. 1997. WHO/TB/97.224.
- 158. World Health Organization. (1999) World Health Report 1999. WHO/99/12368.
- 159. Yu G at el (1988) Risk Factors associated with the pulmonary tuberculosis among sanitary worker in Shanghai. Tubere. 1988; 69: 105-112.
- 160. Zhang LX, Kan GQ (1992) Tuberculosis control programme in Beijing. Tuberc Lung Dis 1992; 73:162-166.

1
