

## LIST OF GRAPHS

<b>Graph No.</b>	<b>Title of Graph</b>	<b>Page No.</b>
<b>4.1</b>	K/S values of samples bleached with Sodium Hypochlorite and dyed with madder using different mordants	<b>80</b>
<b>4.2</b>	K/S values of samples bleached with Hydrogen peroxide and dyed with madder using different mordants	<b>81</b>
<b>4.3</b>	K/S values of samples bleached with Peracetic acid and dyed with madder using different mordants	<b>82</b>
<b>4.4</b>	K/S values of samples bleached with Sodium Hypochlorite dyed with turmeric using different mordant	<b>86</b>
<b>4.5</b>	K/S values of samples bleached with Hydrogen peroxide dyed with turmeric using different mordant	<b>87</b>
<b>4.6</b>	K/S values of samples bleached with Peracetic acid dyed with turmeric using different mordant	<b>88</b>
<b>4.7</b>	K/S values of samples bleached with Sodium Hypochlorite dyed with Eucalyptus Leaves using different mordants	<b>92</b>
<b>4.8</b>	K/S values of samples bleached with Hydrogen Peroxide dyed using Eucalyptus Leaves using different mordants	<b>92</b>
<b>4.9</b>	K/S values of samples bleached with Peracetic acid dyed with Eucalyptus Leaves using different mordant	<b>93</b>
<b>4.10</b>	K/S values of samples bleached with Sodium Hypochlorite dyed with Indian Almond Leaves using different mordants	<b>97</b>
<b>4.11</b>	K/S values of samples bleached with Hydrogen Peroxide dyed with Indian Almond Leaves using different mordants	<b>97</b>
<b>4.12</b>	K/S values of samples bleached with Peracetic acid dyed with Indian Almond Leaves using different mordants	<b>98</b>
<b>4.13</b>	Optical indices of jute fabric bleached with different concentration of Hydrogen Peroxide	<b>107</b>
<b>4.14 (a)</b>	Optical indices of jute fabric bleached with Peracetic acid at pH 6	<b>110</b>
<b>4.14 (b)</b>	Optical indices of jute fabric bleached with Peracetic acid at pH 7	<b>110</b>
<b>4.15</b>	FTIR graph for only woolenized samples	<b>113</b>
<b>4.16</b>	FTIR graph for raw jute samples	<b>114</b>
<b>4.17</b>	FTIR graph for samples for first Woolenized and then bleached with H <sub>2</sub> O <sub>2</sub>	<b>114</b>
<b>4.18</b>	FTIR graph for samples first Woolenized and then bleached with	<b>115</b>

	PAA	
<b>4.19(a)</b>	FTIR graph for samples bleached with H <sub>2</sub> O <sub>2</sub> and then Woolenized	<b>115</b>
<b>4.19(b)</b>	FTIR graph for samples bleached with PAA and then Woolenized	<b>116</b>
<b>4.20</b>	Particle size distribution of synthesized nano copper	<b>119</b>