LIST OF FIGURES

Fig. No.	Title	Page No.
1.1	Diabetes as Global Health Emergency of 21st century by IDF 2019	3
1.2	Number (million) of adults with diabetes for various countries with the diabetes-related health expenditure by IDF 2019	4
1.3	The prevalence rates of diabetes among adults at the national level and per state and union territory	5
2.1	Metabolic actions of insulin in striated muscles, adipose tissue, and liver	19
2.2	Pathogenesis of type 2 diabetes	21
2.3	Estimated total number of adults (20–79 yrs) with diabetes in 2019	23
2.4	The % prevalence of overweight, obesity, hypertension and diabetes among adults by (a) disease/condition, (b) age group,(c) age group and sex, and (d) age group, sex, and residence	24
2.5	The rising prevalence of diabetes in india for last twenty years	26
2.6	Prevalence of diabetes in India by national diabetes and diabetic retinopathy survey report (2019)	27
2.7	Galega Officinalis /French Lilac Or Goat's Rue: Herbal Source of Metformin	29
2.8	Glycemic Management of Type 2 Diabetes Mellitus	34
2.9	Functions of vitamin B12 in human body	45
2.10	Mechanism of diabetic neuropathy	58
2.11	Semmes Weinstein Monofilament (SWM) for sensory testing	62
2.12	Achilles reflex	75
2.13	Jendrassic manoeuver	75
2.14	Neurothesiometer	77
3.1	Experimental design	100

3.2	Physical Assessment of foot -Flat feet	112
3.3	Physical Assessment of foot - Hammer toes	112
3.4	Physical Assessment of foot - Overlapping toes	113
3.5	Physical Assessment of foot - Hallus valgus	113
3.6	Physical Assessment of foot - Charcot foot	113
3.7	Tools used for MNSI: hammer (top), 128 Hz tunning fork (middle), 10g monofilament (bottom)	116
3.8	Jendrassic maneuver performed by the subject	117
3.9	Assessing DPN of the study subject by ankle reflex using hammer	118
3.10	Assessing DPN of the subject by vibration sensation using 128 Hz tuning fork	119
3.11	Assessing DPN of the subject by monofilament testing using 10g monofilament	120
3.12	Vitamin B12 combination supplements available in the market survey rejected for the study	123
3.13	Mecobal - vitamin B12 tablet with sole B12 formulation selected for supplementation	124
3.14	Cal 360: the calcium tablet used for supplementation	124
3.15	Supplementation monitored by counting wrappers of the medicine consumed	125
4.1	Percent distribution of PCI of T2DM adults on metformin	130
4.2	Percentage distribution of various occupations among T2DM adults on metformin	130
4.3	Prevalence of various stages of hypertension from gender perspective	137
4.4	'Yes' response to MNSI history questions for T2DM adults on metformin	143
4.5	Prevalence of DPN among T2DM adults on metformin	144
4.6	Percent prevalence of DPN from gender perspective	145

4.7	Percent prevalence of various grades of DPN among T2DM adults on metformin	145
4.8	Percent distribution of gender among various Grades of DPN	146
4.9	Prevalence of various grades of DPN across gender	146
4.10	Comparison of mean domain scores of QoL among T2DM adults with and without DPN	163
4.11	Box plot for serum B12 distribution of T2DM adults on metformin	170
4.12	Grades of B12 deficiency among T2DM adults Across gender	172
4.13	Grades of anemia among T2DM adults on metformin across gender	173
4.14	Prevalence of anemia among those on iron supplementation and non-iron supplementation	173
4.15	Vegetarianism versus non-vegetarianism across B12 deficiency	176
4.16	Comparison of distribution of serum B12 across present metformin dosage	178
4.17	Comparison of distribution of serum B12 across duration of diabetes	186
4.18	Frequencies of median serum B12 in three groups with increasing duration of diabetes among T2DM adults on metformin (N=155)	187
4.19	Frequencies of median Serum B12(without outliers) in three groups with increasing duration of diabetes among t2dm adults on metformin (N=137)	188
4.20	Prevalence of Serum B12 deficiency by DPN grades	191
4.21	Receiver Operating Characteristic Curve (ROC) For diabetic peripheral neuropathy score (DPN score) related to vitamin B12 deficiency	193
4.22	Prevalence of good glycemic control (HbA1c≤7) and poor glycemic control (HbA1c >7%) by grades of DPN	195
4.23	Quality of life mean domain scores across B12 deficiency	196
4.24	Post supplementation prevalence of B12 deficiency and normal	211

	B12 in between two groups	
4.25	Comparison of pre and post supplementation MNSI physical assessment score of Right Foot	227
4.26	Comparison of pre and post supplementation MNSI scores of physical assessment of left foot	228
4.27	Comparison of impact of supplementation in mean difference (post-pre) of serum B12, HbA1c, DPN score, MNSI history score and four domains of Qol in between the two supplementation groups	260