## LIST OF TABLES

Table no.	Title	Page no.
2.1.1	Production of Banana in India	9
2.1.2	International production of Banana	9
2.1.3	Production of Mandarion (M. Orang, Kinniw, Orange) in India	10
2.1.4	International Production of Oranges	10
2.1.5	Production of Peas in India	11
2.1.6	International Production of Peas	11
2.1.7	Production of Maize in India	12
2.1.8	International production of Maize	12
2.2	Number of Panellists and samples required for sensory test	35
5.2	XOS yield by enzymatic hydrolysis of Xylan at different incubation and pH-5.5 at 40°C	71
5.3	Concentration of XOS among all the four agro wastes	71
5.4	XOS recovery at different levels of pH	77
5.5	Mean growth and SD of <i>Lactobacillus plantarum</i> , <i>Bifidobacterium adolescentis</i> and <i>Escherichia coli</i> with 0.5%, 1% and 2% levels of XOS	97
5.6	Mean growth and SD of <i>Lactobacillus plantarum</i> , <i>Bifidobacterium adolescentis</i> and <i>Escherichia coli</i> with 0.5%, 1% and 2% levels of XOS	98
5.7	Mean and SD values of acetate, butyrate and propionate produced by <i>Lactobacillus plantarum</i> , <i>Bifidobacterium adolescentis</i> and <i>Escherichia coli</i> on its exposure to XOS and glucose being used as control group	102
5.8.1	Organoleptic properties of Paneer Butter Masala	109

5.8.2	Organoleptic properties of <i>Prawn patia</i>	111
5.8.3	Organoleptic properties of Black rice kheer	113
5.8.4	Organoleptic properties of Gajar ka halwa	115
5.9.1	Number of panel members indicating the difference in the organoleptic attributes of <i>Paneer Butter Masala</i> in a difference test	118
5.9.2	Number of panel members indicating the difference in the organoleptic attributes of <i>Prawn patia</i> in a difference test	119
5.9.3	Number of panel members indicating the difference in the organoleptic attributes of <i>Black rice kheer</i> in a difference test	120
5.9.4	Number of panel members indicating the difference in the organoleptic attributes of <i>Gajar ka halwa</i> in a difference test	121
6.1	XOS obtained from 60 g dried powder	135
6.1.1	XOS obtained from 100 g dried powder	135