

LIST OF TABLES

| TABLE NO. | TITLE | PAGE NO. |
|------------------|--|-----------------|
| 2.1 | Classification for Identification of Impaired Glucose Tolerance and Diabetes | 15 |
| 2.2 | Guidelines for Asians (Misra et al 2009) | 74 |
| 2.3 | Guidelines by WHO 1997 | 74 |
| 2.4 | JNC VII Classification for Blood Pressure | 79 |
| 2.5 | ATP III Guidelines for Dyslipidaemia | 82 |
| 4.1 | Score card for “Worksite Policies and Environment” in the Selected Workplace | 135 |
| 4.2 | Score card for “Health Promotion for Employees” in the Selected Workplace | 136 |
| 4.3 | Background Information of the Subjects in the Selected Workplace (n=504) | 148 |
| 4.4 | Addiction Pattern of the Subjects in the Selected Workplace (n=504) | 150 |
| 4.5 | Physical Activity Pattern of the Subjects in the Selected Workplace (n=504) | 152 |
| 4.6 | Dietary Pattern of the Subjects in the Selected Workplace (n=504) | 156 |
| 4.7 | Dietary Pattern of the Subjects in the Selected Workplace (n=504) | 157 |
| 4.8 | Anthropometric Profile of the Subjects in the Selected Workplace (n=504) | 159 |
| 4.9 | Medical History and Current Blood Pressure Profile of the Subjects in the Selected Workplace (n= 504) | 162 |
| 4.10 | Body Mass Index (BMI) & Blood Pressure profile based on gender (Mean \pm S.D.) | 166 |
| 4.11 | Anthropometric Profile, Hypertension Status & Physical Activity Status of Subjects in Relation to Gender (n=504) | 167 |

| | | |
|------|--|-----|
| 4.12 | Differences in Anthropometric Profile, Hypertension Status & Physical Activity Status of Subjects in Relation to Age (n=504) | 168 |
| 4.13 | Difference in Mean Age, BMI and B.P. of Employees Based on Designation in the Selected Workplace (n=504) | 170 |
| 4.14 | Anthropometric & Blood Pressure Profile of the Subjects in Relation to Smoking & Alcohol Consumption (Mean \pm S.D.) | 171 |
| 4.15 | Anthropometric & Blood Pressure Profile of the Subjects in Relation to their BMI Status (n=504) | 172 |
| 4.16 | Hypertension Status of Subjects in Relation to Abdominal Obesity (n=504) | 173 |
| 4.17 | Anthropometric Profile, Hypertension Status, Medical History of Hypertension & Designation level of Subjects in Relation to their Physical Activity Status (n=504) | 175 |
| 4.18 | Verbatim Responses Received from the Subjects in the Selected Workplace | 191 |
| 4.19 | Percent Prevalence of Dyslipidaemia & Abnormal Lipid Profile Fractions among the Consenting 'At Risk' Subjects (n=504) | 191 |
| 4.20 | Biochemical Profile of Consenting 'At Risk' Subjects (Mean \pm S.D.) | 193 |
| 4.21 | Fasting Blood Sugar Profile of 'Consenting 'At Risk' Subjects (n=158) | 194 |
| 4.22 | Percent prevalence of Metabolic Syndrome (MS) and percent distribution of MS risk factors among the Consenting 'At Risk' Subjects (n=158) | 198 |
| 4.23 | Differences in Mean Cholesterol & Fasting Blood Sugar Values of Male and Female Subjects (n=158) | 199 |
| 4.24 | Differences in Mean Cholesterol & Fasting Blood Sugar Values of Smokers and Non Smokers (n=158) | 200 |
| 4.25 | Differences in Mean Cholesterol & Fasting Blood Sugar Values of Physically Active & Physically Inactive Subjects (n=158) | 201 |
| 4.26 | Knowledge, Attitude, Practices and Perceptions of Subjects Regarding Maintenance of Vital Health Parameters (n=126) | 210 |
| 4.27 | Knowledge, Attitude, Practices and Perceptions of Subjects Regarding Maintenance of Healthy Blood Pressure (n=126) | 213 |

| | | |
|------|---|-----|
| 4.28 | Knowledge, Attitude, Practices and Perceptions of Subjects Regarding Maintainance of Healthy Cholesterol Levels (n=126) | 215 |
| 4.29 | Knowledge, Attitude, Practices and Perceptions of Subjects Regarding Maintainance of Healthy Blood Sugar Levels (n=126) | 217 |
| 4.30 | Knowledge, Attitude, Practices and Perceptions of Subjects Regarding Tobacco and Alcohol Consumption (n=126) | 221 |
| 4.31 | Percent Prevalence of Healthy Practices among the Subjects (n=126) | 222 |
| 4.32 | Self-Reported v/s Actual Prevalence of Healthy Practices among the Subjects (n=126) | 224 |
| 4.33 | Impact of Nutrition Health Promotion Programme Intervention on the Knowledge, Attitude and Practices of Subjects Regarding Maintainance of Vital Healthy Parameters (n=83) | 233 |
| 4.34 | Impact of Nutrition Health Promotion Programme Intervention on the Knowledge of Subjects Regarding the Reasons for Weight Gain in Adulthood (n=83) | 234 |
| 4.35 | Impact of Nutrition Health Promotion Programme Intervention on the Knowledge, Attitude and Practices of Subjects Regarding Maintainance of Healthy Blood Pressure Levels (n=83) | 236 |
| 4.36 | Impact of Nutrition Health Promotion Programme Intervention on the Knowledge, Attitude and Practices of Subjects Regarding Maintainance of Healthy Cholesterol Levels i.e. Lipid Profile (n=83) | 239 |
| 4.37 | Impact of Nutrition Health Promotion Programme Intervention on the Knowledge, Attitude and Practices of Subjects Regarding Maintainance of Healthy Blood Sugar Levels (n=83) | 240 |
| 4.38 | Impact of Nutrition Health Promotion Programme Intervention on the Knowledge, Attitude and Practices of Subjects Regarding Tobacco Consumption (n=83) | 242 |
| 4.39 | Impact of Nutrition Health Promotion Programme Intervention on the Knowledge and Perception of Subjects Regarding Healthy Diet & Lifestyle Behaviours (n=83) | 243 |
| 4.40 | Scoring Method used for Knowledge Scores of Subjects Before and After Intervention (maximum score=6) | 245 |

| | | |
|------|---|-------------|
| 4.41 | Scoring Method used for Attitude Scores of Subjects Before and After Intervention (maximum score=2) | 246 |
| 4.42 | Scoring Method used for Evaluating Perception Scores of Subjects Before and After Intervention (maximum score=23) | 247 |
| 4.43 | Scoring Method used for Practice Scores of Subjects Before and After Intervention (maximum score=20) | 248- 249 |
| 4.44 | Overall Knowledge, Attitude, Practice & Perception Scores of Subjects Regarding Before & After Intervention (Mean \pm S.D.) n=83 | 250 |
| 4.45 | Changes in Average Fruit and Vegetable Consumption by the Subjects Before and After Intervention (n=83) | 256 |
| 4.46 | Differences in the Mean Values of Anthropometric Parameters and Blood Pressure Profile in the Study Population Before and After Intervention (n=83) Mean \pm S.D. | 260 |
| 4.47 | Change in the Percent Prevalence of 'At Risk' Population Before and After Intervention (n=83) | 263 |
| 4.48 | Differences in the Mean Values of Lipid Fractions as well as Fasting Blood Sugar in the Study Population Before and After Intervention (n=51) Mean \pm S.D. | 267 |

LIST OF FIGURES

| NO. | TITLE | PAGE NO. |
|------------|---|---------------------|
| 2.1 | Total Deaths by Broad Cause Group, by WHO Region, World Bank Income Group and by Sex, 2008 | 10 |
| 2.2 | Pattern of overall DALYs (age standardized) & NCD related DALYs in India, 2004 | 11 |
| 2.3 | Trend in mean fasting blood glucose of Indians from 1980-2008 (WHO 2010) | 18 |
| 2.4 | Interrelationship of Poverty and NCDs (Global status report on non communicable diseases 2010) | 19 |
| 2.5 | Growth of Urban Population since 1951-2011 in India | 21 |
| 2.6 | Nutrition Transition (Popkin's concept) | 24 |
| 2.7 | Average Dietary Energy Supply Adequacy in the Developing Regions, 1990-92 to 2011-13 | 25 |
| 2.8 | Rise in Dietary Energy Supplies (kcal/person/day) in the world (1990-1992 & 2008-2010) | 26 |
| 2.9 | Changes in Dietary Patterns in the world (1990-1992 and 2007-2009) | 27 |
| 2.10 | Global Prevalence of Tobacco related Diseases and Disabilities in 2005 (WHO Tobacco factsheet 2013) | 30 |
| 2.11 | Age-Standardized prevalence of daily tobacco smoking in adults aged 15+ years, by WHO Region and World Bank Group, comparable estimates, 2008 | 31 |
| 2.12 | Percentage of Adults Age 15 and Above by Detailed Status of Tobacco Use, According to Gender and Residence, GATS India, 2009-2010 | 33 |
| 2.13 | Percentage of Current Tobacco Users According to States/UTs, GATS India, 2009-2010 | 34 |
| 2.14 | Tobacco usage in India (percentage of adults 15+ years) during 2009-2010 | 35 |

| | | |
|------|---|-----|
| 2.15 | Percent Distribution of Tobacco Use According to Gender, GATS India, 2009-2010 | 37 |
| 2.16 | Study results within a country before and after a ban on advertising | 45 |
| 2.17 | Physical Activity prescription for aerobic & muscle strengthening exercises | 46 |
| 2.18 | Percent of total energy from fat and percent energy from saturated fatty acids in countries, based on World Bank income groups (2008) | 56 |
| 2.19 | Global percentages of DALYs attributed to 19 leading risk factors by income group | 62 |
| 2.20 | Health consequences of alcohol usage in India (2004) | 63 |
| 2.21 | Population weighted means of the recorded adult per capita consumption in the WHO region (1961-1999) | 63 |
| 2.22 | India: trend of alcohol consumption (2000-2007) | 64 |
| 2.23 | Countries reporting awareness campaigns on alcohol, 2008 | 71 |
| 2.24 | Age-Standardized prevalence of overweight in adults aged 20+ years, by WHO Region and World Bank Income group, 2008 | 75 |
| 2.25 | Infant and young child overweight trends from 1990 to 2015, by World Bank Income group | 76 |
| 2.26 | Trends in mean BMI of Indians (1980-2008) | 79 |
| 2.27 | Age-Standardized prevalence of raised blood pressure in adults aged 25+ years, by WHO Region and World Bank Income group, comparable estimated (2008) | 85 |
| 2.28 | Trend in mean systolic blood pressure of Indians (1980-2008) | 86 |
| 2.29 | Proportion of countries with funding for NCD activities, by function, WHO Region and World Bank income group, 2010 | 108 |
| 2.30 | National Capacity for Prevention and Control of NCDs 2010 | 109 |
| 3.1 | Experimental Design | 122 |
| 4.1 | Percent Prevalence of Unhealthy Behaviours and Negative Health Status among the Subjects in the Selected Workplace (n=504) | 164 |
| 4.2 | Percent Prevalence of subjects found to be "at risk" by WHO STEPS Methodology (n=504) | 187 |

| | | |
|------|--|-----|
| 4.3 | Distribution of Risk Factors Among Total Study Population (n=504) | 188 |
| 4.4 | Knowledge of subjects regarding reasons for weight gain in adulthood (n=126) | 219 |
| 4.5 | Impact of Nutrition Health Promotion Programme on Knowledge about Key Health Behaviours (n=83) | 252 |
| 4.6 | Percentage of Subjects who Adopted the Recommended Healthy Behaviours (n=83) | 253 |
| 4.7 | Percent Prevalence of Unhealthy Behaviours and Negative Health Status among Study Population Before & After Intervention (n=83) | 255 |
| 4.8 | Percent Prevalence of Unhealthy Behaviours and Negative Health Status among Male Subjects Before and After Intervention (n=72) | 258 |
| 4.9 | Percent Prevalence of Unhealthy Behaviours and Negative Health Status among Female Subjects Before and After Intervention (n=11) | 259 |
| 4.10 | Change in Percent Prevalence of Dyslipidaemia, Lipid Fractions & Impaired Glucose Tolerance among Study Population Before & After Intervention (n=51) | 264 |
| 4.11 | Change in Percent Prevalence of Dyslipidaemia, Lipid Fractions & Impaired Glucose Tolerance among Male Population Before & After Intervention (n=45) | 265 |
| 4.12 | Change in Percent Prevalence of Dyslipidaemia, Lipid Fractions & Impaired Glucose Tolerance among Female Population Before & After Intervention (n=6) | 266 |
| 5.1 | Percent Prevalence of Risk Factors among Study Subjects at Baseline (n=504) | 283 |
| 5.2 | Percent Prevalence of Subjects who Attended Nutrition Health Promotion Programme Sessions Regarding Knowledge & Perception of Key Nutrition & Health Issues at Baseline (n=126) | 287 |
| 5.3 | Percent Prevalence of Healthy Practices Followed by Subjects who Attended the Nutrition Health Promotion Programme Sessions at Baseline (n=126) | 288 |
| 5.4 | Percent Prevalence of Subjects who Attended Nutrition Health Promotion Programme Sessions Exhibiting Healthy Attitudes Regarding Adoption of Healthy Behaviours at Baseline (n=126) | 289 |
| 5.5 | Percent Increase in Subjects who Attended Nutrition Health Promotion Programme Sessions Regarding Knowledge & Perceptions about Key Nutrition & Health Issues Before & After Intervention (n=83) | 294 |

| | | |
|-----|--|-----|
| 5.6 | Percent Increase in Subjects who Attended the Nutrition Health Promotion Programme Sessions Before & After Intervention Regarding Following of Healthy Practices - Self Reported Prevalence (n=83) | 295 |
| 5.7 | Prevalence of Subjects having Unhealthy Behaviors and Negative Health Status Before and After Intervention (n=83) | 296 |
| 5.8 | Change in Percent Prevalence of Dyslipidaemia, Lipid Fractions & Impaired Glucose Tolerance among Study Population Before & After Intervention (n=51) | 297 |

APPENDICES

| SR. NO. | TITLE |
|----------------|---|
| 1 | Employee Consent Letter |
| 2 | Worksite Wellness Index Questionnaire |
| 3 | Modified WHO STEPS Questionnaire (for Baseline Data Collection) |
| 4-A | Power point presentation session slides (Presentation 1) |
| 4-B | Power point presentation session slides (Presentation 2) |
| 5 | A3 size colored posters |
| 6 | A4 size black & white table mats |
| 7 | Point-of-Decision Prompts |
| 8 | Knowledge Attitude Practice (KAP) Questionnaire (for Baseline Data Collection) |
| 9 | Modified WHO STEPS Questionnaire (for Post Intervention Data Collection) |
| 10 | Knowledge Attitude Practice (KAP) Questionnaire (for Post Intervention Collection) |