

LIST OF TABLES

Table no.	TITLE	Page no.
2.1	Classification of weight status according to BMI in Asian adults	29
2.2	Waist circumference cut-off points for increased risk to metabolic	32
4.1	Score Cut-offs for Socio Economic Status	108
4.2	List of seven types of stool which was shown to each subject	110
4.3	Score card for rating defecation pattern	110
4.4	Score cutoffs for type of defecation profile	111
4.5	Score cutoffs for defining degree of constipation	111
4.6	Criteria for - Days and Total Physical Activity Met/Min/Week	113
4.7	Hunger and Satiety Scale	114
4.8	Hunger and Satiety Score Card for Bank Employees	114
4.9	Score Cut-offs for Becks Depression Inventory	115
4.10	Classification of weight status according to BMI in Asian adults	119
4.11	Cut Offs for WC, WHR and WSR for CVD risk	119
4.12	JNC VIII - Classification of Blood Pressure in Adults [Age ≥ 18 years]	120
4.13	Gender-Wise Classification of Body Fat Percentage	121
4.14	MILLIPLEX Analyst Detail Report	134
4.15	Specification and Technical Aspects of FOS Used for Product Development	146
4.16	List of Raw Ingredients Along With Their Sources Used For Product Preparation	147
5.1	Distribution of screened bank employees according to gender [N=650]	161
5.2	Distribution of screened bank employees according to various categories of age	161

Table no.	TITLE	Page no.
5.3	Mean values of anthropometric profile of screened bank employees [N=650]	163
5.4	Mean values for biophysical and biochemical profile of screened bank employees	163
5.5	Distribution of screened bank employees according to Asia pacific BMI classification for Indians [JAPI 2009 & WHO 2000] N=650	165
5.6	Mean values of anthropometric parameters of male bank employees according to BMI classification [N=486]	167
5.7	Mean values of biophysical parameters of male bank employees according to BMI classification [N = 486]	167
5.8	Mean values of anthropometric parameters of female bank employees according to BMI classification [N = 164]	168
5.9	Mean values of biophysical parameters of female bank employees according to BMI classification [N = 164]	168
5.10	Presence of abdominal obesity according to waist circumference [WC] in bank employees subjected to screening	169
5.11	Presence of abdominal obesity according to waist stature ratio [WSR] in bank employees subjected to screening	170
5.12	Presence of central obesity according to waist hip ratio [WHR] in bank employees subjected to screening	171
5.13	Distribution of screened bank employees according to body fat classification	172
5.14	Presence of hypertension in bank employees subjected to screening [N=650]	173
5.15	Association of Body fat with Anthropometric and Biophysical parameters N=650	175
5.16	Percent presence of elevated SBP along-with its coexistence with multiple deranged anthropometric parameters as risk factors for development of NCD's and association amongst them N = 650 [%]	176
5.17	Percent presence of elevated DBP along-with its coexistence with multiple deranged anthropometric parameters as risk factors for development of NCD's and association amongst them N = 650 [%]	177
5.18	Percent presence of hypertension along with its coexistence with multiple deranged anthropometric parameters as risk factors for development of NCD's and association amongst them N = 469 [%]	178
5.19	Background information of non-obese and obese subjects	186
5.20	Mean values for anthropometric parameters of non-obese [150] and obese [150] employees	187

Table no.	TITLE	Page no.
5.21	Mean values for biophysical parameters of non-obese [150] and obese [150] subjects	188
5.22	Mean values for blood pressure measurements of non-obese [150] and obese [150] subjects	188
5.23	Percent presence and association of abdominal obesity [WC] with BMI	189
5.24	Percent presence and association of central obesity [WHR] with BMI	189
5.25	Percent presence of level of fitness according to body-fat %	190
5.26	Odds ratio and relative risk as per level of fitness	190
5.27	Association of BMI with physical activity level in non-obese and obese bank employees	191
5.28	Family history of NCD's in non-obese and obese bank employees and its association with BMI	193
5.29	Personal medical history of non obese and obese bank employees	193
5.30	Frequency of habitual products consumed by non-obese and obese bank employees	195
5.31	Percent consumption of habitual products and its association with obesity in non-obese and obese bank employees	195
5.32	Association of BMI with varying degree of habituation in non-obese and obese subjects	197
5.33	Mean values of anthropometric and biophysical profile of <u>NON-OBESE</u> employees according to their degree of habituation	197
5.34	Percent presence of depression and its association with BMI of non-obese and obese bank employees	198
5.35	Defecation profile of non obese and obese subjects	200
5.36	Defecation pattern according to score analysis and degree of constipation in non-obese and obese bank employees	201
5.37	Frequency of major food group consumption by non-obese and obese bank employees	202
5.38	Mean values of macronutrient and types of fiber intake of non-obese and obese bank employees	204
5.39	Mean hunger scores of non-obese and obese subjects at various meal timings	206
5.4	Mean satiety scores of non-obese and obese subjects at various meal timings	206
5.41	Mean values of gut-microbial profile and difference in their gut colonization pattern of non-obese and obese bank employees	207

Table no.	TITLE	Page no.
5.42	Baseline fasting gut-hormone [pg/ml] profile of non-obese and obese young bank employees	209
5.43	Positive correlation of weight with various parameters of young bank employees of urban Vadodara	210
5.44	Negative correlation of various parameters with weight of young bank employees of urban Vadodara	211
5.45	Stepwise regression model summary for strongest predictor of obesity in young bank employees of urban Vadodara	213
5.46	Mean anthropometric values of obese bank employees before and after intervention	232
5.47	Mean biophysical values of obese bank employees before and after intervention	233
5.48	Mean dietary intakes of obese bank employees before and after intervention	235
5.49	Mean hunger scores of obese bank employees before and after intervention	236
5.50	Mean satiety scores of obese bank employees before and after intervention	237
5.51	Mean depression scores of obese bank employees before and after intervention	238
5.52A/B	Association between defecation profile of obese bank employees before and after intervention	239 – 240
5.53	Association between flatulence profile of obese bank employees before and after intervention	244
5.54	Mean values of fasting plasma gut hormone values of obese subjects before and after intervention	243
5.55	Gut-microbial profile of obese bank employees before and after intervention	244
5.56	Pearson correlation of weight with various parameters of obese bank employees [r value]	245
5.57	Pearson correlation of BMR with various parameters of obese bank employees [r value]	246
5.58	Pearson's correlation of Gut-hormones with various parameters	247
5.59	Pearson's correlation of Gut-microbes with various parameters	248
5.6	Stepwise regression model summary for strongest predictor of obesity in obese bank employees of urban Vadodara	249
5.61	Time point interval difference in anthropometric and biophysical parameters of obese employees in experimental group [N=72]	252
5.62	Weight-loss in obese bank employees according to days of compliance [N = 72]	253

Table no.	TITLE	Page no.
5.63	Physical characteristics of steamed <i>Dudhi Muthiya</i> with FOS addition at varying levels [One Serving]	266
5.64	Physical characteristics of shallow fried <i>Vegetable Cheela</i> with FOS addition at varying levels [One Serving]	268
5.65	Physical characteristics of baked <i>Handwa</i> with FOS addition at varying levels	271
5.66	Physical characteristics of deep fried <i>Vegetable Samosa</i> with FOS addition at varying levels	272
5.67	Effect of varying levels of Fructooligosaccharide addition on the organoleptic quality of steamed <i>Dudhi Muthiya</i>	275
5.68	Effect of varying levels of Fructooligosaccharide addition on the organoleptic quality of shallow fried <i>Vegetable Cheela</i>	279
5.69	Effect of varying levels of Fructooligosaccharide addition on the organoleptic quality of baked <i>Handwa</i>	283
5.70	Effect of varying levels of Fructooligosaccharide addition on the organoleptic quality of deep fried <i>Vegetable Samosa</i>	285