

APPENDIX

CHAPTER I

Figure No.	Caption to Figures	Pg. No.
1.1	Structures of zirconium phosphate	15
1.2	Idealized portion of ZP layer [25]	15
1.3	Schematic diagram of the arrangement of two adjacent macroanions in α -ZP [16]	16
1.4	(a) α -Ti(HPO ₄) ₂ .H ₂ O Octahedra represent TiO ₆ , dark grey tetrahedra represent HPO ₄ and spheres represent water molecules.[d ₀₀₂ =7.6Å][29]	17
	(b) γ -r(PO ₄)(H ₂ PO ₄) ₂ .2H ₂ O Octahedra represent ZrO ₆ , dark grey tetrahedra represent PO ₄ and light grey tetrahedra represent H ₂ PO ₄ and spheres represent water molecules. [d ₀₀₂ =12.2Å]	17
Table No.	Caption to Tables	Pg. No.
1.1	Separation methods	1
1.2	Principle classes of inorganic ion exchangers	10

CHAPTER II

Figure No.	Caption to Figures	Pg. No.
2.1	Schematic representation of the arrangement of layers in Zirconium phosphonates [22]	33
2.2	Structure of Zirconium phosphonates [23]	33
2.3	Scheme of a layered diphosphonate in which both phosphonic groups are bonded to the same face of the inorganic layer [6]	35
2.4	(a) structure of ATMP (b) structure of phosphoric acid	40
2.5	pH titration curves for (A) ZrATMP and (B) TiATMP	48
2.6	(a-d) Concentration and eluent volume optimization for determination of CEC using ZrATMP and TiATMP	50
2.7	(a) EDX; (b) FTIR; (c) TGA; (d) XRD and (e) SEM of ZrATMP	55
2.8	(a) EDX; (b) FTIR; (c) TGA; (d) XRD and (e) SEM of TiATMP	57
2.9	Ion exchange equilibrium [63]	58
2.10	Plot of fractional attainment of equilibrium for Cu ²⁺ -H ⁺ exchange vs. time using ZrATMP	72

2.11	Plot of t vs. time for $\text{Cu}^{2+}\text{-H}^+$ exchange for 0.014 M concentration using ZrATMP	75
2.12	Plots of t vs. time at different temperatures for (a) $\text{Co}^{2+}\text{-H}^+$ (b) $\text{Ni}^{2+}\text{-H}^+$ (c) $\text{Cu}^{2+}\text{-H}^+$ (d) $\text{Zn}^{2+}\text{-H}^+$ (e) $\text{Cd}^{2+}\text{-H}^+$ (f) $\text{Hg}^{2+}\text{-H}^+$ (g) $\text{Pb}^{2+}\text{-H}^+$ exchanges using ZrATMP	76
2.13	(a-g) Arrhenius plots $\ln D_A$ versus $1/T$ for (a) $\text{Co}^{2+} - \text{H}^+$ (b) $\text{Ni}^{2+} - \text{H}^+$ (c) $\text{Cu}^{2+} - \text{H}^+$ (d) $\text{Zn}^{2+} - \text{H}^+$ (e) $\text{Cd}^{2+} - \text{H}^+$ (f) $\text{Hg}^{2+} - \text{H}^+$ (g) $\text{Pb}^{2+} - \text{H}^+$ exchanges using ZrATMP	77
2.14	% Uptake of Co^{2+}	80
2.15	% Uptake of Ni^{2+}	80
2.16	% Uptake of Cu^{2+}	81
2.17	% Uptake of Zn^{2+}	81
2.18	% Uptake of Cd^{2+}	82
2.19	% Uptake of Hg^{2+}	82
2.20	% Uptake of Pb^{2+}	83
2.21	(a-g) Langmuir plots for transition metal ions (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} and heavy metal ions (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} metal ions using ZrATMP	85
2.22	(a-g) Freundlich plots for transition metal ions (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} and heavy metal ions (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} metal ions using ZrATMP	86
2.23	Plot of fractional attainment of equilibrium for $\text{Cu}^{2+}\text{-H}^+$ exchange vs. time using TiATMP	88
2.24	Plot of t vs. time for $\text{Cu}^{2+}\text{-H}^+$ exchange for 0.014 M concentration using TiATMP	91
2.25	Plots of t vs. time plots at different temperatures for (a) $\text{Co}^{2+}\text{-H}^+$ (b) $\text{Ni}^{2+}\text{-H}^+$ (c) $\text{Cu}^{2+}\text{-H}^+$ (d) $\text{Zn}^{2+}\text{-H}^+$ (e) $\text{Cd}^{2+}\text{-H}^+$ (f) $\text{Hg}^{2+}\text{-H}^+$ (g) $\text{Pb}^{2+}\text{-H}^+$ exchanges using TiATMP	92
2.26	(a-g) Arrhenius plots $\ln D_A$ versus $1/T$ for (a) $\text{Co}^{2+} - \text{H}^+$; (b) $\text{Ni}^{2+} - \text{H}^+$; (c) $\text{Cu}^{2+} - \text{H}^+$; (d) $\text{Zn}^{2+} - \text{H}^+$; (e) $\text{Cd}^{2+} - \text{H}^+$; (f) $\text{Hg}^{2+} - \text{H}^+$ (g) $\text{Pb}^{2+} - \text{H}^+$ exchanges using TiATMP	93
2.27	% Uptake of Co^{2+}	96
2.28	% Uptake of Ni^{2+}	96
2.29	% Uptake of Cu^{2+}	97
2.30	% Uptake of Zn^{2+}	97
2.31	% Uptake of Cd^{2+}	98
2.32	% Uptake of Hg^{2+}	98
2.33	% Uptake of Pb^{2+}	99
2.34	(a-g) Langmuir Plots for transition metal ions (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} and heavy metal ions (e) Cd^{2+} (f) Hg^{2+} (g) Pb^{2+} metal ions using TiATMP	101

2.35	(a-g) Freundlich plots for transition metal ions (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} and heavy metal ions (e) Cd^{2+} (f) Hg^{2+} (g) Pb^{2+} metal ions using TiATMP	102
2.36	Types of elution curve (a) linear isotherm (b) classical isotherm and (c) anti classical isotherm	108
2.37	Types of elution curve with (a) controlled flow (b) tailing/not good flow and (c) ideal time break.	110
2.38	(a-g) Breakthrough curves for transition (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; and heavy (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} metal ions using ZrATMP	117
2.39	(a-g) Elution behaviour of transition metal ions and heavy metal ions using 0.2 M HNO_3 as eluent (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; (e) Cd^{2+} (f) Hg^{2+} and (g) Pb^{2+} using ZrATMP	118
2.40	(a-g) Binary separations of transition and heavy metal ions using ZrATMP (a) Ni^{2+} - Cu^{2+} (b) Co^{2+} - Cu^{2+} , (c) Zn^{2+} - Ni^{2+} , (d) Co^{2+} - Ni^{2+} (e) Hg^{2+} - Pb^{2+} (f) Hg^{2+} - Cd^{2+} (g) Cd^{2+} - Pb^{2+}	120
2.41	Ternary separations of transition and heavy metal ions using ZrATMP (a) Co^{2+} - Ni^{2+} - Cu^{2+} and (b) Hg^{2+} - Cd^{2+} - Pb^{2+}	121
2.42	Plot of % retention in K_d values versus number of cycles using ZrATMP	121
2.43	(a-g) Breakthrough curves for transition (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; and heavy (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} metal ions using TiATMP	125
2.44	(a-g) Elution behaviour of transition and heavy metal using 0.2 M HNO_3 as eluent (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; (e) Cd^{2+} ; (f) Hg^{2+} and (g) Pb^{2+} using TiATMP	126
2.45	(a-g) Binary separations of transition and heavy metal ions using TiATMP (a) Ni^{2+} - Cu^{2+} (b) Co^{2+} - Cu^{2+} , (c) Zn^{2+} - Cu^{2+} , (d) Ni^{2+} - Zn^{2+} (e) Hg^{2+} - Cd^{2+} (f) Cd^{2+} - Pb^{2+} (g) Hg^{2+} - Pb^{2+}	127
2.46	Ternary separations of transition and heavy metal ions using TiATMP (a) Ni^{2+} - Co^{2+} - Cu^{2+} and (b) Hg^{2+} - Cd^{2+} - Pb^{2+}	129
2.47	Plot of % retention in K_d values versus number of cycles using TiATMP	129

Table No.	Caption to Tables	Pg. No.
2.1	Optimization of reaction conditions for synthesis of ZrATMP	45
2.2	Optimization of reaction conditions for synthesis of TiATMP	45
2.3	Characterization of ZrATMP	54
2.4	Characterization of TiATMP	56

2.5	<i>Thermodynamic parameters evaluated for M^{2+} - H^{+} exchange at various temperatures using ZrATMP</i>	72
2.6	<i>Kinetic parameters evaluated for M^{2+}-H^{+} exchange using ZrATMP</i>	75
2.7	<i>% Uptake of metal ions varying pH using ZrATMP</i>	79
2.8	<i>Equilibrium time determination (varying temperature) using ZrATMP</i>	79
2.9	<i>% Uptake of Co^{2+} using ZrATMP</i>	80
2.10	<i>% Uptake of Ni^{2+} using ZrATMP</i>	80
2.11	<i>% Uptake of Cu^{2+} using ZrATMP</i>	81
2.12	<i>% Uptake of Zn^{2+} using ZrATMP</i>	81
2.13	<i>% Uptake of Cd^{2+} using ZrATMP</i>	82
2.14	<i>% Uptake of Hg^{2+} using ZrATMP</i>	82
2.15	<i>% Uptake of Pb^{2+} using ZrATMP</i>	83
2.16	<i>Langmuir and Fruendlich constants evaluated for transition and heavy metal ions using ZrATMP</i>	84
2.17	<i>Thermodynamic parameters evaluated for M^{2+}-H^{+} exchange at various temperatures using TiATMP</i>	88
2.18	<i>Kinetic parameters evaluated for M^{2+}-H^{+} exchange using TiATMP</i>	91
2.19	<i>% Uptake of metal ions varying pH using TiATMP</i>	95
2.20	<i>Equilibrium time determination (varying temperature) using TiATMP</i>	95
2.21	<i>% Uptake of Co^{2+} using TiATMP</i>	96
2.22	<i>% Uptake of Ni^{2+} using TiATMP</i>	96
2.23	<i>% Uptake of Cu^{2+} using TiATMP</i>	97
2.24	<i>% Uptake of Zn^{2+} using TiATMP</i>	97
2.25	<i>% Uptake of Cd^{2+} using TiATMP</i>	98
2.26	<i>% Uptake of Hg^{2+} using TiATMP</i>	98
2.27	<i>% Uptake of Pb^{2+} using TiATMP</i>	99
2.28	<i>Langmuir and Fruendlich constants evaluated for transition and heavy metal ions using TiATMP</i>	100
2.29	<i>Distribution coefficient (K_d) values ($mL \cdot g^{-1}$) evaluated varying metal ion concentration in aqueous medium using ZrATMP</i>	116
2.30	<i>BTC ($mmol \cdot g^{-1}$) and K_d ($mL \cdot g^{-1}$) in aqueous and various electrolyte media using ZrATMP</i>	116
2.31	<i>Percentage elution (% E) of metal ions in different electrolyte</i>	119

	<i>media using ZrATMP</i>	
2.32	<i>Binary separations of transition and heavy metal ions using ZrATMP</i>	119
2.33	<i>Ternary separations of transition and heavy metal ions using ZrATMP</i>	121
2.34	<i>Distribution coefficient (K_d) values ($\text{mL}\cdot\text{g}^{-1}$) varying metal ion concentration in aqueous medium using TiATMP</i>	124
2.35	<i>BTC ($\text{mmol}\cdot\text{g}^{-1}$) and K_d ($\text{mL}\cdot\text{g}^{-1}$) values evaluated in aqueous and various electrolyte media using TiATMP</i>	124
2.36	<i>Percentage elution (%E) of metal ions in different electrolyte media using TiATMP</i>	127
2.37	<i>Binary separations of transition and heavy metal ions using TiATMP</i>	127
2.38	<i>Ternary separations of transition and heavy metal ions using TiATMP</i>	129

CHAPTER III

Figure No.	Caption to Figures	Pg. No.
3.1	<i>Proposed Structure of ZrD Gel</i>	153
3.2	<i>EDX of ZrD</i>	153
3.3	<i>FTIR spectrum of ZrD and Cu^{2+} chelated ZrD</i>	153
3.4	<i>pH titration curve for ZrD</i>	153
3.5	<i>Proposed Structure of Cu^{2+} chelated ZrD</i>	153
3.6	<i>UV-DRS Spectrum of ZrD and Cu^{2+} chelated ZrD</i>	154
3.7	<i>EDX of Cu^{2+} chelated ZrD</i>	154
3.8	<i>NMR Spectra of ZrD</i>	154
3.9	<i>NMR Spectra of Cu^{2+} chelated ZrD</i>	154
3.10	<i>TGA of ZrD</i>	154
3.11	<i>XRD of ZrD</i>	155
3.12	<i>SEM of ZrD</i>	155
3.13	<i>Plot of fractional attainment of equilibrium for Cu^{2+}-H^+ exchange vs. time using ZrD</i>	155
3.14	<i>(a-g) Elution behaviour of transition and heavy metal ions using 0.2 M HNO_3 as eluent (a) Co^{2+}; (b) Ni^{2+}; (c) Cu^{2+}; (d) Zn^{2+}; (e) Cd^{2+}; (f) Hg^{2+}; (g) Pb^{2+} using ZrD</i>	156
3.15	<i>(a-f) Binary separations of transition and heavy metal ions using</i>	157

ZrD (a) Zn^{2+} - Co^{2+} (b) Zn^{2+} - Cu^{2+} (c) Ni^{2+} - Cu^{2+} (d) Co^{2+} - Cu^{2+} (e) Cd^{2+} - Hg^{2+} (f) Cd^{2+} - Pb^{2+}

3.16	EDX of ZrD treated with CuCl_2	158
3.17	EDX of ZrD treated with HgCl_2	158
3.18	EDX of tap water Sample	158
3.19	EDX of ZrD treated with tap water sample	158
3.20	Plot of % retention in K_d values versus number of cycles using ZrD	158
3.21	Proposed Structure of ZrT Gel	166
3.22	EDX of ZrT	166
3.23	FTIR spectrum of ZrT and Cu^{2+} chelated ZrT	166
3.24	pH titration curve for ZrT	166
3.25	Proposed Structure of Cu^{2+} chelated ZrT	166
3.26	UV-DRS Spectrum of ZrT and Cu^{2+} chelated ZrT	167
3.27	EDX of Cu^{2+} chelated ZrT	167
3.28	NMR Spectra of ZrT	167
3.29	NMR Spectra of Cu^{2+} chelated ZrT	167
3.30	TGA of ZrT	167
3.31	XRD of ZrT	168
3.32	SEM of ZrT	168
3.33	Plot of fractional attainment of equilibrium for Cu^{2+} - H^+ exchange vs. time ZrT	168
3.34	(a-g) Elution behaviour of transition and heavy metal ions using 0.2 M HNO_3 as eluent (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} using ZrT	169
3.35	(a-f) Binary separations of transition and heavy metal ions using ZrT (a) Co^{2+} - Cu^{2+} (b) Co^{2+} - Zn^{2+} (c) Ni^{2+} - Cu^{2+} (d) Zn^{2+} - Cu^{2+} (e) Cd^{2+} - Hg^{2+} (f) Cd^{2+} - Pb^{2+}	170
3.36	EDX of ZrT treated with CuCl_2	171
3.37	EDX of ZrT treated with HgCl_2	171
3.38	EDX of tap water sample	171
3.39	EDX of ZrT treated with tap water sample	171
3.40	Plot of % retention in K_d values versus number of cycles using ZrT	171

Table No.	Caption to Tables	Pg. No.
3.1	Optimization of reaction conditions for synthesis of ZrD	143
3.2	Optimization of reaction conditions for synthesis of ZrT	143

3.3	<i>Physical and ion exchange characteristics of ZrD</i>	159
3.4	<i>Effect of pH on adsorption of metal ions using ZrD</i>	159
3.5	<i>Distribution coefficient (K_d) values ($\text{mL}\cdot\text{g}^{-1}$) evaluated varying metal ion concentration in aqueous medium using ZrD</i>	160
3.6	<i>Distribution coefficient (K_d) ($\text{mL}\cdot\text{g}^{-1}$) evaluated at optimum condition in aqueous and various electrolyte media/concentration using ZrD</i>	160
3.7	<i>Percentage elution (%E) of metal ions in different electrolyte media using ZrD</i>	160
3.8	<i>Binary separations of metal transition and heavy ions using ZrD</i>	161
3.9	<i>Simultaneous uptake of Cu^{2+}, Hg^{2+} and Cl^- using ZrD</i>	161
3.10	<i>Simultaneous removal of cations and anions in tap water samples using ZrD</i>	161
3.11	<i>Physical and ion exchange characteristics of ZrT</i>	172
3.12	<i>Effect of pH on adsorption of metal ions using ZrT</i>	172
3.13	<i>Distribution coefficient (K_d) values ($\text{mL}\cdot\text{g}^{-1}$) evaluated varying metal ion concentration in aqueous medium using ZrT</i>	173
3.14	<i>Distribution coefficient (K_d) ($\text{mL}\cdot\text{g}^{-1}$) evaluated at optimum condition in aqueous and various electrolyte media/concentration using ZrT</i>	173
3.15	<i>Percentage elution (% E) of metal ions in different electrolyte media using ZrT</i>	173
3.16	<i>Binary separations of transition and heavy metal ions using ZrT</i>	174
3.17	<i>Simultaneous uptake of Cu^{2+}, Hg^{2+} and Cl^- using ZrT</i>	174
3.18	<i>Simultaneous removal of cations and anions in tap water samples using ZrT</i>	174

CHAPTER IV

Figure No.	Caption to Figures	Pg. No.
4.1	<i>Experimental set up for impedance measurements. Two Ag/AgCl electrodes are used as CEs, and potential across the membrane (BM) is measured with two REs using Haber– Luggin capillaries salt bridges (HLC). The solution flux (SF) is directed to the membrane to avoid concentration changes at interfacial zone</i>	190
4.2	<i>Schematic diagram of experimental setup used for metal ion transport studies</i>	193

4.3	<i>Schematic experimental set up for the determination of membrane permselectivity by membrane potential measurements</i>	194
4.4	<i>ATR Spectra of (A) ZrT particle (B) Cross linked ZrTETA-55 membrane</i>	206
4.5	<i>¹H-NMR spectrum of ZrTETA-55 membrane</i>	206
4.6	<i>Schematic structure of ZrTETA-55 membrane</i>	206
4.7	<i>SEM images of ZrT gel particles</i>	206
4.8	<i>SEM image of cross section of ZrTETA-55 membrane</i>	206
4.9	<i>SEM image of cross section of ZrTETA-55 membrane</i>	206
4.10	<i>DSC of ZrTETA-X membranes</i>	207
4.11	<i>DMA curves for ZrTETA-X membranes</i>	207
4.12	<i>Adsorption capacity of ZrTETA-55 membrane for different metal ions</i>	207
4.13	<i>Diffuse reflectance spectrum (DRS) for ZrTETA-55 membrane before and after Cu²⁺ chelation</i>	207
4.14	<i>Proposed structure for Cu²⁺ chelated ZrTETA-55 membrane</i>	207
4.15	<i>WXR pattern of ZrTETA-55 and Cu²⁺ Chelated ZrTETA-55 membrane</i>	207
4.16	<i>EDX of ZrTETA-55 membrane</i>	208
4.17	<i>EDX of Cu²⁺ chelated ZrTETA-55 membrane</i>	208
4.18	<i>Membrane conductivity data in equilibration of different metal salt solutions (0.01 M)</i>	208
4.19	<i>i-v characteristics for ZrTETA-X membranes in equilibration with 0.10M NaCl solution</i>	208
4.20	<i>t_i^m values in equilibration with 0.1M metal ion solutions across ZrTETA-X membranes</i>	208
4.21	<i>Variation of electro osmotic flux (J_v) with applied voltage for different ZrTETA-X membranes</i>	208
4.22	<i>Electro-transport flux for different metal ions (J) (0.10 M) across ZrTETA-X membranes at 10 mA/cm² applied current density</i>	209
4.23	<i>Ionic flux data across different ZrTETA-X membranes for various metal ions (0.10M) under 10 mA/cm² applied current density</i>	209
4.24	<i>Separation factor (SF) of different equi-molar metal ion mixtures (0.10 M) for ZrTETA-X membranes under 10 mA/cm² applied current density</i>	209

4.25	<i>Effect of pH on Cu²⁺ adsorption towards crosslinked ZrTETA-55 hybrid membrane</i>	215
4.26	<i>Effect of contact time on the adsorption of Cu²⁺ at different concentrations using ZrTETA-55 hybrid membrane</i>	215
4.27	<i>Effect of Cu²⁺ concentration on adsorption (%) using crosslinked ZrTETA-55 hybrid membrane at pH 7.0</i>	215
4.28	<i>Effect of adsorbent dose for Cu²⁺ (50mg.L⁻¹) adsorption using crosslinked ZrTETA-55 hybrid membrane</i>	215
4.29	<i>Vant Hoff plot [log K_c vs. 1/T]</i>	215
4.30	<i>Pseudo first order plots for Cu²⁺ adsorption on crosslinked ZrTETA-55 hybrid membrane at different concentration</i>	215
4.31	<i>Pseudo second order plot for Cu²⁺ adsorption on crosslinked ZrTETA-55 hybrid membrane at different concentration (pH 7.0)</i>	216
4.32	<i>Freundlich plots for Cu²⁺ adsorption on crosslinked ZrTETA-55 hybrid membrane under different experimental conditions</i>	216
4.33	<i>Langmuir plot for Cu²⁺ adsorption on crosslinked ZrTETA-55 hybrid membrane under different experimental conditions</i>	216

Table No.	Caption to Tables	Pg. No.
4.1	<i>Properties of commercial ion exchange membranes [1-7]</i>	180
4.2	<i>ϕ_w, Φ_{H_2O}, IUC, $r(\text{\AA})$, X_m, and λ_w values for different ZrTETA-X membranes</i>	210
4.3	<i>Oxidative and Hydrolytic Stabilities for ZrTETA-X Chelating Membranes</i>	210
4.4	<i>Characteristic values of i-v curves for ZrTETA-X membranes in equilibration with 0.10 M NaCl solutions</i>	210
4.5	<i>Thermodynamic parameters and correlation coefficients (R^2) at different temperatures for the adsorption of Cu²⁺ on crosslinked ZrTETA-55 hybrid membrane at pH 7.0</i>	217
4.6	<i>Pseudo-first- and second-order kinetic constants and correlation coefficients (R^2) for Cu²⁺ adsorption at pH 7.0 using ZrTETA-55 hybrid membrane</i>	217
4.7	<i>Freundlich and Langmuir constants and correlation coefficients (R^2) for Cu²⁺ adsorption on ZrTETA-55 hybrid membrane at pH 7.0</i>	217
4.8	<i>Langmuir correlation coefficients (R_L) for Cu²⁺ adsorption at different concentration on ZrTETA-55 hybrid membrane at pH 7.0</i>	217

Scheme No.	Caption to Schemes	Pg. No.
4.1	<i>Synthesis of ZrTETA-X cross linked membrane</i>	186

