

CHAPTER II

Figure No.	Caption to Figures	Pg. No.
2.1	Schematic representation of the arrangement of layers in Zirconium phosphonates [22]	33
2.2	Structure of Zirconium phosphonates [23]	33
2.3	Scheme of a layered diphosphonate in which both phosphonic groups are bonded to the same face of the inorganic layer [6]	35
2.4	(a) structure of ATMP (b) structure of phosphoric acid	40
2.5	pH titration curves for (A) ZrATMP and (B) TiATMP	48
2.6	(a-d) Concentration and eluent volume optimization for determination of CEC using ZrATMP and TiATMP	50
2.7	(a) EDX (b) FTIR (c) TGA (d) XRD and (e) SEM of ZrATMP	55
2.8	(a) EDX (b) FTIR (c) TGA (d) XRD and (e) SEM of TiATMP	57
2.9	Ion exchange equilibrium [63]	58
2.10	Plot of the fractional attainment of equilibrium for Cu^{2+} - H^{+} exchange vs. time using ZrATMP	72
2.11	Plots of τ vs. time for Cu^{2+} - H^{+} exchange for 0.014 M concentration using ZrATMP	75
2.12	Plots of τ vs. time at different temperatures for (a) Co^{2+} - H^{+} (b) Ni^{2+} - H^{+} (c) Cu^{2+} - H^{+} (d) Zn^{2+} - H^{+} (e) Cd^{2+} - H^{+} (f) Hg^{2+} - H^{+} (g) Pb^{2+} - H^{+} exchanges using ZrATMP	76
2.13	(a-g) Arrhenius plots $\ln D_A$ versus $1/T$ for (a) Co^{2+} - H^{+} (b) Ni^{2+} - H^{+} (c) Cu^{2+} - H^{+} (d) Zn^{2+} - H^{+} (e) Cd^{2+} - H^{+} (f) Hg^{2+} - H^{+} (g) Pb^{2+} - H^{+} exchanges using ZrATMP	77
2.14	% Uptake of Co^{2+} using ZrATMP	80
2.15	% Uptake of Ni^{2+} using ZrATMP	80
2.16	% Uptake of Cu^{2+} using ZrATMP	81
2.17	% Uptake of Zn^{2+} using ZrATMP	81
2.18	% Uptake of Cd^{2+} using ZrATMP	82
2.19	% Uptake of Hg^{2+} using ZrATMP	82
2.20	% Uptake of Pb^{2+} using ZrATMP	83
2.21	(a-g) Langmuir plots for transition metal ions (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} and heavy metal ions (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} metal ions using ZrATMP	85
2.22	(a-g) Freundlich plots for transition metal ions (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} and heavy metal ions (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} metal ions using ZrATMP	86
2.23	Plot of the fractional attainment of equilibrium for Cu^{2+} - H^{+} exchange vs. time using TiATMP	88
2.24	Plots of τ vs. time for Cu^{2+} - H^{+} exchange for 0.014 M concentration using TiATMP	91

2.25	Plots of τ vs. time plots at different temperatures for (a) Co^{2+} - H^+ (b) Ni^{2+} - H^+ (c) Cu^{2+} - H^+ (d) Zn^{2+} - H^+ (e) Cd^{2+} - H^+ (f) Hg^{2+} - H^+ (g) Pb^{2+} - H^+ exchanges using TiATMP	92
2.26	(a-g) Arrhenius plots $\ln D_A$ versus $1/T$ for (a) $\text{Co}^{2+} - \text{H}^+$; (b) $\text{Ni}^{2+} - \text{H}^+$; (c) $\text{Cu}^{2+} - \text{H}^+$; (d) $\text{Zn}^{2+} - \text{H}^+$; (e) $\text{Cd}^{2+} - \text{H}^+$; (f) $\text{Hg}^{2+} - \text{H}^+$ (g) $\text{Pb}^{2+} - \text{H}^+$ exchanges using TiATMP	93
2.27	% Uptake of Co^{2+} using TiATMP	96
2.28	% Uptake of Ni^{2+} using TiATMP	96
2.29	% Uptake of Cu^{2+} using TiATMP	97
2.30	% Uptake of Zn^{2+} using TiATMP	97
2.31	% Uptake of Cd^{2+} using TiATMP	98
2.32	% Uptake of Hg^{2+} using TiATMP	98
2.33	% Uptake of Pb^{2+} using TiATMP	99
2.34	(a-g) Langmuir Plots for transition metal ions (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} and heavy metal ions (e) Cd^{2+} (f) Hg^{2+} (g) Pb^{2+} metal ions using TiATMP	101
2.35	(a-g) Freundlich plots for transition metal ions (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} and heavy metal ions (e) Cd^{2+} (f) Hg^{2+} (g) Pb^{2+} metal ions using TiATMP	102
2.36	Types of elution curve (a) linear isotherm (b) classical isotherm and (c) anti classical isotherm	108
2.37	Types of elution curve with (a) controlled flow (b) tailing/not good flow and (c) ideal time break.	110
2.38	(a-g) Breakthrough curves for transition (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; and heavy (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} metal ions using ZrATMP	117
2.39	(a-g) <i>Elution behaviour of transition metal ions and heavy metal ions using 0.2 M HNO_3 as eluent</i> (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; (e) Cd^{2+} (f) Hg^{2+} and (g) Pb^{2+} using ZrATMP	118
2.40	(a-g) Binary separations of transition and heavy metal ions using ZrATMP (a) Ni^{2+} - Cu^{2+} (b) Co^{2+} - Cu^{2+} , (c) Zn^{2+} - Ni^{2+} , (d) Co^{2+} - Ni^{2+} (e) Hg^{2+} - Pb^{2+} (f) Hg^{2+} - Cd^{2+} (g) Cd^{2+} - Pb^{2+}	120
2.41	Ternary separations of transition and heavy metal ions using ZrATMP (a) Co^{2+} - Ni^{2+} - Cu^{2+} and (b) Hg^{2+} - Cd^{2+} - Pb^{2+}	121
2.42	A plot of % Retention in K_d values versus number of cycles using ZrATMP	121
2.43	(a-g) Breakthrough curves for transition (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; and heavy (e) Cd^{2+} ; (f) Hg^{2+} ; (g) Pb^{2+} metal ions using TiATMP	125
2.44	(a-g) Elution behaviour of transition and heavy metal using 0.2 M HNO_3 as eluent (a) Co^{2+} ; (b) Ni^{2+} ; (c) Cu^{2+} ; (d) Zn^{2+} ; (e) Cd^{2+} ; (f) Hg^{2+} and (g) Pb^{2+} using TiATMP	126

2.45	(a-g) Binary separations of transition and heavy metal ions using TiATMP (a) Ni^{2+} - Cu^{2+} (b) Co^{2+} - Cu^{2+} , (c) Zn^{2+} - Cu^{2+} , (d) Ni^{2+} - Zn^{2+} (e) Hg^{2+} - Cd^{2+} (f) Cd^{2+} - Pb^{2+} (g) Hg^{2+} - Pb^{2+}	127
2.46	Ternary separations of transition and heavy metal ions using TiATMP (a) Ni^{2+} - Co^{2+} - Cu^{2+} and (b) Hg^{2+} - Cd^{2+} - Pb^{2+}	129
2.47	A plot of % Retention in K_d values versus number of cycles using TiATMP	129

Table No.	Caption to Tables	Pg. No.
2.1	Optimization of reaction conditions for synthesis of ZrATMP	45
2.2	Optimization of reaction conditions for synthesis of TiATMP	45
2.3	Characterization of ZrATMP	54
2.4	Characterization of TiATMP	56
2.5	Thermodynamic parameters evaluated for $M^{2+} - H^{+}$ exchange at various temperatures using ZrATMP	72
2.6	Kinetic parameters evaluated for $M^{2+}-H^{+}$ exchange using ZrATMP	75
2.7	% Uptake of metal ions varying pH using ZrATMP	79
2.8	Equilibrium time determination (varying temperature) using ZrATMP	79
2.9	% Uptake of Co^{2+} using ZrATMP	80
2.10	% Uptake of Ni^{2+} using ZrATMP	80
2.11	% Uptake of Cu^{2+} using ZrATMP	81
2.12	% Uptake of Zn^{2+} using ZrATMP	81
2.13	% Uptake of Cd^{2+} using ZrATMP	82
2.14	% Uptake of Hg^{2+} using ZrATMP	82
2.15	% Uptake of Pb^{2+} using ZrATMP	83
2.16	Langmuir and Freundlich constants evaluated for transition and heavy metal ions using ZrATMP	84
2.17	Thermodynamic parameters evaluated for $M^{2+}-H^{+}$ exchange at various temperatures using TiATMP	88
2.18	Kinetic parameters evaluated for $M^{2+}-H^{+}$ exchange using TiATMP	91
2.19	% Uptake of metal ions varying pH using TiATMP	95
2.20	Equilibrium time determination (varying temperature) using TiATMP	95
2.21	% Uptake of Co^{2+} using TiATMP	96
2.22	% Uptake of Ni^{2+} using TiATMP	96
2.23	% Uptake of Cu^{2+} using TiATMP	97
2.24	% Uptake of Zn^{2+} using TiATMP	97
2.25	% Uptake of Cd^{2+} using TiATMP	98
2.26	% Uptake of Hg^{2+} using TiATMP	98
2.27	% Uptake of Pb^{2+} using TiATMP	99
2.28	Langmuir and Freundlich constants evaluated for transition and	100

	heavy metal ions using TiATMP	
2.29	Distribution coefficient (K_d) values ($\text{mL}\cdot\text{g}^{-1}$) evaluated varying metal ion concentration in aqueous media using ZrATMP	116
2.30	BTC ($\text{mmol}\cdot\text{g}^{-1}$) and K_d ($\text{mL}\cdot\text{g}^{-1}$) in aqueous and various electrolyte media using ZrATMP	116
2.31	Percentage Elution (% E) of metal ions in different electrolyte media using ZrATMP	119
2.32	Binary separations of transition and heavy metal ions using ZrATMP	119
2.33	Ternary separations of transition and heavy metal ions using ZrATMP	121
2.34	Distribution coefficient (K_d) values ($\text{mL}\cdot\text{g}^{-1}$) varying metal ion concentration in aqueous media using TiATMP	124
2.35	BTC ($\text{mmol}\cdot\text{g}^{-1}$) and K_d ($\text{mL}\cdot\text{g}^{-1}$) values evaluated in aqueous and various electrolyte media using TiATMP	124
2.36	Percentage elution (%E) of metal ions in different electrolyte media using TiATMP	127
2.37	Binary separations of transition and heavy metal ions using TiATMP	127
2.38	Ternary separations of transition and heavy metal ions using TiATMP	129
