

APPENDIX

CHAPTER I

Figure No.	Caption to Figures	Pg. No.
1.1	<i>Parallel and sequential reactions.</i>	10
1.2	<i>Structures of zirconium phosphate.</i>	22
1.3	<i>Idealized portion of ZP layer.</i>	22
1.4	<i>Schematic diagram of the arrangement of two adjacent macroanions in -ZP.</i>	23
1.5	<i>(a) $-Ti(HPO_4)_2 \cdot H_2O$ octahedra represent TiO_6, dark grey tetrahedra represent HPO_4 and spheres represent water molecules [$d_{002}=7.6\text{\AA}$].</i>	24
	<i>(b) $-r(PO_4)(H_2PO_4)_2 \cdot 2H_2O$ octahedra represent ZrO_6, dark grey tetrahedra represent PO_4 and light grey tetrahedra represent H_2PO_4 and spheres represent water molecules [$d_{002}=12.2\text{\AA}$].</i>	24
Table No.	Caption to Tables	Pg. No.
1.1	<i>History of Industrial catalytic processes.</i>	04
1.2	<i>E-factor for various segments of chemical industry.</i>	11
1.3	<i>ASTM procedures for catalyst characterization.</i>	14

CHAPTER II

Figure No.	Caption to Figures	Pg. No.
2.1	<i>(a) Highly branched networks of silicon.</i>	42
	<i>(b) Condensation reactions leading to closed ring 3D structure.</i>	
	<i>(c) SEM showing Ostwald ripening mechanism.</i>	
	<i>(d) Polymerization behavior of aqueous silica; A= in presence of salts/acidic medium, B= alkaline medium.</i>	
2.2	<i>Sequential steps involved in sol-gel process.</i>	44
2.3	<i>(a) FTIR spectra of ZrPW.</i>	57
	<i>(b) TGA plot of ZrPW.</i>	57
	<i>(c) XRD pattern of ZrPW.</i>	57
	<i>(d) SEM image of ZrPW.</i>	58
	<i>(e) NH_3-TPD patterns of ZrPW.</i>	58
2.4	<i>(a) FTIR spectra of TiPW.</i>	60
	<i>(b) TGA plot of TiPW.</i>	60
	<i>(c) XRD pattern of TiPW.</i>	60
	<i>(d) SEM image of TiPW.</i>	61
	<i>(e) NH_3-TPD patterns of TiPW.</i>	61

2.5	(a) FTIR spectra of SnPW.	63
	(b) TGA plot of SnPW.	63
	(c) XRD pattern of SnPW.	63
	(d) SEM image of SnPW.	64
	(e) NH ₃ -TPD patterns of SnPW.	64
2.6	(a) FTIR spectra of 12-TPA/ZrO ₂ -20.	66
	(b) TGA plot of 12-TPA/ZrO ₂ -20.	66
	(c) XRD pattern of 12-TPA.	66
	(d) XRD pattern of 12-TPA/ZrO ₂ -20.	67
	(e) SEM image of ZrO ₂ .	67
	(f) SEM image of 12-TPA/ZrO ₂ -20.	67
	(g) NH ₃ -TPD patterns of 12-TPA/ZrO ₂ -20.	68
2.7	(a) FTIR spectra of 12-TPA/TiO ₂ -20.	70
	(b) TGA plot of 12-TPA/TiO ₂ -20.	70
	(c) XRD pattern of 12-TPA.	70
	(d) XRD pattern of 12-TPA/ZrO ₂ -20.	71
	(e) SEM image of ZrO ₂ .	71
	(f) SEM image of 12-TPA/ZrO ₂ -20.	71
	(g) NH ₃ -TPD patterns of 12-TPA/ZrO ₂ -20.	72
2.8	(a) FTIR spectra of 12-TPA/SnO ₂ -20.	74
	(b) TGA plot of 12-TPA/SnO ₂ -20.	74
	(c) XRD pattern of 12-TPA.	74
	(d) XRD pattern of 12-TPA/SnO ₂ -20.	75
	(e) SEM image of SnO ₂ .	75
	(f) SEM image of 12-TPA/SnO ₂ -20.	75
	(g) NH ₃ -TPD patterns of 12-TPA/SnO ₂ -20.	76
Table No.	Caption to Tables	Pg. No.
2.1	Optimization of parameters for synthesis of ZrPW.	48
2.2	Optimization of parameters for synthesis of TiPW.	49
3.3	Optimization of parameters for synthesis of SnPW.	49
2.4	Optimization of loading of 12-TPA (wt. %) onto M(IV)O ₂ .	50
2.5	Characterization of ZrPW.	56
2.6	Characterization of TiPW.	59
2.7	Characterization of SnPW.	62
2.8	Characterization of 12-TPA/ZrO ₂ -20.	65
2.9	Characterization of 12-TPA/TiO ₂ -20.	69
2.10	Characterization of 12-TPA/SnO ₂ -20.	73

CHAPTER III

Figure No.	Caption to Figures	Pg. No.
3.1	<i>Possible protonated intermediates in esterification reaction.</i>	87
3.2	<i>Optimization of reaction time for synthesis of EA.</i>	90
3.3	<i>Optimization of amount of catalyst for synthesis of EA.</i>	90
3.4	<i>Optimization of mole ratio of reactants for synthesis of EA.</i>	90
3.5	<i>Comparison of % yields of EA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	91
3.6	<i>Comparison of % yields of PA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	92
3.7	<i>Comparison of % yields of BA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	93
3.8	<i>Comparison of % yields of BzA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	94
3.9	<i>EDX of fresh ZrPW.</i>	95
3.10	<i>EDX of spent ZrPW (in synthesis of EA).</i>	95
3.11	<i>EDX of fresh 12-TPA/ZrO₂-20.</i>	95
3.12	<i>EDX of spent 12-TPA/ZrO₂-20 (in synthesis of EA).</i>	95
3.13	<i>Optimization of reaction time for synthesis of DEM.</i>	105
3.14	<i>Optimization of amount of catalyst for synthesis of DEM.</i>	105
3.15	<i>Optimization of mole ratio of reactants for synthesis of DEM.</i>	105
3.16	<i>Comparison of % yields of DEM at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	106
3.17	<i>Comparison of % yields of DES at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	107
3.18	<i>Comparison of % yields of DOP at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	108
3.19	<i>Comparison of % yields of DBP at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	109
3.20	<i>EDX of fresh ZrPW.</i>	110
3.21	<i>EDX of spent ZrPW (in synthesis of DEM).</i>	110
3.22	<i>EDX of fresh 12-TPA/ZrO₂-20.</i>	110
3.23	<i>EDX of spent 12-TPA/ZrO₂-20 (in synthesis of DEM).</i>	110
Table No.	Caption to Tables	Pg. No.
3.1	<i>Optimization of reaction conditions for synthesis of EA using ZrPW and 12-TPA/ZrO₂-20.</i>	89
3.2	<i>Synthesis of mono ester - ethyl acetate (EA) at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	91
3.3	<i>Synthesis of mono ester - propyl acetate (PA) at optimized</i>	92

	<i>condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	
3.4	<i>Synthesis of mono ester - butyl acetate (BA) at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	93
3.5	<i>Synthesis of mono ester - benzyl acetate (BzA) at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	94
3.6	<i>Elemental analysis by EDX for both fresh and spent ZrPW and 12-TPA/ZrO₂-20 in the synthesis of EA.</i>	95
3.7	<i>Optimization of reaction conditions for synthesis of DEM using ZrPW and 12-TPA/ZrO₂-20.</i>	104
3.8	<i>Synthesis of diester – diethyl malonate (DEM) at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	106
3.9	<i>Synthesis of diester – diethyl succinate (DES) at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	107
3.10	<i>Synthesis of diester – dioctyl phthalate (DOP) at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	108
3.11	<i>Synthesis of diester – dibutyl phthalate (DBP) at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	109
3.12	<i>Elemental analysis by EDX for both fresh and spent ZrPW and 12-TPA/ZrO₂-20 in the synthesis of DEM.</i>	110

Scheme No.	Caption to Schemes	Pg. No.
3.1	<i>Esterification reaction.</i>	78
3.2	<i>Synthesis of monoesters (EA, PA, BA and BzA).</i>	85
3.3	<i>Proposed mechanism for esterification of ethanol with acetic acid using M(IV)PWs and 12-TPA/M(IV)O₂.</i>	88
3.4	<i>Schematic presentation of diester formation.</i>	96
3.5	<i>Synthesis of DEM and DES.</i>	100
3.6	<i>Synthesis of DBP and DOP.</i>	100

CHAPTER IV

Figure No.	Caption to Figures	Page No.
4.1	<i>Benzo pyrone structures.</i>	118
4.2	<i>Optimization of reaction time for synthesis of 7H4MC.</i>	133
4.3	<i>Optimization of reaction temperature for synthesis of 7H4MC.</i>	133
4.4	<i>Optimization of amount of catalyst for synthesis of 7H4MC.</i>	133
4.5	<i>Optimization of reaction time under MW irradiation for synthesis of 7H4MC.</i>	133
4.6	<i>Optimization of mole ratio of reactants for synthesis of 7H4MC.</i>	133
4.7	<i>Comparison of % yields of 7H4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under conventional heating.</i>	134
4.8	<i>Comparison of % yields of 7H4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under MW heating.</i>	135
4.9	<i>Comparison of % yields of 7,8DH4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under conventional heating.</i>	136

4.10	<i>Comparison of % yields of 7,8DH4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under MW heating.</i>	137
4.11	<i>Comparison of % yields of 5,7DH4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under conventional heating.</i>	138
4.12	<i>Comparison of % yields of 5,7DH4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under MW heating.</i>	139
4.13	<i>Comparison of % yields of 6H4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under conventional heating.</i>	140
4.14	<i>Comparison of % yields of 6H4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under MW heating.</i>	141
4.15	<i>Comparison of % yields of 6N4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under conventional heating.</i>	142
4.16	<i>Comparison of % yields of 6N4MC at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20 under MW heating.</i>	143
4.17	<i>EDX of fresh ZrPW.</i>	144
4.18	<i>EDX of spent ZrPW for synthesis of 7H4MC.</i>	144
4.19	<i>EDX of fresh 12-TPA/ZrO₂-20.</i>	144
4.20	<i>EDX of spent 12-TPA/ZrO₂-20 for synthesis of 7H4MC.</i>	144
Table No.	Caption to Tables	Page No.
4.1	<i>Optimization of reaction conditions for Pechmann condensation of resorcinol (R) with methyl acetoacetate (MA) using ZrPW and 12-TPA/ZrO₂-20 under conventional and microwave (MW) heating.</i>	132
4.2	<i>Pechmann condensation of resorcinol (R) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under conventional heating.</i>	134
4.3	<i>Pechmann condensation of resorcinol (R) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under microwave (MW) heating.</i>	135
4.4	<i>Pechmann condensation of pyrogallol (Py) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under conventional heating.</i>	136
4.5	<i>Pechmann condensation of pyrogallol (Py) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under microwave (MW) heating.</i>	137
4.6	<i>Pechmann condensation of phloroglucinol (Ph) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under conventional heating.</i>	138
4.7	<i>Pechmann condensation of phloroglucinol (Ph) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under microwave (MW) heating.</i>	139
4.8	<i>Pechmann condensation of hydroquinone (Hq) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under conventional heating.</i>	140
4.9	<i>Pechmann condensation of hydroquinone (Hq) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under microwave (MW) heating.</i>	141

4.10	<i>Pechmann condensation of p-nitrophenol (pNp) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under conventional heating.</i>	142
4.11	<i>Pechmann condensation of p-Nitrophenol (pNp) with methyl acetoacetate (MA) using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimum condition under microwave (MW) heating.</i>	143
4.12	<i>Elemental analysis by EDX for both fresh and spent ZrPW and 12-TPA/ZrO₂-20 in the synthesis of 7H4MC.</i>	144
4.13	<i>Characterization of coumarin derivatives.</i>	145

Caption to Schemes

Scheme No.	Caption	Page No.
4.1	<i>Perkin reaction.</i>	119
4.2	<i>Knoevenagel reaction.</i>	119
4.3	<i>Reaction pathways for the synthesis of coumarin derivatives by Knoevenagel condensation.</i>	120
4.4	<i>Reformatsky Reaction.</i>	121
4.5	<i>Synthesis of 7-hydroxy-4-methyl coumarin.</i>	127
4.6	<i>Synthesis of 7,8-dihydroxy-4-methyl coumarin.</i>	127
4.7	<i>Synthesis of 5,7-dihydroxy-4-methyl coumarin.</i>	127
4.8	<i>Synthesis of 6-hydroxy-4-methyl coumarin.</i>	127
4.9	<i>Synthesis of 6-nitro-4-methyl coumarin.</i>	127
4.10	<i>Mechanism for Pechmann condensation.</i>	130
4.11	<i>Proposed mechanism for the Pechmann condensation of phenols and α-keto ester using M(IV)PWs and 12-TPA/M(IV)O₂.</i>	131

CHAPTER V

Figure No.	Caption to Figures	Pg. No.
5.1	<i>Optimization of reaction time for synthesis of p-BT.</i>	165
5.2	<i>Optimization of reaction temperature for synthesis of p-BT.</i>	165
5.3	<i>Optimization of catalyst amount for synthesis of p-BT.</i>	165
5.4	<i>Optimization of mole ratio of reactants for synthesis of p-BT.</i>	165
5.5	<i>Comparison of % yields of p-BT at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	166
5.6	<i>Comparison of % yields of 4-MA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	167
5.7	<i>Comparison of % yields of 3,4-DMA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	168
5.8	<i>EDX of fresh ZrPW.</i>	169
5.9	<i>EDX of spent ZrPW in synthesis of p-BT.</i>	169
5.10	<i>EDX of fresh 12-TPA/ZrO₂-20.</i>	169
5.11	<i>EDX of spent 12-TPA/ZrO₂-20 in synthesis of p-BT.</i>	169
5.12	<i>Optimization of reaction time for synthesis of 2-MA.</i>	178
5.13	<i>Optimization of reaction temperature for synthesis of 2-MA.</i>	178

5.14	<i>Optimization of catalyst amount for synthesis of 2-MA.</i>	178
5.15	<i>Optimization of mole ratio of reactants for synthesis of 2-MA.</i>	178
5.16	<i>Comparison of % yields of 2-MA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	179
5.17	<i>Comparison of % yields of 2-NA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	180
5.18	<i>Comparison of % yields of 1,3-DHA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	181
5.19	<i>Comparison of % yields of 1,4-DHA at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	182
5.20	<i>EDX of fresh ZrPW.</i>	183
5.21	<i>EDX of spent ZrPW in synthesis of 2-MA.</i>	183
5.22	<i>EDX of fresh 12-TPA/ZrO₂-20.</i>	183
5.23	<i>EDX of spent 12-TPA/ZrO₂-20 in synthesis of 2-MA.</i>	183
Table No.	Caption to Tables	Pg. No.
5.1	<i>Optimization of reaction conditions for alkylation of toluene with benzyl chloride using ZrPW and 12-TPA/ZrO₂-20.</i>	164
5.2	<i>Friedel-Crafts alkylation of toluene with benzyl chloride at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	166
5.3	<i>Friedel-Crafts acylation of anisole with acetyl chloride at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	167
5.4	<i>Friedel-Crafts acylation of veratrole with acetyl chloride at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	168
5.5	<i>Elemental analysis by EDX for both fresh and spent ZrPW and 12-TPA/ZrO₂-20 in alkylation of toluene with benzyl chloride at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	169
5.6	<i>Optimization of reaction conditions for preparation of anthraquinone derivative from toluene and phthalic anhydride by using ZrPW and 12-TPA/ZrO₂-20.</i>	177
5.7	<i>Synthesis of anthraquinone derivative from toluene and phthalic anhydride by using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimized condition.</i>	179
5.8	<i>Synthesis of anthraquinone derivative from nitrobenzene and phthalic anhydride by using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimized condition.</i>	180
5.9	<i>Synthesis of anthraquinone derivative from resorcinol and phthalic anhydride by using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimized condition.</i>	181
5.10	<i>Synthesis of anthraquinone derivative from hydroquinone and phthalic anhydride by using M(IV)PWs and 12-TPA/M(IV)O₂-20 at optimized condition.</i>	182

5.11	<i>Elemental analysis by EDX for both fresh and spent ZrPW and 12-TPA/ZrO₂-20 in the synthesis of 2-MA.</i>	183
5.12	<i>Characterization of anthraquinone derivatives.</i>	184
Scheme No.	Caption to Schemes	Pg. No.
5.1	<i>Reaction scheme for Friedel Crafts alkylation.</i>	150
5.2	<i>Reaction scheme for t-butylation of 1,4-dimethoxybenzene.</i>	150
5.3	<i>Reaction scheme for Friedel Crafts alkylation using bromonium ion.</i>	150
5.4	<i>Reaction scheme for Friedel Crafts acylation.</i>	152
5.5	<i>Friedel-Crafts alkylation of toluene with benzyl chloride.</i>	159
5.6	<i>Friedel-Crafts acylation of anisole with acetyl chloride.</i>	159
5.7	<i>Friedel-Crafts acylation of veratrole with acetyl chloride.</i>	159
5.8	<i>Reaction mechanism for Friedel Crafts alkylation.</i>	161
5.9	<i>Reaction mechanism for Friedel Crafts acylation.</i>	162
5.10	<i>Proposed mechanism of Friedel Crafts alkylation of toluene using M(IV)PWs and 12-TPA/M(IV)O₂.</i>	162
5.11	<i>Proposed mechanism of Friedel Crafts acylation of anisole using M(IV)PWs and 12-TPA/M(IV)O₂.</i>	163
5.12	<i>Friedel–Crafts reaction between phthalic anhydride and substituted benzenes.</i>	170
5.13	<i>Friedel-Crafts reaction of toluene with phthalic anhydride to prepare 2-methyl anthraquinone (2-MA).</i>	173
5.14	<i>Friedel-Crafts reaction of nitrobenzene with phthalic anhydride to prepare 2-nitro anthraquinone (2-NA).</i>	173
5.15	<i>Friedel-Crafts reaction of resorcinol with phthalic anhydride to prepare 1,3-dihydroxy anthraquinone (1,3-DHA).</i>	173
5.16	<i>Friedel-Crafts reaction of hydroquinone with phthalic anhydride to prepare 1,4-dihydroxy anthraquinone(1,4-DHA).</i>	174
5.17	<i>Proposed mechanism for synthesis of 2-methyl anthraquinone using solid acid catalyst.</i>	176

CHAPTER VI

Figure No.	Caption to Figures	Pg. No.
6.1	<i>Optimization of reaction time for synthesis of diacetals.</i>	203
6.2	<i>Optimization of catalyst amount for synthesis of diacetals.</i>	203
6.3	<i>Optimization of mole ratio of reactants for synthesis of diacetals.</i>	203
6.4	<i>Comparison of % yields of diacetal derivative of benzaldehyde at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	204
6.5	<i>Comparison of % yields of diacetal derivative of</i>	205

	<i>cyclohexanone at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	
6.6	<i>Comparison of % yields of diacetal derivative of acetophenone at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	206
6.7	<i>Comparison of % yields of diacetal derivative of benzophenone at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	207
6.8	<i>EDX of fresh ZrPW.</i>	208
6.9	<i>EDX of spent ZrPW in the synthesis of benzaldehyde diacetals.</i>	208
6.10	<i>EDX of fresh 12-TPA/ZrO₂-20.</i>	208
6.11	<i>EDX of spent 12-TPA/ZrO₂-20 in the synthesis of benzaldehyde diacetal.</i>	208
Table No.	Caption to Tables	Pg. No.
6.1	<i>Optimization of reaction conditions for preparation of diacetal from benzaldehyde and PET using ZrPW and 12-TPA/ZrO₂-20.</i>	200
6.2	<i>Optimization of reaction conditions for preparation of diacetal from cyclohexanone and PET using ZrPW and 12-TPA/ZrO₂-20.</i>	201
6.3	<i>Optimization of reaction conditions for preparation of diacetal from acetophenone and PET using ZrPW and 12-TPA/ZrO₂-20.</i>	202
6.4	<i>Synthesis of diacetal derivative of benzaldehyde at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	204
6.5	<i>Synthesis of diacetal derivative of cyclohexanone at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	205
6.6	<i>Synthesis of diacetal derivative of acetophenone at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	206
6.7	<i>Synthesis of diacetal derivative of benzophenone at optimized condition using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	207
6.8	<i>Elemental analysis by EDX for fresh and spent ZrPW and 12-TPA/ZrO₂-20 in the synthesis of benzaldehyde diacetal.</i>	208
6.9	<i>Characterization of diacetal derivatives.</i>	209
Scheme No.	Caption to Schemes	Pg. No.
6.1	<i>Acetalization of carbonyl compound.</i>	190
6.2	<i>General mechanism for acetalization of carbonyl compounds.</i>	195
6.3	<i>Reaction of benzaldehyde with PET to give diacetal of benzaldehyde.</i>	195
6.4	<i>Reaction of cyclohexanone with PET to give diacetal of cyclohexanone.</i>	195
6.5	<i>Reaction of acetophenone with PET to give diacetal of acetophenone.</i>	195
6.6	<i>Reaction of benzophenone with PET to give diacetal of benzophenone.</i>	195

6.7	<i>Proposed mechanism for acetalization of cyclohexanone with pentaerythritol using M(IV)PWs and 12-TPA/M(IV)O₂-20.</i>	199
-----	--	-----

CHAPTER VII

Table No.	Caption to Tables	Pg. No.
7.1	<i>Comparison of % yields of mono ester – EA using TBMA salt, TMBA salts and 12-TPA/M(IV)O₂-20.</i>	217
7.2	<i>Comparison of % yields of diester – DOP using TMA salts, TBMA salt, TMBA salts and 12-TPA/M(IV)O₂-20.</i>	218
7.3	<i>Comparison of % yield of 7-hydroxy-4-methyl coumarin using TMA salts, TMBA salts and 12-TPA/M(IV)O₂-20.</i>	218
7.4	<i>Comparison of % yield of p-benzyl toluene using TMA salts, TMBA salts [M(IV)PWs] and 12-TPA/M(IV)O₂-20.</i>	219
7.5	<i>Comparison of % yield of anthraquinone derivatives using M(IV)PWs, 12-TPA/M(IV)O₂-20 and Montmorillonite KSF.</i>	220
7.6	<i>Comparison of % yield of diacetal derivative of benzaldehyde using TMA salts, TMBA salts [M(IV)PWs] and 12-TPA/M(IV)O₂-20.</i>	220
7.7	<i>% yields of the products formed using TiPW at optimized condition.</i>	221