

List of Figures

2.1	Parent and daughter Regge trajectories (J, M^2) for charmonia (left), bottomonia (middle) and B_c (right) mesons with natural parity ($P = (-1)^J$).	34
2.2	Parent and daughter Regge trajectories (J, M^2) for charmonia (left), bottomonia (middle) and B_c (right) mesons with unnatural parity ($P = (-1)^{J+1}$).	35
2.3	Parent and daughter Regge trajectories $(n_r \rightarrow M^2)$ for charmonia (left), bottomonia (middle) and B_c (right) mesons	35
4.1	Wood-Saxon potential with variation in potential depth	48
5.1	Diagram describing meson mass operator.	54
5.2	Quark model diagrams for the D -meson leptonic decay	57
5.3	Quark model diagrams for the D -meson semileptonic decay	59
5.4	The form factors for semileptonic $D \rightarrow K, \pi$, $D_{(s)}^+ \rightarrow \eta^{(\prime)}$ and $D_s^+ \rightarrow K^0$ transitions with comparison to LCSR, LFQM and CQM.	64
5.5	The form factors for semileptonic $D_s^+ \rightarrow \phi$ (left) and $D_s^+ \rightarrow K^*(892)^-$ (right) transitions with comparison to LFQM, HM χ T and CQM.	65
5.6	The form factors for semileptonic $D^+ \rightarrow K^*$ (left), $D \rightarrow \rho$ (middle) and $D \rightarrow \omega$ (right) transitions with comparison to LFQM, HM χ T, CQM and CLEO data.	66
5.7	$D \rightarrow \pi(K)$ form factors obtained in our model (solid lines) and in LQCD calculations (dots with error bar) by ETM collaboration.	67