

List of figures

No.	Title	Pg. No.
1.1	Image of <i>Jatropha curcas</i>	9
1.2	Images of <i>Jatropha curcas</i>	19
	←(A) Young <i>Jatropha curcas</i> plant with both flowers and developing → seedpods	
	←(B) <i>Jatropha curcas</i> inflorescence containing both male staminate flowers →(M) and female pistillate flowers (F)	
	←(C) Cross-section of a <i>Jatropha curcas</i> seedpod containing three → developing seeds	
	(D) Mature seeds of <i>Jatropha curcas</i>	
1.3	Schematic representation of RFLP	28
1.4	A schematic representation of PCR	29
1.5	Principle of RAPD-PCR technique	31
1.6	A schematic flow chart showing the principle of the AFLP method	33
1.7	Kennedy Pathway	36
1.8	Pictorial representation of Ti plasmid	40
1.9	Schematic representation of plant bacterial interaction	42
1.10	The ChvE/VirA/VirG signal transducing system	43
✓ 1.11	Cloning of gene of interest into Ti plasmid and its use in plant transformation	45
1.12	Plant transformation using binary vectors	46
2.1	Callus initiation at the cut margin	69
2.2	Emerging shoot buds from callus	70
2.3	Shoot bud initiation and proliferation	73
2.4	Regenerants achieved from callus	75

2.5	Root formation	77
2.6	Emergence of new shoot bud	78
2.7	Multiplication of nodal culture	78
3.1	Summary of culturing of carrot somatic embryos	85
3.2	A comparison of somatic and zygotic embryogenesis	86
3.3	Percent Embryogenesis in MS medium supplemented with 2,4-D	90
3.4	Embryogenesis in MS medium supplemented with 2,4-D	91
3.5	A-F: Cells seen in suspension culture; A- Elongated cells B -Round cells, C - Globular embryos, D - Cell aggregate with a globular embryo, E - Heart shaped cells, F - Torpedo shaped cells.	94
3.6	G-J: Stages in direct somatic embryogenesis as observed on solid media: G-Globular embryos seen at the edge, H: Small nodular embryos seen at the edge and veinlets, I- Heart shaped embryo, J- Direct somatic embryogenesis	95
3.7	Comparative auxin levels in MS +2.4, 3.6 and 4.0mg/l 2,4-D	96
3.8	Comparative auxin levels in MS +0.4, 0.8 and 1.2mg/l 2,4-D+(10%)CW	97
3.9	Comparative auxin levels in MS +0.4, 0.8 and 1.2mg/l 2,4-D+(2%)CW	97
4.1	A) The general pathway of TAG synthesis and accumulation.	105
	B) The bottlenecks in engineering plants to accumulate TAG	105
4.2	Predicted topological maps of DGAT1 and DGAT2	107
4.3	Map of pE172	115
4.4	Map of pE309	116
4.5	Growth of transformants on LA plates containing tetracyclin(10 µg/ml),rifampicin(5 µg/ml) and kanamycin(50 µg/ml).	119
4.6	A) Conformation of transformation (pE172)	120
	B) Conformation of transformation (pE309)	121

4.7	Co-cultivation of transformed <i>Agrobacterium</i> and <i>Jatropha</i> leaf discs	123
4.8	A) Somatic embryos on the edges of cotyledonary leaf discs	124
	B) Co-cultivation of somatic embryos on MS medium	124
	C) Transformed embryos on MS+ Kan	124
4.9	Effect of explants pre-culture period on transformation efficiency	126
4.10	Effect of wounding on transformation efficiency	127
4.11	Effect of acetosyringone on transformation efficiency	128
4.12	Effect of period of co-cultivation on transformation efficiency	129
5.1	Electrophoresis of Genomic DNA of <i>J.curcas</i>	144
5.2	Electrophoresis of <i>J.curcas</i> DNA digested with <i>Hind</i> III on 1% agarose gel	145
5.3	Electrophoretic separation of genomic DNA extracted from leaf using different incubation time at 65°C	147
5.4	Amplification of <i>J.curcas</i> by RAPD in varying MgCl ₂	152
5.5	RAPD profile generated with RAN3	157
5.6	RAPD profile generated with RAN10	157
5.7	RAPD profile generated with RAN14	158
5.8	RAPD profile generated with RBA14	158
5.9	RAPD profile generated with RBA13	159
5.10	RAPD profile generated with RFU6	159
5.11	RAPD profile generated with RFU10	160
5.12	Phylogenetic tree analysis	160