



LITERATURE CITED



- Achard, F., Eva, H., and Mayaux, P. (2001). Tropical forest mapping from coarse spatial resolution satellite data: production and accuracy assessment issues. *International Journal of Remote sensing*, **22**, 2741-2762.
- Achard, R., Eva, H. D., Stibig, H. J., Mayaux, P., Gallego, J., Richards, T., et al. (2002). Determination of deforestation rates of the world's humid tropical forests. *Science*, **297**(5583), 999-1002.
- Alcoverro, T., Cerbian, E., and Ballesteros, E. (2001). The photosynthetic capacity of the seagrass *Posidonia oceanica*: Influence of nitrogen and light. *Journal of Experimental Marine Biology*, **261**, 107-120.
- Allison, S.D., and Vitousek, P.M. (2004). Rapid nutrient cycling in leaf litter from invasive plants in Hawaii. *Oecologia*, **141**(4), 612-619.
- Aplin, P., Atkinson, P., and Curran, P. (1997). Fine spatial resolution satellite sensors for the next decade. *International Journal of Remote Sensing*, **18**, 3873-3881.
- Arvidson, T., Gasch, J., and Goward, S.N. (2001). Landsat 7's long-term acquisition plan – an innovative approach to building a global imagery archive. *Remote Sensing of Environment*, **78**, 13-26.
- Asner, G.P. (1998). Biophysical and biochemical sources of variability in canopy reflectance. *Remote sensing of Environment*, **64**(3), 234-253.
- Asner, G.P., and Heidebrecht, K.B. (2003). Imaging spectroscopy for desertification studies: comparing AVIRIS and EO-1 Hyperion in Argentina Drylands. *IEEE Transactions on Geoscience and Remote Sensing*, **41**, 1283-1296.
- Asner, G.P., and Lobell, D.B. (2000). A biogeophysical approach for automated SWIR unmixing of soils and vegetation. *Remote Sensing of Environment*, **74**, 94-112.
- Asner, G.P., and Wessman, C.A. (1997). Scaling PAR absorption from the leaf to landscape level in spatially heterogeneous ecosystems. *Ecological Modelling*, **101**, 145-163.
- Asner, G.P., Braswell, B.H., Schimel, D.S., and Wessman, C.A. (1998a). Ecological research needs from multi-angle remote sensing data. *Remote Sensing of Environment*, **63**, 155-165.
- Asner, G.P., Wessman, C.A., Bateson, C.A., and Privetter, J.L. (2000). Impact of tissue, canopy, and landscape factors on the hyperspectral reflectance variability of arid ecosystems. *Remote Sensing of Environment*, **74**, 69-84.
- Aspinall, R.J., Andrew Marcus, W., and Boardman, J.W. (2002). Considerations in collecting, processing, and analyzing high spatial resolution hyperspectral data for environmental investigations. *Journal of Geographical Systems*, **4**, 15-29.
- Bahuguna, V.K. (1999). Fire situation in India. *International Forest Fire News*. Publication No. **26**, 23-27.
- Bajjouk, T., Guillaumont, B., and Populus, J. (1996). Application of airborne imaging spectrometry system data to intertidal seaweed classification and mapping. *Hydrobiologia*, **326/327**, 463-471.
- Bajwa, S. G., Bajcsy, P., Groves, P., and Tian, L. F. (2004). Hyperspectral image data mining for band selection in agricultural application. *Transactions of the ASAE*, **43**(3), 895-907.
- Balaguru, B., Britto, S.J., Nagamrurugan, N., Natarajan, D., Soosairaj, S., Ravipaul, S., and Arockiasamy, D.I. (2003). Vegetation mapping and slope characteristics in Shervaryan Hills, Eastern Ghats using remote sensing and GIS. *Current Science*, **85**(5), 645-653.
- Barrett, E. C., and Curtis, L. F. (1999). *Introduction to Environmental Remote Sensing*, Stanley Thornes Ltd., pp. 457.

- Belanger, M. J., Miller, J. R., and Boyer, M. G. (1995). Comparative relationships between some red edge parameters and seasonal leaf chlorophyll concentrations, *Canadian Journal of Remote Sensing*, **21**, 16-21.
- Bhat, D.M., Naik, M.B., Patagar, S.G., Hegde, G.T., Kanade, Y.G., Hegde, G.N., Shastri, C.M., and Shetti, D.M. (2000). Forest Dynamics in tropical rain forests of Uttara Kannada district in Western Ghats, India. *Current Science*, **79**(7), 975-985.
- Bierregaard, R. O. Jr., Lovejoy, T. E., Kapos, V., Dos Santos, A.A., and Hutchings, R.W. (1992). The biological dynamics of tropical rain forest fragments. *Bioscience*, **42**, 859-866.
- Binaghi, E., Gallo, I., Boschetti, M., and Brivio, P.A. (2004). A neural adaptive model for hyperspectral data classification under minimal training conditions. Proceedings of Image and Signal Processing for Remote Sensing X, 13-16 September 2004, Maspalomas, Gran Canaria, pp. 173-181.
- Blackburn, G. A. (1998). Quantifying chlorophylls and carotenoids at leaf and canopy scales: An evaluation of some hyperspectral approaches. *Remote Sensing of Environment*, **66**, 273-285.
- Blackburn, G.A., and Milton, E.J. (1995). Seasonal variations in the spectral reflectance of deciduous tree canopies. *International Journal of Remote Sensing*, **16**(4), 709-720.
- Boardman, J.W. (1993). Automated spectral unmixing of AVIRIS data using convex geometry concepts, *Summaries of the 4th Annual JPL Airborne Geoscience Workshop*, Pasadena, CA, JPL Publ. 93-26, 1.
- Boardman, J.W., and Kruse, F.A. (1994). Automated spectral analysis: A geological example using AVIRIS data, northern Grapevine Mountains, Nevada: in *Proceedings, Tenth Thematic Conference, Geologic Remote Sensing*, 9-12 May 1994, San Antonio, Texas. p I-407 – I-418.
- Boardman, J.W., Kruse, F.A., and Green, R.O. (1995). Mapping target signatures via partial unmixing of AVIRIS data. Summaries of the 5th Annual JPL Airborne Geoscience Workshop, JPL Publication 95-1, **1**, 23-26.
- Boochs, F., Kupfer, G., Dockter, K., and Kuhbauch, W. (1990). Shape of the red-edge as vitality indicator for plants. *International Journal of Remote Sensing*, **11**(10), 1741-1753.
- Borregaard, T., Nielsen, H., Norgaard, L., and Have, H. (2000). Crop-weed discrimination by line imaging spectroscopy. *Journal of Agriculture & Engineering Research*, **75**, 389-400.
- Botkin, D. B., Estes, J. E., McDonald, R. M., and Wilson, M. V. (1984). Studying the Earth's vegetation from space. *BioScience*, **34**, 508.
- Brandtberg, T. (1999). Automatic individual tree based analysis of high spatial resolution aerial images on naturally regenerated boreal forests. *Canadian Journal of Forest Research*, **29**, 1461-1478.
- Brown, J.K., and Bevins, C.D. (1986). Surface fuel loadings and predicted fire behavior for vegetation types in the northern Rocky Mountains. US Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT. Res. Note: INT-358, pp. 9.
- Bubier, J.L, Rock, B.N., and Crill, P.M. (1997). Spectral Reflectance Measurements of Boreal Wetland and Forest Mosses. *Journal of Geophysical Research – Atmospheres*, **102**, (D24) 29483 – 29494.
- Buddenbaum, H., Schlerf, M., and Hill, J. (2005). Classification of coniferous tree species and age classes using hyperspectral data and geostatistical methods. *International Journal of Remote Sensing*, **26** (24, 20), 5453-5465.

- Cardner, K.L., Reinersman, P., Chen, R.F., Muller-Karger, F., Davis, C.O., and Hamilton, M.K. (1993). AVIRIS calibration and application in coastal oceanic environments. *Remote Sensing of Environment*, **44**(2), 205-216.
- Carrere, V., and Conel, J. (1993). Recovery of atmospheric water vapor total column abundance from imaging spectrometer data around 940nm – sensitivity analysis and application to AVIRIS data. *Remote sensing of Environment*, **44**(2-3), 179-204.
- Carter, G. A. (1993). Responses of leaf spectral reflectance to plant stress. *American Journal of Botany*, **80**(3), 239-243.
- Castro-Esau, K.L., Sanchez-Azofeifa, G.A., and Caelli, T. (2004). Discrimination of lianas and trees with leaf-level hyperspectral data. *Remote Sensing of Environment*, **90**, 353–372.
- Chen, F., Niu, Z., Sun, G.Y., Wand, C. Y., and Teng, J. (2007). Using low-spectral-resolution images to acquire simulated hyperspectral images. *International Journal of Remote Sensing*, **29**(10), 2963-2980.
- Cho, M. A., and Skidmore, A. K. (2006). A new technique for extracting the red edge position from hyperspectral data: The linear extrapolation method. *Remote Sensing of Environment*, **101**(2), 181-193.
- Chokkalingam, U., and White, A. (2001). Structure and spatial patterns of trees in old-growth northern hardwood and mixed forests of northern Maine. *Plant Ecology*, **156**(2), 139–160.
- Christian, B., and Krishnayya, N.S.R. (2007). Spectral signatures of teak (*Tectona grandis* L.) using hyperspectral (EO-1) data. *Current Science*, **93**(9), 1291-1296.
- Christian, B., and Krishnayya, N.S.R. (2008). Discrimination of floor cover of dry deciduous forest using Hyperion (EO-1) data. In press.
- Clark, D.A. (2004a). Tropical forests and global warming: Slowing it down or speeding it up? *Frontiers of Ecology & Environment*, **2**, 73–80.
- Clark, D.A. (2007). Detecting tropical forests' responses to global climatic and atmospheric change: Current challenges and a way forward. *Biotropica*, **39**(1), 4-19.
- Clark, D.A., Piper, S.C., Keeling, C.D., and Clark, D.B. (2003). Tropical rain forest tree growth and atmospheric carbon dynamics linked to interannual temperature variation during 1984-2000. *Proceedings of the National Academy of Sciences of the United States of America*, **100**(10), 5852-5857.
- Clark, M.L., Roberts, D.A., and Clark, D.B. (2005). Hyperspectral discrimination of tropical rain forest tree species at leaf to crown scales. *Remote sensing of Environment*, **96**, 375-398.
- Clark, R. N. (1999). Spectroscopy of rocks and minerals, and principles of spectroscopy. *Manual of Remote Sensing*, Vol-3, Remote Sensing for the Earth Sciences (A.N. Rencz, editor), John Wiley and Sons, New York, pp-137.
- Clark, R.N., and Roush, T.L. (1984). Reflectance spectroscopy: Quantitative analysis techniques for remote sensing applications. *Journal of Geographical Research*, **89**, 6329-6340.
- Clark, R.N., and Swayze, G.A. (1995). Automated spectral analysis: Mapping minerals, amorphous materials, environmental materials, vegetation, water, ice, and snow, and other materials: The USGS Tricorder Algorithm. *Lunar and Planetary Science xxvi*, 255-256.
- Clevers, J.G.P.W., de Jong, S.M., Epema, G.F., Van der Meer, F., Bakker, W.H., Skidmore, A.K., Scholte, K.H. (2002). Derivation of the red edge index using MERIS standard band setting. *International Journal of Remote Sensing*, **23**(16), 3169-3184.
- Clevers, J.G.P.W., Kooistra, L., and Salas, E.A.L. (2004). Study of heavy metal contamination in river floodplains using the red-edge position in spectroscopic data. *International Journal of Remote Sensing*, **25**(19), 3883-3895.

- Cochrane, M.A. (2000). Using vegetation reflectance variability for species level classification of hyperspectral data. *International Journal of Remote Sensing*, **21**, 2075-2087.
- Cochrane, M.A., Alencar, A., Schulze, M.D., Souza, C.M., Nepstad, D.C., and Lefebvre, P. (1999). Positive feedbacks in the fire dynamic of closed canopy tropical forests. *Science*, **284**(5421), 1832-1835.
- Congalton, R.G. (1991). A review of assessing the accuracy of classifications of remotely sensed data. *Remote Sensing of Environment*, **37**, 35-46.
- Congalton, R.G., and K. Green. (1999). *Assessing the Accuracy of Remotely Sensed Data: Principles and Practices*, Lewis Publishers, New York, pp. 137.
- Cooke, T. (1967). *The Flora of Presidency of Bombay*, Botanical Survey of India, Calcutta.
- Cramer, W., Bondeau, A., Woodward, F. I., Prentice, I. C., Betts, R. A., Brovkin, V., Cox, P. M., Fisher, V., Foley, A., Friend, A. D., Kucharik, C., Lomas, M. R., Raman Kutty, N., Sitch, S., Smith, B., White, A., and Young-Molling, C. (2001). Global response of terrestrial ecosystem structure and function to CO₂ and climate change: Results from six dynamic global vegetation models. *Global Change and Biology*, **7**, 357-373.
- Crowley, J.K. (1993). Mapping playa evaporate mineral with AVIRIS data: a first report from Death Valley, California. *Remote Sensing of Environment*, **44**(2-3), 337-356.
- Curran, P. J., Dungan, J. L., Macler, B. A., and Plummer, S. E. (1991). The effect of a red leaf pigment on the relationship between red edge and chlorophyll concentration. *Remote Sensing of Environment*, **35**, 69-76.
- Curran, P. J., Dungan, J. L., Macler, B. A., Plummer, S. E., and Peterson, D. L. (1992). Reflectance spectroscopy of fresh whole leaves for the estimation of chemical concentration. *Remote Sensing of Environment*, **39**(2), 153-166.
- Curran, P. J., Windham, W. R., and Gholz, H. L. (1995). Exploring the relationship between reflectance red edge and chlorophyll concentration in slash pine leaves. *Tree Physiology*, **15**, 203-206.
- Curran, P.J. (1989). Remote sensing of foliar chemistry. *Remote Sensing of Environment*, **30**(3), 271-278.
- Curran, P.J., Foody, G.M., Lucas, R.M., Honzak, M., and Grace, J. (1997). The carbon balance of tropical forests: From the local to the regional scale. In P.R. van Gardingen, Foody, G.M., and Curran, P.J. (Eds.), *Scaling-up from cell to landscape*. Cambridge: Cambridge University Press, pp.201-227.
- Dai, C.D., and Lei, L.P. (1989). The spectral characteristics and optimal band combination of TM image. *Chinese Remote Sensing of Environment*, **4**, 282-292.
- Danson, F. M., Steven, M. D., Malthus, T. J., and Clark, J. A. (1992). High-spectral resolution data for determining leaf water content. *International Journal of Remote Sensing*, **13**, 461-470.
- Danson, F.M. (1995). Developments in the remote sensing of forest canopy structure. In *Advances in Environmental Remote Sensing*, edited by F.M. Danson and S. E. Plummer (Chichester: John Wiley & Sons), pp. 53-69.
- Danson, F.M., and Plummer, S.E. (1995). Red-edge response to forest leaf area index. *International Journal of Remote Sensing*, **16**(1), 183-188.
- Datt, B. 2000. Recognition of eucalyptus forest species using hyperspectral reflectance data, In *Proceedings of the IEEE International Geosciences Remote Sensing Symposium*, Honolulu, Hawaii, **4**, pp. 1405-1407.

- Datt, B., Mc Vicar, T.R., Jupp, D.L.B., and Pearlman, J.S. (2003). Preprocessing EO-1 Hyperion hyperspectral data to support the application of agricultural indexes. *IEEE Transaction Geoscience Remote Sensing*, **41**(2), 1246-1259.
- Daughtry, C. S. T., Gallo, K. P., Goward, S. N., Prince, S. D., and Kustas, W. P. (1992). Spectral estimates of absorbed radiation and phytomass production in corn and soybean canopies. *Remote Sensing of Environment*, **39**(2), 141-152.
- Daughtry, C.S.T., Hunt Jr. E.R., and McMurtrey III. J.E. (2004). Assessing crop residue cover using shortwave infrared reflectance, *Remote Sensing of Environment*, **90**: 126-134.
- Daughtry, C.S.T., McMurtrey, J.E.III., Nagler, P.L., Kim, M.S., and Chappelle, E.W. (1996). Spectral reflectance of soils and crop residues. In *near Infrared Spectroscopy: The Future Waves* (A.M.C. Davies and Phil Williams, Eds.), NIR publications, Chichester, UK, pp. 505-511.
- Dawson, S.P., and Dennison, W.C. (1996). Effects of ultraviolet and photosynthetically active radiation on five seagrass species. *Marine Biology* **125**, 629-638.
- Dawson, T. P., and Curran, P. J. (1998). A new technique for interpolating red edge position. *International Journal of Remote Sensing*, **19**(11), 2133-2139.
- Dawson, T. P., Curran, P. J., North, P. R. J., and Plummer, S. E. (1999). The propagation of the foliar biochemical absorption features in forest canopy reflectance: a theoretical analysis. *Remote Sensing of Environment*, **67**, 147-159.
- De Jong, B.H.J., Gaona, S.O., Santiago, M.A.C., Marcial, N.R., and Cairns, M.A. (2000). Carbon flux and patterns of land-use/land-cover change in the Selva Lacandona, Mexico. *Ambio*, **29**(8), 504-511.
- de Vasconcelos, J.J.P., Paul, J.C.U., Silva, S., Pereira, J.M.C., Caetono, M.S., Catry, F.X., and Oliveira, T.M. (1998). Regional fuel mapping using knowledge based system approach. In *proceedings: III international conference on forest fire research and 14th conference on fire and forest meteorology*, **2**, pp. 2111-2133.
- DeFries, R.S., and Townshend, J.R.G. (1999). Global land cover characterization from satellite data: from research to operational implementation? *Global Ecology and Biogeography*, **8**, 367-379.
- Demarez, V., Gastellu-Etchegorry, J.P., Mougin, E., Marty, G., and Proisy, C. (1999). Seasonal variation of leaf chlorophyll content of a temperate forest. Inversion of the PROSPECT model. *International Journal of Remote Sensing*, **20**, 879-894.
- Dennis, R.A., and Colfer, C.P. (2006). Impacts of land use and fire on the loss and degradation of lowland forest between 1983-2000 in East Kutai District, East Kalimantan. *Singapore Journal of Tropical geography*, **27**(1), 36-58.
- Dennison, P.E., and Roberts, D.A (2003). Endmember selection for multiple endmember spectral mixture analysis using endmember average RMSE. *Remote Sensing of Environment*, **87**, 295-309.
- Deota, B.S. (1991). Geological studies in the South Gujarat Quarternary landscape with special reference to environmental planning and management. Ph.D Thesis, M.S.University, Baroda.
- Dixon, R.K., and Turner, D.P. (1991). *Environmental Pollution*, **73**, 245.
- Dixon, R.K., Brown, S., Houghton, R.A., Solomon, A.M., Trexler, M.C., and Wisniewski, J. (1994). Carbon pools and flux of global forest ecosystems. *Science*, **262**, 185-190.
- Doherty, R.M., Hulme, M., and Jones, C.G. (1999). A gridded reconstruction of land and ocean precipitation for the extended tropics from 1974 to 1994. *International Journal of Climatology*, **19**, 119-142.

- Drake, N.A., Mackin, S., and Settle, J.J. (1999). Mapping Vegetation, Soils, and Geology in Semiarid Shrublands Using Spectral Matching and Mixture Modeling of SWIR AVIRIS Imagery. *Remote Sensing of Environment*, **68**, 12 - 25.
- Drohan, P.J., Stout, S.L., and Petersen, G.W. (2002). Sugar maple (*Acer saccharum* Marsh.) decline during 1979–1989 in northern Pennsylvania. *Forest Ecology and Management*, **170**(1–3), 1–17.
- Elands, B.H.M., and T.N. O’Leary, (2002). The myth of forests: a reflection of the variety of their rural identifies in Europe and the role of forests in it. In *The Changing Role of Forestry in Europe; perspectives for rural development*. K.F. weirsum and B.H.M Elands (eds). Wageningen University, Wageningen, the Netherlands 25-50.
- Elvidge, C.D., and Portigal, F.P. (1990). Change detection in vegetation using 1989 AVIRIS data. *SPIE Imaging Spectroscopy of the Terrestrial Environment*. SPIE, Orlando, FL, 178-1899.
- Elvidge, D. E. (1990). Visible and near infrared reflectance characteristics of dry plant materials. *International Journal of Remote Sensing*, **11**(10), 1775-1795.
- ERDAS user guide.
- Ferwerda, J. G., Skidmore, A. K., and Mutanga, O. (2005). Nitrogen detection with hyperspectral normalized ratio indices across multiple plant species. *International Journal of Remote Sensing*, **26**(18), 4083-4095.
- Ferwerda, J. G., Skidmore, A. K., and Stein, A. (2006). A bootstrap procedure to select hyperspectral wavebands related to tannin content. *International Journal of Remote Sensing*, **27**(7), 1413-1424.
- Field, D.B., Behrenfeld, M.J., Randerson, J.T., and Falkowski, P. (1998). Primary production of the biosphere: Integrating terrestrial and oceanic components. *Science*, **281**, 237–240.
- Filella, I., and Penuelas, J. (1994). The red edge position and shape as indicators of plant chlorophyll content, biomass and hydric status. *International Journal of Remote Sensing*, **15**(7), 1459-1470.
- Foley, B., McIlwee, A., Lawler, I., Agragones, I., woolnough, A. P., and Berding, N. (1998). Ecological applications of near infrared spectroscopy - a tool for rapid, cost effective prediction of the composition of plant and animal tissues and aspect of animal performance. *Oecologia*, **116**, 293-305.
- Foley, J.A., Dfries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N., and Snyder, P.K. (2005). Global consequences of land use. *Science*, **309**, 570–574.
- Foody, G.M. (2003). Remote sensing of tropical forest environment: Towards the monitoring of environmental resources for sustainable development. *International Journal of Remote Sensing*, **24**(20), 4035-4046.
- Foody, G.M., Boyd, D.S., and Cutler, M.E.J. (2003). Predictive relations of tropical forest biomass from Landsat TM data and their transferability between regions. *Remote Sensing of Environment*, **85**(4), 463-474.
- Foody, G.M., Palubinskas, G., Lucas, R.M., Honzak, M., and Curran, P.J. (1996). Identifying terrestrial carbon sinks: Classification of successional stages in regenerating tropical forest from Landsat TM data. *Remote sensing of Environment*, **55**, 205-216.
- Forest survey of India, (1996). State of Forest Report.
- Forest Survey of India, (2003). State of Forest Report.

- Foreword to the Earth Observing 1 Spectral Issue (2003). *IEEE Transaction Geoscience Remote Sensing*, **41**(6), 1147-1148.
- Fourty, Th., Baret, F., Jacquemoud, S., Schmuck, G., and Verdebout, J. (1996). Leaf optical properties with explicit description of its biochemical composition: direct and inverse problems. *Remote Sensing of Environment*, **56**, 104-117.
- Frank, T. D. (1988). Mapping dominant vegetation communities in the Colorado Rocky Mountain Front Range with Landsat Thematic Mapper and digital terrain data. *Photogrammetric Engineering and Remote Sensing*, **54**, 1727-1734.
- Franklin, S. E. (1994). Discrimination of subalpine forest species and canopy density using CASI, SPOT, PLA, and Landsat TM data. *Photogrammetric Engineering and Remote Sensing*, **60**, 1233-1241.
- Fuller, D.O. (2006). Tropical forest monitoring and remote sensing: A new era of transparency in forest governance? *Singapore Journal of Tropical Geography*, **27**, 15-29.
- Fung, I.Y., Doney, S.C., Lindsay, K., and John, J. (2005). Evolution of carbon sinks in a changing climate. *Proceedings of National Academic Science, USA*, **102**, 11201-11206.
- Fung, T., Ma, F.Y., and Siu, W.L. (1999). Hyperspectral data analysis for subtropical tree species recognition. *Symposium proceedings, IGRASS'98, sensing and managing the environment*, **3**, 1298-1300.
- Gamon, J. A., and Surfus, J. S. (1999). Assessing leaf pigment content and activity with a reflectometer. *New Phytologist* **143**(1), 105-117.
- Gamon, J.A., Field, C.B., Roberts, D.A., Ustin, S.L., and Valentini, R. (1993). Functional patterns in an annual grassland during and AVIRIS overflight. *Remote Sensing of Environment*, **44**, 239-254.
- Gao, B., and Goetz, A.F.H. (1990). Column atmospheric water vapor and vegetation liquid water retrievals from airborne imaging spectrometer data. *Journal of Geophysical Research*, **95**(D4), 3549-3564.
- Gausman, H.W. (1985). Plant leaf optical properties in visible and near infrared light, Lubbock, Texas, Texas Tech Press.
- Geist, H.J., and Lambin, E.F. (2002). Proximate causes and underlying driving forces of tropical deforestation. *Bioscience*, **52**, 143-150.
- Gilabert, M.A., García-Haro, F.J., and Meliá, J. (2000). A mixture modeling approach to estimate vegetation parameters for heterogeneous canopies in remote sensing, *Remote Sensing of Environment*, **72**, 328-345.
- Giri, C., Zhu, Z., and Reed, B. (2005). A comparative analysis of the Global Land Cover-2000 and MODIS land covers data sets. *Remote Sensing of Environment*, **94**, 123-132.
- Gitelson, A. A., Merzlyak, M. N., and Lichtenthaler, H. K. (1996). Detection of red-edge position and chlorophyll content by reflectance measurements near 700 nm. *Journal of plant physiology*, **148**, 494-500.
- Goetz, A.F.H., Vane, G., Solomon, J.E., and Rock, B.N. (1985). Imaging spectrometry for earth remote sensing. *Science*, **211**, 1147-1153.
- Gong, P., Pu, R.L., and Yu, B. (1997). Conifer species recognition: An exploratory analysis of in situ hyperspectral data. *Remote Sensing of Environment*, **62**, 189-200.
- Goodenough, D.G., Dyk, A., Olaf Niemann, K., and Pearlman, J. (2003) Processing Hyperion and ALI for Forest Classification. *IEEE Transaction Geoscience Remote Sensing*, **41**(6), 1321-1331.
- Goparaju, L., Tripathi, A., and Jha, C.S. (2005). Forest fragmentation impacts on phytodiversity-An analysis using remote sensing and GIS. *Current Science*, **88**(25), 1264-1274.

- Gower, S. T., Kucharik, C. J., and Norman, J. M. (1999). Direct and Indirect Estimation of Leaf Area Index, fAPAR, and Net Primary Production of Terrestrial Ecosystems. *Remote Sensing of Environment*, **70**(1), 29-51.
- Gracia, M., and Ustin, S.L. (2001). Detection of interannual vegetation responses to climatic variability using AVIRIS data in a coastal savanna in California, *IEEE Transactions on Geosciences and Remote Sensing*, **39**, 1480-1490.
- Grant, L. (1987). Diffuse and secular characteristics of leaf reflectance. *Remote Sensing of Environment*, **22**, 309-322.
- Green, A.A., Berman, M., Switzer, P., and Craig, M.D. (1998). A transformation for ordering multispectral data in terms of image quality with implications for noise removal. *IEEE Transaction Geoscience Remote Sensing*, **26**, 65-74.
- Green, R. O., Conel, J. E., and Chrien, T. G. (1992). Airborne Visible-Infrared Imaging Spectrometer (AVIRIS): Sensor System, In-flight Calibration and Reflection Calculation. *Jet Propulsion Laboratory, ISSSR'92 Proceedings*, **1**, pp. 198.
- GSFC Systems Engineering Seminar: EO-1 Results, 2002.
- Guyot, G., and Baret, F. (1988). Utilisation de la haute résolution spectrale pour suivre l'état des couverts végétaux. Proceedings of the 4th International colloquium on spectral signatures of objects in remote sensing. ESA SP-287, Assois, France, pp. 279-286.
- Hamilton, M.K., Davis, C.O., Rhea, W.J., Pilorz, S.H., and Carder, K.L. (1993). Estimating chlorophyll content and bathymetry of Lake Tahoe using AVIRIS data. *Remote Sensing of Environment*, **44**(2), 217-230.
- Hansen, J., Ruedy, R., Glascoe, J., and Sato, M. (1999). GIS analysis of surface temperature change. *Journal of Geophysical Research*, **104**, 30997-31022.
- Hansen, M.C., and Reed, B. (2000). A comparison of the IGBP Discover and University of Maryland 1 km global land covers products. *International Journal of Remote Sensing*, **21**, 1365-1373.
- Hoffbeck, J.P., and Landgrebe, D.A. (1996). Covariance matrix estimation and classification with limited training data. *IEEE Transactions on Pattern Analysis & Machine Intelligence*, **18**(7), 763-767.
- Horler, D.N.H., Dockray, M., and Barber, J. (1983). The red edge of plant leaf reflectance. *International Journal of Remote Sensing*, **4**(2), 273-288.
- Houghton, R. A., and Woodwell, G. M. (1981). Biotic contributions to the global carbon cycle: the role of remote sensing; *Proc Seventh International Symposium on Machine Processing of Remote Sensed Data*, West Lafayette, Indiana, pp. 593-602.
- Hubbell, S. P. (1997). A unified theory of biogeography and relative species abundance and its application to tropical rain forests and coral reefs. *Coral Reefs*, **16**, S9-S21 (suppl).
- Hubbell, S. P. (2001). The unified neutral theory of biodiversity and biogeography. Princeton, NJ: Princeton University Press.
- Hughes, L. (2000). Biological consequences of global warming: is the signal already apparent? *Trends in Ecology and Evolution*, **15**, 56-61.
- Hulme, M., Osborn, T.J., and Johns, T.C. (1998). Precipitation sensitivity to global warming: comparison of observations with HadCM2 simulations. *Geophysical Research Letters*, **25**, 3379-3382.
- Hunt, E. R., McMurtrey, J. E., Parker Williams, A. E., and Corp, L. A. (2004). Spectral characteristics of leafy spurge (*Euphorbia esula*) leaves and flower bracts. *Weed Science*, **52**(4), 492-497.

- Hunt, E.R.J., Everitt, J.H., Ritchie, J.C., Moran, S.M., Booth, D.T., Anderson, G.L., Clark, P.E., and Seyfried, M.S. (2003). Applications and Research Using Remote Sensing for Rangeland Management. *Photogrammetric Engineering and Remote Sensing*, **69**(6), 675-693.
- ImSpec, 2004. The ACORN home page, URL: <http://www.imspec.com/index.html>.
- Ingle, J.D. (1988). *Spectrochemical Analysis*, Prentice-Hall, Inc., Engelwood Cliffs, New Jersey, pp. 424-425.
- Iverson, L. R., Cook, E. A., and Graham, R. L. (1989). A technique for extrapolating and validating forest cover across large regions: calibrating AVHRR data with TM data. *International Journal of Remote Sensing*, **10**, 1805-1812.
- Iverson, L.R., Dale, M.E., Scott, C.T., and Prasad, A. (1997). A GIS derived integrated moisture index to predict forest composition and productivity of Ohio forests (USA), *Landscape Ecology*, **12**(5), 331-348.
- Jackson, Q., and Landgrebe, D.A. (2001). An adaptive classifier design for high-dimensional data analysis with a limited training data set. *IEEE Transactions on Geoscience and Remote Sensing*, **39**(12), 2664-2679.
- Jacobsen, A. (2000). Analysing Airborne Optical Remote Sensing Data from a Hyperspectral Scanner and Implications for Environmental Mapping and Monitoring - results from a study of CASI data and Danish semi-natural, dry grasslands. PhD thesis. National Environmental Research Institute, Denmark, pp. 148.
- Jacquemoud, S., Ustin, S. L., Verdebout, J., Schmuck, G., Andreoli, G., and Hosgood, B. (1996). Estimating leaf biochemistry using the PROSPECT leaf optical properties model. *Remote Sensing of Environment*, **56**, 194-202.
- Jago, R.A., Cutler, M.E.J., and Curran, P.J. (1999). Estimating canopy chlorophyll concentration from field and airborne spectra. *Remote sensing of Environment*, **68**, 217-224.
- Jain, A., Ravan, S.A., Singh, R.K., Das, K.K., and Roy, P.S. (1996). Forest fire risk modeling using remote sensing and geographical information systems. *Current science*, **70**(10), 928-933.
- Jakubauskas, M.E. (1996). Thematic mapper characterization of lodgepole pine seral stages in Yellowstone National Park, USA. *Remote Sensing of Environment*, **56**, 118-132.
- Jakubauskas, M.E., and Price, K.E. (1997). Empirical relationships between structural and spectral factors of Yellowstone Lodgepole pine forests. *Photogrammetric Engineering and Remote Sensing*, **63**(12), 1375-1381.
- Jaykumar, S., Arockiasamy, D.I., and John B.S. (2002). Conserving forests in the Eastern Ghats through remote sensing and GIS – A case study in Kolli hills. *Current Science*, **82**, 1259-1266.
- Johnson, L. F., Hlavka, C. A., and Peterson, D. L. (1994). Multivariate analysis of AVIRIS data for canopy biochemical estimation along the Oregon transect. *Remote Sensing of Environment*, **47**, 216-230.
- Jupp, D.P.L., Datt, B., Lovell, J., and King, E. (2002). EO-1/ Hyperion data workshop notes, Canberra, Australia, CSIRO Office of space science and Applications, Earth Observation Centre, April 2002.
- Kallio, K., Kutser, T., Hannonen, T., Koponen, S., Pulliainen, J., Vepsäläinen, J., and Pyhälähti, T. (2001). Retrieval of water quality from airborne imaging spectrometry of various lake types in different seasons. *Science of the Total Environment*, **268**, 59-77.

- Kasturirangan, K. (2004). Science and technology of imaging from space. *Current Science*, **87**(5), 584-601.
- Keane, R.E., Rollins, M.G., McNicoll, C.H., and Persons, R.A. (2002). Integrating ecosystem sampling, gradient modeling, remote sensing, and ecosystem simulation to create spatially explicit landscape inventories. US Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, CO. RMRS-GTR-92, pp. 61.
- Kent, M., and Coker, P. (1992). *Vegetation description and analysis, A practical approach*. London, John Wiley & Sons.
- Kerr, R.A. (2006). Climate change: a worrying trend of less ice, higher seas. *Science*, **311**, 1698-1701.
- King, T. V. V., Clark, R. N., and Swayze, G. A. (2000). Applications of imaging spectroscopy data: a case study at Summitville, Colorado. In F. Kuehn, T. V. V. King, B. Hoerig, & D. C. Peters (Eds.), *Remote sensing for site characterization* New York: Springer, pp. 164-185.
- Kirchner, J. W. (2002). The Gaia hypothesis: fact, theory, and wishful thinking. *Climate Change*, **52**, 391-408.
- Knapp, A. K., and Carter, G. A. (1998). Variability in leaf optical properties among 26 species from a broad range of habitat. *American Journal of Botany*, **85**(7), 940-946.
- Knipling, E. B. (1970). Physical and physiological basis for the reflectance of visible and near-infrared radiation from vegetation. *Remote Sensing of Environment*, **1**, 155-159.
- Koger, C. H., Shaw, D.R., Reddy, K.N., and Bruce, L.M. (2004). Detection of pitted morning glory (*Ipomoea lacunosa*) with hyperspectral remote sensing. II Effects of vegetation ground cover and reflectance properties. *Weed Science*, **52**, 230-235.
- Kokaly, R. F., Clark, R. N., and Livo, K. E. (1998). Mapping the biology and mineralogy of Yellowstone National Park using imaging spectroscopy. In R. O. Green (Ed.), *Summaries of the 7th annual JPL airborne earth sciences workshop*. JPL Publications, **97-21**, 245-254.
- Kokaly, R.F. (2001). Investigating a physical basis for spectroscopic estimates of leaf nitrogen concentration. *Remote Sensing of Environment*, **75**, 153-161.
- Kokaly, R.F., and Clark, R.N. (1999). Spectroscopic determination of leaf biochemistry using band-depth analysis of absorption features and stepwise multiple linear regression. *Remote Sensing of Environment*, **67**, 267-287.
- Kokaly, R.F., Despain, D.G., Clark, R.N., and Eric Livo, K. (2003). Mapping vegetation in Yellowstone National Park using spectral feature analysis of AVIRIS data. *Remote Sensing of Environment*, **84**, 437-456.
- Korobov, R.M., and Railyan, V.Y. (1993). Canonical correlation relationships among spectral and phytometric variables for twenty winter wheat fields. *Remote Sensing of Environment*, **43**, 1-11.
- Kraft, M., Weigel, H., Mejer, G., and Brandes, F. (1996). Reflectance measurements of leaves for detecting visible and non-visible ozone damage to crops. *Journal of plant physiology*, **148**, 148-154.
- Kruse, F.A., Leafkoff, A.B., and Dietz, J.B. (1993). Expert System-Based Mineral Mapping in northern Death Valley, California/Nevada using the Airborne Visible/Infrared Imaging Spectrometer (AVIRIS). *Remote Sensing of Environment*, special issue on AVIRIS, May-June 1993, **44**, 309-336.

- Kumar, L., and Skidmore, A. K. (1998). Use of derivative spectroscopy to identify regions of difference between some Australian eucalypt species. *Proceedings of the 9th Australasian Remote Sensing and Photogrammetry Conference*. Sydney, Australia.
- Kumar, L., Schmidt, K., Dury, S., and Skidmore, A. (2001). Imaging Spectrometry & Vegetation Science in *Imaging Spectrometry, Basic Principles and Prospective Applications*, Edited by Van Der Meer, F. D., Kluwer Academic Publishers, pp.111.
- Lamb, D.W., Steyn-Ross M., Schaare P., Hanna, M.M., Silvester, W., and Steyn-Ross, A. (2002). Estimating leaf nitrogen concentration in ryegrass (*Lolium* spp.) pasture using the chlorophyll red-edge: theoretical modeling and experimental observations. *International Journal of Remote Sensing*, **23**(18), 3619-3648.
- Lambert, N.J., Ardo, J., Rock, B.N., and Vogelmann, J.E. (1995). Spectral characterization and regression-based classification of forest damage in Norway spruce stands in Czech Republic using Landsat Thematic Mapper data. *International Journal of Remote Sensing*, **16**, 1261-1287.
- Landgrebe, D. (2003). *Signal Theory Methods in Multispectral Remote Sensing* (New York: Wiley-Interscience).
- Landgrebe, D.A. (1999). Some Fundamentals and Methods for Hyperspectral Image Data Analysis. *Proceedings of the International Society for Optical Engineering (SPIE): Systems and Technologies for Clinical Diagnostics and Drug Discovery II* (January 24-25, 1999, San Jose, California, USA), **3603**, 104-113.
- Lass, L. W., Thill, D. C., Shafii, B., and Prather, T. S. (2002). Detecting Spotted Knapweed (*Centaurea maculosa*) with Hyperspectral Remote Sensing Technology. *Weed Technology*, **16**(2), 426-432.
- Laurance, W. F., and Bierregaard, R.O. (1997). *Tropical forest remnants* (Chicago: University of Chicago Press).
- Laurance, W.F., Oliveira, A.A., Laurance, S.G., Condir, R., Nascimento, H.E.M., and Sanchez-Thorin, A.C. (2004). Pervasive alteration of tree communities in undisturbed Amazonian forests. *Nature*, **438**, 171-175.
- Lawrence, W.T., Shimabukuro, Y.E., and Gao, B. (1993). Analysis of Hyperspectral AVIRIS Image Data over a Mixed-Conifer Forest in Maine. *Proceedings: 7th Brazilian Remote Sensing Symposium* (Curitiba, Brazil, May 10-14, 1993), pp. 129-155.
- Leckie, D.G., Gougeon, F.A., Walsworth, N., and Paradine, D. (2003). Stand delineation and composition estimation using semiautomated individual tree crown analysis, *Remote Sensing of Environment*, **85**(3), 355-369.
- Lee, K. S., Kin, S. H., Ma, J. R., Kook, M. J., and Shin, J. (2005). Spectral variation of vegetative areas in leaf-off season observed by hyperspectral data. *IEEE*, pp.4347-4350.
- Leica, 2004. ATCOR for Imagine, Data Sheet, URL: <http://www.geosystems.de/atcor/index.html>.
- Leigh, E.G., Jr., Davidkar, P., Dick, C.W., Puyravaud, J.P., Terborgh, J., Ter Steege, H., and Wright, S.J. (2004). Why do some tropical forests have so many species of trees? *Biotropica*, **36**, 447-463.
- Lewis, M., Jooste, V., and De Gasparis, A. (2000). Hyperspectral discrimination of Arid vegetation. *Proceedings 28th International Symposium on Remote Sensing of Environment*, pp. 148-151.

- Li, L., Ustin, S.L., and Lay, M. (2005). Application of multiple endmember spectral mixture analysis (MESMA) to AVIRIS imagery for coastal salt marsh mapping: a case study in China Camp, CA, USA. *International Journal of Remote Sensing*, **26**(23), 5193-5207.
- Lichtenthaler, H.K., Gitelson, A.A., and Lang, M.J. (1996). Non-destructive determination of chlorophyll content of leaves of a green and aurea mutant tobacco by reflectance measurements. *Journal of Plant Physiology*, **148**, 483-493.
- Lillesand, T.M., and Kiefer, R.W. (2000). *Remote sensing and image interpretation* (fourth edition). New York: John Wiley & sons, Inc.
- Longstaff, B.J., and Dennison, W. C. (1999). Seagrass survival during pulsed turbidity events: The effects of light deprivation on the seagrasses *Halodule pinifolia* and *Halophila ovalis*. *Aquatic Botany*, **65**, 105-121.
- Lugo, A. E., and Brown, S. (1992). Tropical Forests as Sinks of Atmospheric Carbon, *Forest Ecological Management*, **54**, 239-255.
- Lumme, J.H. (2004). Classification of vegetation and soil using Imaging spectrometer data. *Proceedings XXth ISPRS Congress*, Istanbul, Turkey. July 12-23.
- Luther, J. E., and Carroll, A. L. (1999). Development of an index of balsam fir vigor by foliar spectral reflectance. *Remote Sensing of Environment*, **69**, 241-252.
- Malhi, Y., and Grace, J. (2000). Tropical forest and atmospheric carbon dioxide. *Trends in Ecology and Evolution*, **15**, 332-337.
- Malingreau, J. P. (1991). Remote Sensing for Tropical Forest Monitoring: an overview; in *Remote Sensing and Geographical Information Systems for Resource Management in Developing Countries* (eds) A S-Belward and C R Valenzuela (Dordrecht: Kluwer), pp. 253-278.
- Mark, C.A., Bushey, C.L., and Smetanka, W. (1995). Fuel model identification and mapping for fire behavior prediction in the Absaroka-Beartooth Wilderness, Montana and Wyoming. In *Proceedings: symposium on fire in wilderness and park management*. USDA Forest Service General Technical Report INT-GTR- 320, pp. 227-229.
- Martin, M. E., and Aber, J. D. (1997b), Estimation of forest canopy lignin and nitrogen concentration and ecosystem processes by high spectral resolution remote sensing. *Ecological Applications*, **7**, 431-443.
- Martin, M. E., Newman, S. D., Aber, J. D., and Congalton, J. C. (1998). Determining forest species composition using high spectral resolution remote sensing data. *Remote Sensing of Environment*, **65**, 249-254.
- Maslekar, A.R. (1974). Remote Sensing and its scope in Indian Forestry. *Indian Forester*, **100**, 192-201
- Mayaux, P., Achard, F., and Malingreau, J.P. (1998). Global tropical forest area measures derived from coarse resolution maps at a global level: a comparison with other approaches. *Environmental conservation*, **25**, 37-52.
- Mayaux, P., Holmgren, P., and Achard, F. et al. (2005). Tropical Forest cover change in the 1990s and options for future monitoring. *Philosophical Transactions of the Royal Society B-Biological Sciences*, **360**, 373-384.
- McCarty, J.P. (2001). Ecological consequences of recent climate change. *Conservation. Biology*, **15**, 320-331.
- Melillo, J.M. (1999). Warm, warm on the range. *Science*, **283**, 183-4.
- Melillo, J.M., McGuire, A. D., Kicklighter, D.W., Moore III, B., Vorosmarty, C. J., and Schloss, A. L. (1993). Global Climate Change and Terrestrial Net Primary Production, *Nature* **363**, 234-240.

- Merton, R.N. (1998). Monitoring community hysteresis using spectral shift analysis and the red-edge vegetation stress index. *Proceedings of the Seventh JPL airborne Earth Science Workshop*. Jet Propulsion Laboratory, Pasadena, CA, 275-284.
- Merton, R.N. 1992. Multi-temporal analysis of community scale vegetation stress with imaging spectroscopy. Ph.D. Thesis, Geography Department, University of Auckland, New Zealand, pp. 492.
- Miller, J.R., Jiyou, W., Boyer, M.G., Belanger, M., and Hare, E.W. (1991). Seasonal patterns in leaf reflectance red-edge characteristics. *International Journal of Remote Sensing*, **12**, 1509-1523.
- Ministry of Environment and Forest, 1994.
- Moran, E.F., Brondizio, E., Mausel, P., and Wu, Y. (1994). Integrating Amazonian vegetation, land-use, and satellite data. *Bioscience*, **44**, 329-338.
- Murli, K.S., Siddapa, R., Ganeshiah, K.N., and Umashankar, R. (1998). Does forest type classification reflect spatial dynamics of vegetation? An analysis using GIS technique. *Current Science*, **75**, 220-227.
- Mutanga, O., and Skidmore, A. K. (2004). Integrating imaging spectroscopy and neural networks to map grass quality in the Kruger National Park, South Africa. *Remote Sensing of Environment*, **90**(1), 104-115.
- Mutanga, O., Skidmore, A. K., and Prins, H. H. T. (2004). Predicting in situ pasture quality in the Kruger National Park, South Africa, using continuum-removed absorption features. *Remote Sensing of Environment*, **89**(3), 393-408.
- Mutanga, O., Skidmore, A. K., and van Wieren, S. (2003). Discriminating tropical grass (*Cenchrus ciliaris*) canopies grown under different nitrogen treatments using spectroradiometry. *ISPRS Journal of Photogrammetry and Remote Sensing*, **57**(4), 263-272.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A., and Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, **403**, 853-858.
- Mygatt, E. (2006). World's forests continue to shrink. Earth policy institute.
- Myneni, R.B., Ross, J., and Asrar, G. (1989). A review on the theory of photon transport in leaf canopies. *Agriculture, Forestry and Meteorology*, **45**, 1-153.
- Nagendra, H. (2001). Using remote sensing to assess biodiversity. *International Journal of Remote Sensing*, **22**, 2377-2400.
- Navalgund, R.R. (2006). Indian earth observation systems: An overview. *Asian Journal of Geoinformatics*, **6**, 17-25.
- Navalgund, R.R., Jayaraman, V., and Roy, P.S. (2007). Remote sensing applications: An overview. *Current Science*, **93**(12), 1747-1766.
- Nemani, R.R., Keeling, C.D., Hashimoto, H., Jolly, W.M., Piper, S.C., Tucker, C.J., Myneni, R.B., and Running, S.W. (2003). Climate-driven increases in global terrestrial net primary productivity from 1982 to 1999. *Science*, **300**, 1560-1563.
- Niemann, K. O. (1995). Remote sensing of forest stand age using airborne spectrometer data. *Photogrammetric Engineering and Remote Sensing*, **61**, 1119-1127.
- Nolin, A.W., and Dozier, J. (1993). Estimating snow grain size using AVIRIS data. *Remote Sensing of Environment*, **44**(2), 231-238.
- O'Neill, A.L., Hardy, S., Fraser, S.J., and McCloy, K.R. (1990). Leaf morphology and the spectral reflectance of some Australian plant species, *In Proceedings of the 5th Australasian Remote Sensing Conference*, Perth, Australia, pp. 1096-1105.
- Ottmar, R.D., Vihnanek, R., and Alvarado, E. (1994). Forest health assessment: Air quality tradeoffs. *In proceedings of the 12th international conference on fire and forest*

- Meteorology*, 26-28 October 1993, Jekyll Island, Georgia. (Society of American Foresters: Bethesda, MD), pp. 47-62.
- Palacios-Orueta, A.P., and Ustin, S.L. (1996). Multivariate Statistical Classification of Soil Spectra. *Remote Sensing of Environment*, **57** (2), 108-118.
- Parker Williams, A., and Hunt, E. R. (2002). Estimation of leafy spurge cover from hyperspectral imagery using mixture tuned matched filtering. *Remote Sensing of Environment*, **82**(2-3), 446-456.
- Parmesan, C., and Yohe, G.A. (2003). Globally coherent fingerprint of climate change impacts across natural systems. *Nature*, **421**, 37-42.
- Pe'rez-Llorens, J.L., Strother, S., and Niell, F.X. (1994). Species differences in short-term pigment levels in four Australian seagrasses in response to dessication and rehydration. *Bot. Mar.* **37**, 91-95.
- Pearlman, J.S., Barry, P.S., Segal, C.C., Shepanski, J., Beiso, D., and Carman, S.L. (2003). Hyperion, a Space-Based Imaging Spectrometer. *IEEE Transaction Geoscience Remote Sensing*, **41**(6), 1150-1173.
- Peñuelas, J., Baret, F., and Filella, I. (1995). Semi-empirical indices to assess carotenoid/chlorophyll a ratio from leaf spectral reflectance. *Photosynthetica* **31**(2), 221-230.
- Peñuelas, J., Field, C. B., Griffin, K., and Gamon, J. A. (1993a). Assessing community type, plant biomass, pigment composition and photosynthetic efficiency of aquatic vegetation from spectral reflectance. *Remote Sensing of Environment*, **46**, 110-118.
- Peñuelas, J., Filella, I., Biel, C., Serrano, L., and Save, R. (1993b). The reflectance at the 950-970 regions as an indicator of plant water status. *International Journal of Remote Sensing*, **14**(10), 1887-1905.
- Peñuelas, J., Pinol, J., Ogaya, R., and Filella, I. (1997). Estimation of plant water concentration by the reflectance Water Index WI (R900/R970). *International Journal of Remote Sensing*, **18**(13), 2869-2875.
- Pereira, V.F.G., Congalton, R.G., and Zarín, D.J. (2002). Spatial and temporal analysis of a tidal floodplain landscape – Aranpi, Brazil – using geographic information systems and remote sensing. *Photogrammetric Engineering and Remote Sensing*, **68**, 463-472.
- Peterson, D.L., Aber, J.D., Matson, P.A., Card, D.H., Swanberg, N., Wessman, C., and Spanner, M. (1988). Remote sensing of forest canopy and leaf biochemical contents. *Remote sensing of Environment*, **24**, 85- 108.
- Peterson, D.L., Spanner, M. A., Running, S.W., and Teuber, K.B. (1987). Relationship of Thematic Mapper Simulator data to leaf area index of temperate coniferous forests. *Remote Sensing of Environment*, **2**, 323-341.
- Pinz, A. (1998). *Tree isolation and species classification*. International Forum Automated Interpretation of High Spatial Resolution Digital Imagery for Forestry, pp. 127-139.
- Plourde, L.C., Ollinger, S.V., Smith, M., and Martin, M.E. (2007). Estimating Species Abundance in a Northern Temperate Forest Using Spectral Mixture Analysis. *Photogrammetric Engineering & Remote Sensing*, **73**(7).
- Poorter, L., Oberbauer, S.F., and Clark, D.B. (1995). Leaf optical-properties along a vertical gradient in a tropical rain forest canopy in Costa-Rica. *American Journal of Botany*, **82**(10), 1257-1263.
- Portigal, F., Holasek, R., Mooradian, G., Owensby, P., Dicksion, M., Fene, M., Elliot, M., Hall, E., and Driggett, D. (1997). Vegetation classification using red-edge first derivative and green peak statistical moment indices with the Advanced Airborne Hyperspectral

- Imaging System (AAHIS). *Third International Airborne Remote Sensing Conference and Exhibition, Copenhagen, Denmark, 7-10 July 1997*, vol. II (Ann Arbor, MI: ERIM), pp. 789-797.
- Price, J.C. (1994). How unique are spectral signatures? *Remote Sensing of Environment*, **49**, 181-186.
- Pu, R., Gong, P., Biging, G.S., and Larrieu, M.R. (2003). Extraction of red edge optical parameters from Hyperion data for estimation of forest leaf area index. *IEEE Transactions on Geoscience and Remote Sensing*, **41**(4), 916-921.
- Puttock, G.D., Timossi, I., and Davis, L.S. (1998). BOREAL: A tactical planning system for forest ecosystem management. *Forestry Chronicle*, **74**(3), 413-420.
- Quigley, T.M., Graham, R.T., and Heynes, R.W. (1996). An integrated scientific assessment for ecosystem management in the interior Columbia River Basin and portions of the Klamath and Great Basins. USDA Forest Service, Pacific Northwest Research Station General Technical Report PNW-GTR- 382, pp. 126.
- Rahman, M.M., Csaplovics, E., Koch, B., and Köhl, M. (2004). Interpretation of Tropical Vegetation Using Landsat ETM+ Imagery. In Altan MO (ed.) *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, **XXXV** (B-YF), Istanbul, pp 157-162.
- Ravindranath, N.H., Somashekhar, B.S., and Gadgil, M. (1997). Carbon flow in Indian forests. *Climate change*, **35**, 297-320.
- Ray, S.S., Das G., Singh, J.P., and Panigrahy, S. (2006). Evaluation of hyperspectral indices for LAI estimation and discrimination of potato crop under different irrigation treatments. *International Journal of Remote Sensing*, **27**(24), 5373-6387.
- Research Systems, Inc., ENVI Version 3.6, 2002, URL: <http://www.RSInc.com/envi>.
- Richards, J.A., and Xiuping, J. (1998). *Remote Sensing Digital Image Analysis*, 3rd edn (Berlin: Springer).
- Roberts, D.A., Belson, B.W., Adams, J.B., and Palmer, F. (1998). Spectral changes with leaf aging in Amazon caatinga. *Trees*, **12**, 315-325.
- Roberts, D.A., Dennison, P.E., Gardner, M.E., Hetzel, Y., Ustin, S.L., and Lee, C.T. (2003). Evaluation of the potential of Hyperion for fire danger assessment by comparison to the Airborne Visible/Infrared Imaging Spectrometer. *IEEE Transactions on Geoscience and Remote Sensing*, **41**(6), 1297-1309.
- Roberts, D.A., Green, R.O., and Adams, J.B. (1997). Temporal and spatial patterns in vegetation and atmospheric properties from AVIRIS. *Remote Sensing of Environment*, **62**, 223-240.
- Roberts, D.A., Numata, I., Holmes, K., Batista, G., Krug, T., and Moteiro, A., (2002). Large area mapping of land-cover change in Rondonia using multitemporal spectral mixture analysis and decision tree classifiers. *Journal of Geophysical Research, D: Atmospheres*, **107**(20), 8073.
- Roberts, D.A., Smith, M.O., and Adams, J.B. (1993). Green vegetation nonphotosynthetic vegetation and soils in AVIRIS data. *Remote Sensing of Environment*, **44**(2), 255-269.
- Roberts, D.A., Ustin, S.L., Ogunjemiyo, S., Greenberg, J., Dobrowski, S. Z., and Chen, J. (2004). Spectral and structural measures of Northwest forest vegetation at leaf to landscape scales. *Ecosystems*, **7**(5), 545-562.

- Rock, B.N., Williams, D. L., Moss, D.M. Lauten, G. N., and Kim, M. (1994). High-Spectral Resolution Field and Laboratory Optical Reflectance Measurements of Red Spruce and Eastern Hemlock Needles and Branches. *Remote Sensing of Environment*, **47**, 176-189.
- Rosso, P.H., Ustin, S.L., and Hastings, A. (2005). Mapping marshland vegetation of San Francisco Bay, California, using hyperspectral data. *International Journal of Remote Sensing*, **26**(23), 5169-5191.
- Roy, P. S. (1993). Remote Sensing for Forest Ecosystem Analysis and Management: in *Environmental studies in India* (ed.) M Balakrishnan (New Delhi: Oxford and IBH), pp. 335-363.
- Running, S.W., Nemani, R., Peterson, D.L., Band, L., Potts, D.F., Pierce, L.L., and Spanner, M.A. (1989). Mapping regional forest evapotranspiration and photosynthesis by coupling satellite data with ecosystem simulation.
- Sabnis, S.D., and Amin, J.C. (1992). Eco-Environmental studies of Sardar Sarovar Environs. The M.S.University of Baroda.
- Sader, S.A., Sever, T., Smoot, J.C., and Richards, M. (1994). Forest change estimates for the northern Peten region of Guatemala-1986 to 1990. *Human Ecology*, **22**(3), 317-332.
- Saksena, A. K., and Tyagi, S. (2002). Management plan for SWS, Rajpipla (East) Division, Forest Department, Wild life wing, Gujarat state, pp. 125-136.
- Salas, W.A., Ducey, M.J., Rignot, E., and Skole, D. (2002). Assessment of JERS-1 SAR for monitoring secondary vegetation in Amazonia: I. Spatial and temporal variability in backscatter across a chrono-sequence of secondary vegetation stands in Rondonia. *International Journal of Remote Sensing*, **23**(7), 1357-1379.
- Salisbury, J.W., Milton, N.M., and Walsh, P.A. (1987). Significance of nonisotropic scattering from vegetation for geobotanical remote sensing. *Remote Sensing of Environment*, **8**, 997-1009.
- Salvatori, R., Grignetti, A., Casacchia, R., and Mandrone, S. (2003). The role of spatial resolution in vegetation studies by hyperspectral airborne images. EPA-Conference on Spectral Remote Sensing of Vegetation, US EPA Environmental Sciences Division, Las Vegas, NV, 12–14 March 2003. Available at: www.epa.gov/esd/land-sci/srsv/images/salvatori-casacchia.pdf (accessed 26 September 2006).
- Sampson, P.H., Treitz, P.M., and Mohammed, G.H. (2001). Remote sensing of forest condition in tolerant hardwoods: An examination of spatial scale, structure and function. *Canadian Journal of remote Sensing*, **27**(3), 232-246.
- Sanchez-Azofeifa, G. A., Castro, K., Rivard, B., Kalascka, M., and Harriss, B. (2003). Remote sensing research priorities in tropical dry forest environments. *Biotropica*, **35**(2), 134-142.
- Sanchez-Azofeifa, G. A., Harris, R. C., and Skole, D. L. (2001). Deforestation in Costa Rica: A quantitative analysis using remote sensing imagery. *Biotropica*, **33**(3), 378-384.
- Sari, M., Sonmez, N.K., and Kurklu, A. (2005). Determination of seasonal variations in solar energy utilization by the leaves of Washington navel orange trees (*Citrus sinensis* L. Osbeck). *International Journal of Remote Sensing*, **26**, 3295-3307.
- Schmidt, K. S., and Skidmore, A. K. (2001). Exploring spectral discrimination of grass species in African rangelands. *International Journal of Remote Sensing*, **22**(17): 3421 - 3434.
- Schmidt, K.S., and Skidmore, A.K. (2003). Spectral discrimination of vegetation types in a coastal wetland. *Remote Sensing of Environment*, **85**, 92-108.
- Scholes, R.J., and Archer, S.R. (1997). Tree grass interactions in savannas. *Annual Review of Ecological Systems*, **28**, 514-544.

- Schriever, J.R., and Congalton, R.G. (1995). Evaluating seasonal variability as an aid to cover-type mapping from Landsat Thematic Mapper data in the Northeast. *Photogrammetric Engineering and Remote Sensing*, **61**(3), 321-327.
- Scurlock, J. M. O., and Prince, S. D. (1993). Remote sensing of biomass and productivity. In D. O. Hall, J. M. O. Scurlock & H. R. Bolhar- Nordenkamp (Ed.), *Photosynthesis and production in a changing environment: A field and laboratory manual*, London: Chapman & Hall, pp. 23-46.
- Sgrenzaroli, M., de Grandi, G.F., and Eva, H et al. (2002). Tropical forest cover monitoring: estimates from the GRFM JERS-1 radar mosaics using wavelet zooming techniques and validation. *International Journal of Remote Sensing*, **7**, 1329-1355.
- Shaw, D. T., Malthus, T. J., and Kupiec, J. A. (1998). High-spectral resolution data for monitoring Scots pine (*Pinus sylvestris* L.) regeneration. *International Journal of Remote Sensing*, **19**, 2601-2608.
- Shaw, G.A., and Bruke, H.K. (2000). Spectral Imaging for remote sensing. *Lincoln laboratory Journal*, **14**(1).
- Shih, T.Y., On the Atmospheric Correction for a Hyperion Scene. *GIS development*, 2004.
- Shippert, P. (1992). Introduction to Hyperspectral Image Analysis. Ph.D Thesis, Geography Department, University of Auckland, New Zealand, pp. 492-504.
- Shippert, P. (2000). Introduction to Hyperspectral Image analysis. Research Systems, www.researchsystems.com.
- Shriner, D.S., and Street, R.B. (1998). North America. In: Watson RT, Zinyowera MC, Moss RH, editors. *The regional impacts of climate change*. New York: Cambridge University Press. pp 253-330.
- Shukla, R.S., and Chandel, P.S. (1989). Plant ecology. S. Chand & Company, New Delhi.
- Silvestri, S., Marani, M., and Marani, A. (2003). Hyperspectral remote sensing of salt marsh vegetation, morphology and soil topography. *Physics and Chemistry of the Earth, Parts A/B/C*, **28**(1-3), 15-25.
- Singh, T.P., Singh, S., Roy, P.S., and Rao, B.S.P. (2002). Vegetation mapping and characterization in west Siang District of Arunachal Pradesh, India – a satellite remote sensing based approach. *Current Science*, **83**, 1221-1230.
- Skole, D., and Tucker, C. (1993). Tropical deforestation and habitat fragmentation in the Amazon-satellite data from 1978 to 1988. *Science*, **260**(5116), 1905-1910.
- Skole, D.L., Chomentowski, W.H., and Salas, W.A. (1994). Physical and human dimensions of deforestation in Amazonia. *Bioscience*, **44**, 314-322.
- Smith, K.L., Steven, M.D., and Colls, J.J. (2004). Use of hyperspectral derivative ratios in the red edge region to identify plant stress responses to gas leak. *Remote Sensing of Environment*, **92**, 207-217.
- Smith, M. O., Ustin, S. L., Adams, J. B., and Gillespie, A. R. (1990). Vegetation in deserts: I. A regional measure of abundance from multispectral images. *Remote Sensing of Environment*, **31**, 1-26.
- Soukupová, J., Rock , B. N., and Albrechtova, J. (2002). Spectral characteristics of lignin and soluble phenolics in the near infrared- a comparative study. *International Journal of Remote Sensing*, **23**(15), 3039-3055.
- Spetich, M.A., Parker, G.R., and Gustafson, E.J. (1997). Spatial and temporal relationships of old-growth and secondary forests in Indiana, USA. *Natural Areas Journal*, **17**(2), 118-130.

- Sphere, G.M. (2005). Application of Phenology to assist in Hyperspectral species classification of a Northern Hardwood Forest. Master's Thesis, Environmental Science Programme, Rochester Institute of Technology, Rochester, New York.
- Srivastava, V.K. (2004). Biodiversity in the Western Ghats districts of Maharashtra and Goa. SAC/RESIPA/FLPG/FED/DOS-DBT/SR/02/2004, Forestry and Environment Division, Space Application Centre, Ahmedabad.
- Steininger, M.K. (1996). Tropical secondary forest regrowth in the Amazon: Age, area and change estimation with Thematic Mapper data. *International Journal of Remote Sensing*, **17**, 9-27.
- Steininger, M.K. (2000). Satellite estimation of tropical secondary forest above-ground biomass: data from Brazil and Bolivia. *International Journal of Remote Sensing*, **21**, 1139-1157.
- Steininger, M.K., Tucker, C.J., Townshend, J.R.G., Killeen, T.J., Desch, A., and Bell, V. (2001). Tropical deforestation in Bolivian Amazon. *Environmental Conservation*, **28**(2), 127-134.
- Streeck, N.A., Rundquist, D. and Connot, J. (2002). Estimating Residual Wheat Dry Matter from Remote Sensing Measurements, *Photogrammetric Engineering and Remote Sensing*, **68**(11): 1193-1201.
- Sudhakar, S., Pujar, G.S., Rajashekhar, G., Gharai, B., and Murthy, M.S.R (2004). Utilization of Resourcesat – I data for advanced forestry applications, *NNRMS Bulletin*, pp. 49-54.
- Thenkabail, P.S. (1999). Characterization of the alternative to slash-and-burn benchmark research area representing the Congolese rainforests of Africa using near-real-time SPOT HRV data. *International Journal of Remote Sensing*, **20**(5), 839-877.
- Thenkabail, P.S. (2002). Optimal Hyperspectral narrowbands for discriminating agricultural crops. *Remote Sensing Reviews*, **20**, 257-291.
- Thenkabail, P.S., Enclona, E.A., Ashton, M.S., Legg, C., and Jean De Dieu, M. (2004). Hyperion, IKONOS, ALI, and ETM+ sensors in the study of African rainforests. *Remote Sensing of Environment*, **90**, 23-43.
- Thenkabail, P.S., Smith, R.B., and De Paul, E. (2000). Hypersepctral vegetation indices and their relationships with agricultural crop characteristics. *Remote Sensing of Environment*, **71**, 158-182.
- Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L. J., and Collingham, Y.C. (2004). Extinct risk from climate change. *Nature*, **427**(6970), 145-148.
- Tiwari, K. P. (1978). A comparative evaluation of landuse and forest type classification and mapping with aerial photograph vis a vis conventional ground stock mapping—a case study in Tehri Garhwal, *Indian Forester*, **104**, 747-767
- Todd, W. S., Hoffer, M. R., and Milchunas, G. D. (1998). Biomass estimation on grazed and ungrazed rangelands using spectral indices. *International Journal of Remote Sensing*, **19**(3), 427-438.
- Tomar, M. S. (1976). Use of aerial photographs in working plans. *Indian Forester*, **102**, 98-10.
- Townshend, J.R.G., Bell, V., and Desch, A. (1995). The NASA Landsat Pathfinder Humid Tropical Deforestation Project. In *Land Satellite Information in the Next Decade*, ASPRS Conference Proceedings, IV-76-IV-87. ASPRS, Virginia.
- Trigg, S.N., Curran, L.M., and McDonald A.K. (2006). Testing the utility of Landsat 7 satellite data for continued monitoring of forest cover in protected areas of Southeast Asia. *Singapore Journal of Tropical geography*, **27**(1), 59-79.
- Tsai, F., and Philpot, W. (1998). Derivative analysis of hyperspectral data. *Remote Sensing of Environment*, **66**, 41–51.

- Tueller, P.T. (1987). Remote sensing science applications in arid environments. *Remote Sensing of Environment*, **23**, 143-154.
- Tuomisto, H., Poulsen, A.D., Ruokolainen, K., Moran, R.C., Quintana, C., and Celi, J. (2003). Linking floristic patterns with soil heterogeneity and satellite imagery in Ecuadorian Amazonia. *Ecological Applications*, **13**(2), 352-371.
- Turner, B.L., Clark, W.C., Kates, R.W., Richards, J.F., Mathews, J.T., and Meyer, W.B. (Eds.), (1990). *The Earth as Transformed by Human Action: Global and Regional Changes in the Biosphere over the Past 300 Years* (Cambridge: Cambridge University Press).
- Ungar, S.G., Pearlman, J.S., Mendenhall, J. A., and Reuter, D. (2003). Overview of the earth observing one (EO-1) mission. *IEEE Transactions on Geoscience and Remote Sensing*, **41**(6), 2003.
- USGS, (2004). EO-1 FAQ, [url://http://eo1.usgs.gov/dataproducts/faq#hyperion](http://eo1.usgs.gov/dataproducts/faq#hyperion).
- Ustin, S.L., and Xiao, Q.F. (2001). Mapping successional boreal forests in interior central Alaska. *International Journal of Remote Sensing*, **22**, 1779-1797.
- Ustin, S.L., Smith, M.O., and Adams, J.B. (1993). Remote sensing of ecological processes: A strategy for developing and testing ecological models using spectral mixture analysis. In J. Ehrlinger and C. Field [eds.], *scaling ecological processes between leaf and landscape*. Academic.
- Ustin, S.L., Wessman, C.A., Curtiss, B., Kasischke, E., Way J., and Vanderbilt, V.C. (1991). Opportunities for using the EOS imaging spectrometers and synthetic aperture radar in ecological models. *Ecology*, **72**, 1934-1945.
- Ustin, S.L., Smith, M.O., Jacquemoud, S., Verstraete, M., and Govaerts, Y. (1999). Geobotany: vegetation mapping in earth science. In *Remote Sensing for the Earth Sciences. Manual of Remote Sensing 3rd Edition*, Vol.3, John Wiley and Sons, Inc., New York, pp. 189-248.
- Vaiphasa, C., Ongsomwang, S., Vaiphasa, T., and Skidmore, A. K. (2005). Tropical mangrove species discrimination using hyperspectral data: a laboratory study. *Estuarine, Coastal and Shelf Science*, **65**, 371-379.
- Van Aardt, J.A.N., and Wynne, R.H. (2001). Spectral separability among six southern tree species. *Photogrammetric Engineering and Remote Sensing*, **67**, 1367-1375.
- Vane, G., and Goetz, A.F.H. (1988). Terrestrial Imaging Spectroscopy. *Remote Sensing of Environment*, **24**, 1-29.
- Vogelmann, T. C. 1993. Plant-tissue optics. *Annual Review of Plant Physiology and Plant Molecular Biology*, **44**, 231-251.
- Volden, E., Solberg, A. S., and Huseby, R.B. (1998). Forest Classification Using Spectrometer and SAR Data. *Proceedings: 1998 Institute of Electrical and Electronics Engineers, Inc., (IEEE) International Geoscience and Remote Sensing Symposium, IGARSS. Part 5 (of 5)*(Seattle, WA, USA), pp. 2732-2736.
- Wessman, C.A., Aber, J.D., and Peterson, D.L. (1989). An evaluation of imaging spectrometry for estimating forest canopy chemistry. *International Journal of Remote Sensing*, **10**, 1293-1316.
- Wessman, C.A., Aber, J.D., Peterson, D.L., and Melillo, J.M. (1988). Remote sensing of canopy chemistry and nitrogen cycling in temperate forest ecosystems. *Nature*, **335**, 154-156.
- Wessman, C.A., Bateson, C.A., and Benning, T.L. (1997). Detecting fire and grazing patterns in tallgrass prairie using spectral mixture analysis. *Ecological Applications*, **7**, 493-511.
- Westman, W.E., and Price, C.V. (1987). Remote detection of air pollution stress to vegetation: laboratory level studies. *Proceedings of the 1987 International Geoscience and Remote*

- Sensing Symposium, Ann Arbor, MI, 18-21 May 1987* (Ann Arbor: University of Michigan), pp. 451-456.
- White, A., Cannell, G.R., and Friend, A.D. (2000). CO₂ stabilization, climate change and the terrestrial carbon sink. *Global Change Biol.*, **6**, 817-833.
- White, J.D., Kroh, G.C., and Pinder, J.E. (1995). Forest mapping at Lassen Volcanic National Park, California, using Landsat TM data and a geographical information system. *Photogrammetric Engineering and Remote Sensing*, **61**, 299-305.
- Whitmore, T.C. (1998). *An introduction to tropical rainforests*. Oxford: Oxford University Press.
- Wilson, B.A., Ow, C.F.Y., Heathcott, M., Milne, D., McCaffrey, T.M., Ghitter, G., and Franklin, S.E. (1994). Landsat MSS classification of fire fuel types in Wood Buffalo National Park, northern Canada. *Global Ecology and Biogeography Letter*, **4**, 33-39.
- Woodcock, C.E., Collins, J.B., Gopal, S. V., Jakabhazy, D., Li, X., Macomber, S., Ryherd, S., Harward, V.J., Levitan, J., and Wu, Y. (1994). Mapping forest vegetation using Landsat TM imagery and a canopy reflectance model, *Remote Sensing of Environment*, **50**(3), 240-254.
- Wright, S.J. (2002). Plant diversity in tropical forests: A review of mechanisms of species coexistence. *Oecologia*, **130**, 1-14.
- Wulder, M.A., White, J.C., Niemann, K.O., and Nelson, T. (2004). Comparison of airborne and satellite high spatial resolution data for the identification of individual trees with local maxima filtering. *International Journal of Remote Sensing*, **25**, 2225-2232.
- Xiao, J., and Moody, A. (2004). Trends in vegetation activity and their climatic correlates: China 1982 to 1998. *International Journal of Remote Sensing*, **25**, 5669-5689.
- Yamano, H., Chen, J., and Tamura, M. (2003). Hyperspectral identification of grassland vegetation in Xilinhot, Inner Mongolia, China. *International Journal of Remote Sensing*, **24**(15), 3171-3178.
- Yu, B., Ostland, M., Gong, P., and Pu, R. (1999). Penalized discriminant analysis of in situ hyperspectral data for conifer species recognition. *IEEE Transaction Geoscience Remote Sensing*, **37**, 2569-2577.
- Zarco-Tejada, P. J., Berjon, A., Lopez-Lozano, R., Miller, J. R., Martin, P., Cachorro, V., Gonzalez, M. R., and de Frutos, A. (2005). Assessing vineyard condition with hyperspectral indices: Leaf and canopy reflectance simulation in a row-structured discontinuous canopy. *Remote Sensing of Environment*, **99**(3), 271-287.
- Zarco-Tejada, P. J., Pushnik, J. C., Dobrowski, S., and Ustin, S. L. (2003). Steady state chlorophyll a fluorescence detection from canopy derivative reflectance and double-peak red-edge effects. *Remote Sensing of Environment*, **84**(2), 283-294.
- Zarco-Tejada, P., and Miller, J. (1999). Land cover mapping at BOREAS using red edge spectral parameters from CASI imagery. *Journal of Geophysical Research*, **104**, 921-933.
- Zhang, J., Rivard, B., Sanchez-Azofeifa, A., and Castro-Esau, K. (2006). Intra-and inter-class spectral variability of tropical tree species at La Selva, Costa Rica: Implications for species identification using HYDICE imagery. *Remote Sensing of Environment*, **105**, 129-141.
- Zhu, Z., and Waller, E. (2003). Global forest cover mapping for the United Nations Food and Agriculture Organization Forest Resources Assessment 2000 programme. *Forest science*, **49**, 369-379.