

List of Figures

Chapter 1: Introduction

Fig. 1.1. Structure of Ubiquitin (PDB:1UBQ). Five anti-parallel β -sheets are shown in blue color, brown color displays the presence of α -helix and a 3_{10} helix.	9
Fig. 1.2. Lysine residues of Ubiquitin (PDB: 1UBQ).	11
Fig. 1.3. Met1-linked linear diubiquitination (PDB: 2W9N).	12
Fig. 1.4. A) Lys-63 linked diubiquitination (PDB: 2JF5), B) Lys-48 linked diubiquitination (PDB: 1AAR).	13
Fig. 1.5. Chains of ubiquitin with different branching patterns and their functions.	14
Fig. 1.6. Substrate protein degradation by ubiquitin proteasome pathway.	16
Fig. 1.7. Reaction scheme of E1 enzyme.	19
Fig. 1.8. A) Comparison of ubiquitin activating enzyme (UBA1) with bacterial ancestor MoeB and ThiF proteins, B) Schematic diagram of sequence relationships between MoeB and UBA1, with conserved sequence motifs indicated by colored blocks. <i>E. coli</i> MoeB and residues 1-250 and 400-660 of human Uba1 are 17% and 22% identical, respectively.	21
Fig. 1.9. Structure of MoeB-MoaD complex (PDB: 1JWA). The N-terminal subdomain of MoeB adopts a variation of the Rossmann fold and is involved in ATP (in Green) binding and catalyzes the adenylation reaction. The C-terminal subdomain contains an antiparallel, four-stranded β - sheet, which makes direct contact with the globular domain of MoaD.	23
Fig. 1.10. Structure of conserved UBC domain of E2 (PDB: 2ESK)	26
Fig. 1.11. HECT and RING E3 ubiquitin ligases.	29

Fig. 1.12. Domain architecture of yeast Rsp5 protein. N-terminus contains a C2 domain followed by several WW domains. The HECT domain is situated on C-terminus, which has catalytic cysteine to form a bond with ubiquitin.	31
Fig. 1.13. A) RING domain, B) U-Box domain.	32
Fig. 1.14. Structure of 26S proteasome.	34

CHAPTER 2: Cloning, Purification and Structural Characterization of Four Domains of Ubiquitin Activating Enzyme E1

Fig. 2.1. Linear representation of yeast ubiquitin activating enzyme E1 and its domains namely, AAD (Active Adenylation Domain), FCCH (First Catalytic Cysteine Half-domain), 4HB (Four Helix Bundle), AAD (Active Adenylation domain), SCCH (Second Catalytic Cysteine Half-domain) and UFD (Ubiquitin Fold Domain).	45
Fig. 2.2. Cartoon diagram of the E1-Ub complex (PDB ID: 3CMM). E1 contains 39% helical (46 helices; 403 residues) 16% β -sheet (47 strands; 170 residues) structure. Domain IAD is in cyan, AAD is in purple, FCCH is in green, SCCH is in blue, UFD is in red and 4HB domain is in pale cyan. Ubiquitin is displayed in yellow and the catalytic cysteine of E1 is in pink.	46
Fig. 2.3. Schematic diagram of the work plan; A) cloning of all four fragments namely; FCCH, 4HB, SCCH and UFD, B) protein purification C) structural analysis of all four peptides by Fluorescence and Circular Dichroism (CD) spectroscopic techniques.	48
Fig. 2.4. Vector map of pET28a.	50
Fig. 2.5. Diagrammatic representation of the cloning strategy. PCR amplified desired fragments encoding domains were double digested with <i>NheI</i> and <i>XhoI</i> enzymes. Vector pET28a was also	

double digested with the same enzymes to form complementary sticky ends for ligation of amplicons to construct the desired clone.	52
Fig. 2.6. Gel picture showing pETRPB1 clone in pET28a backbone.	58
Fig. 2.7. Gel picture showing pETRPB2 clone in pET28a backbone.	59
Fig. 2.8. Gel picture showing pETRPB3 clone in pET28a backbone.	60
Fig. 2.9. Gel picture showing pETRPB4 clone in pET28a backbone.	61
Fig. 2.10. Gel picture showing purification of FCCH peptide.	62
Fig. 2.11. Gel picture showing purification of 4HB peptide.	63
Fig. 2.12. Gel picture showing purification of SCCH peptide.	64
Fig. 2.13. Gel picture showing purification of UFD peptide.	65
Fig. 2.14. Far-UV CD spectra of domains of ubiquitin activating enzyme E1 namely, A) FCCH, B) 4HB, C) SCCH and D) UFD peptides.	68
Fig. 2. 15. Guanidine Hydrochloride denaturation spectra of A) FCCH, B) 4HB, C) SCCH, D) UFD.	71
Fig. 2.16. λ_{\max} values of all four peptides FCCH, 4HB, SCCH and UFD plotted against guanidine hydrochloride concentrations.	72
Fig. 2.17. Fluorescence emission spectra of extrinsic fluorophore ANS bound to the domains of E1 namely; A) FCCH, B) 4HB, C) SCCH, and D) UFD domains in the presence of different concentrations of the denaturant guanidine hydrochloride.	74

CHAPTER 3: Cloning and Functional Characterization of Four Domains of Ubiquitin Activating Enzyme E1

Fig. 3. 1. Schematic diagram of the work plan. A) subcloning of all four DNA fragments encoding FCCH, 4HB, SCCH and UFD into yeast vector, B) Transformation of recombinant plasmids into yeast C) Functional studies of all four peptides by heat stress and antibiotic stress.	80
Fig. 3. 2. Vector map of Yep96 carrying wild type-yeast ubiquitin gene.	82

Fig. 3.3. Gel picture showing YRPB1 clone in Yep96 backbone.	86
Fig. 3.4. Gel picture showing YRPB2 clone in Yep96 backbbone.	87
Fig. 3.5. Gel picture showing YRPB3 clone in Yep96 backbone.	88
Fig. 3. 6. Gel picture showing YRPB4 clone in Yep96 backbone.	89
Fig. 3.7. Growth curves of MHY501 cells expressing domains FCCH, 4HB, SCCH, UFD of ubiquitin activating enzyme E1, A) without inducer B) with CuSO ₄ inducer. Untransformed MHY501 cells were used as negative control.	91
Fig. 3. 8. Survival of MHY501 strain expressing the domains of ubiquitin activating enzyme E1 under heat stress: A) without inducer B) with CuSO ₄ inducer.	93
Fig. 3.9. Effect of overexpression of domains on cell growth under antibiotic stress.	95

Chapter 4: Structural and Functional Studies on β -bulge Mutants of Ubiquitin

Fig. 4.1. Schematic diagram of work plan A) Purification β -bulge mutants Q2N, E64G and S65D, B) Purification ubiquitin activating enzyme E1, C) Incubation of mutant ubiquitin and E1 proteins, D) western blot analysis.	103
Fig. 4.2. Expression and purification of ubiquitin wild type and β -bulge mutants.	109
Fig. 4.3. Expression of E1 protein.	110
Fig. 4. 4. Western blot analysis of ubiquitin activation by E1.	112
Fig. 4.5. CD spectra of Ubiquitin (F45W) and β -bulge mutants. A) Ubiquitin F45W, B) Ubiquitin Q2N, C) Ubiquitin E64G, D) Ubiquitin S65D with each mutant showing effect of different pH point.	114
Fig. 4.6. CD spectra showing effect of different pH range on Ubiquitin (F45W) and β -bulge mutants. A) pH 2.0, B) pH 4.0, C) pH 6.0, D) pH 8.0.	116