

# *Bibliography*

## Bibliography

- Alpi, A.F., Pace, P.E., Babu, M.M., and Patel, K.J. (2008). Mechanistic insight into site-restricted monoubiquitination of FANCD2 by Ube2t, FANCL, and FANCI. *Molecular cell* 32, 767-777.
- Alves-Rodrigues, A., Gregori, L., and Figueiredo-Pereira, M.E. (1998). Ubiquitin, cellular inclusions and their role in neurodegeneration. *Trends Neurosci* 21, 516-520.
- Amerik, A.Y., and Hochstrasser, M. (2004). Mechanism and function of deubiquitinating enzymes. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1695, 189-207.
- Andersen, P.L., Zhou, H., Pastushok, L., Moraes, T., McKenna, S., Ziola, B., Ellison, M.J., Dixit, V.M., and Xiao, W. (2005). Distinct regulation of Ubc13 functions by the two ubiquitin-conjugating enzyme variants Mms2 and Uev1A. *J Cell Biol* 170, 745-755.
- Aravind, L., and Koonin, E.V. (2000). The U box is a modified RING finger - a common domain in ubiquitination. *Curr Biol* 10, R132-134.
- Bartke, T., Pohl, C., Pyrowolakis, G., and Jentsch, S. (2004). Dual role of BRUCE as an antiapoptotic IAP and a chimeric E2/E3 ubiquitin ligase. *Molecular cell* 14, 801-811.
- Batista, P.R., Costa, M.G., Pascutti, P.G., Bisch, P.M., and de Souza, W. (2011). High temperatures enhance cooperative motions between CBM and catalytic domains of a thermostable cellulase: mechanism insights from essential dynamics. *Phys Chem Chem Phys* 13, 13709-13720.
- Bedford, L., Lowe, J., Dick, L.R., Mayer, R.J., and Brownell, J.E. (2011). Ubiquitin-like protein conjugation and the ubiquitin-proteasome system as drug targets. *Nat Rev Drug Discov* 10, 29-46.
- Betts, M. J., & Russell, R. B. (2003). Amino acid properties and consequences of substitutions, In R. B. Michael & C. G. Ian (eds), Bioinformatics for geneticists (pp. 289–316). John Wiley & Sons,Ltd.
- Bosanac, I., Phu, L., Pan, B., Zilberleyb, I., Maurer, B., Dixit, V.M., Hymowitz, S.G., and Kirkpatrick, D.S. (2011). Modulation of K11-linkage formation by variable loop residues within UbcH5A. *J Mol Biol* 408, 420-431.

## Bibliography

- Bucciantini, M., Giannoni, E., Chiti, F., Baroni, F., Formigli, L., Zurdo, J., Taddei, N., Ramponi, G., Dobson, C.M., and Stefani, M. (2002). Inherent toxicity of aggregates implies a common mechanism for protein misfolding diseases. *Nature* *416*, 507-511.
- Buetow, L., and Huang, D.T. (2016). Structural insights into the catalysis and regulation of E3 ubiquitin ligases. *Nature reviews Molecular cell biology* *17*, 626-642.
- Burroughs, A.M., Jaffee, M., Iyer, L.M., and Aravind, L. (2008). Anatomy of the E2 ligase fold: implications for enzymology and evolution of ubiquitin/Ub-like protein conjugation. *J Struct Biol* *162*, 205-218.
- Ceccarelli, D.F., Tang, X., Pelletier, B., Orlicky, S., Xie, W., Plantevin, V., Neculai, D., Chou, Y.C., Ogunjimi, A., Al-Hakim, A., *et al.* (2011). An allosteric inhibitor of the human Cdc34 ubiquitin-conjugating enzyme. *Cell* *145*, 1075-1087.
- Chen, D.C., Yang, B.C., and Kuo, T.T. (1992). One-step transformation of yeast in stationary phase. *Curr Genet* *21*, 83-84.
- Chen, P., Johnson, P., Sommer, T., Jentsch, S., and Hochstrasser, M. (1993). Multiple ubiquitin-conjugating enzymes participate in the in vivo degradation of the yeast MAT alpha 2 repressor. *Cell* *74*, 357-369.
- Chen, Z., and Pickart, C.M. (1990). A 25-kilodalton ubiquitin carrier protein (E2) catalyzes multi-ubiquitin chain synthesis via lysine 48 of ubiquitin. *The Journal of biological chemistry* *265*, 21835-21842.
- Chen, Z.J., Niles, E.G., and Pickart, C.M. (1991). Isolation of a cDNA encoding a mammalian multiubiquitinating enzyme (E2-25K) and overexpression of the functional enzyme in Escherichia coli. *The Journal of biological chemistry* *266*, 15698-15704.
- Chou, P.Y., and Fasman, G.D. (1974). Conformational parameters for amino acids in helical, beta-sheet, and random coil regions calculated from proteins. *Biochemistry* *13*, 211-222.
- Chuang, S.M., and Madura, K. (2005). *Saccharomyces cerevisiae* Ub-conjugating enzyme Ubc4 binds the proteasome in the presence of translationally damaged proteins. *Genetics* *171*, 1477-1484.

## Bibliography

- Cook, B.W., and Shaw, G.S. (2012). Architecture of the catalytic HPN motif is conserved in all E2 conjugating enzymes. *Biochem J 445*, 167-174.
- Cook, W.J., Jeffrey, L.C., Xu, Y., and Chau, V. (1993). Tertiary structures of class I ubiquitin-conjugating enzymes are highly conserved: crystal structure of yeast Ubc4. *Biochemistry 32*, 13809-13817.
- Cook, W.J., Martin, P.D., Edwards, B.F.P., Yamazaki, R.K., and Chau, V. (1997). Crystal Structure of a Class I Ubiquitin Conjugating Enzyme (Ubc7) from *Saccharomyces cerevisiae* at 2.9 Å Resolution. *Biochemistry 36*, 1621-1627.
- Dagert, M., and Ehrlich, S.D. (1979). Prolonged incubation in calcium chloride improves the competence of *Escherichia coli* cells. *Gene 6*, 23-28.
- de Pril, R., Fischer, D.F., Roos, R.A., and van Leeuwen, F.W. (2007). Ubiquitin-conjugating enzyme E2-25K increases aggregate formation and cell death in polyglutamine diseases. *Mol Cell Neurosci 34*, 10-19.
- Deffenbaugh, A.E., Scaglione, K.M., Zhang, L., Moore, J.M., Buranda, T., Sklar, L.A., and Skowyra, D. (2003). Release of ubiquitin-charged Cdc34-S - Ub from the RING domain is essential for ubiquitination of the SCF(Cdc4)-bound substrate Sic1. *Cell 114*, 611-622.
- Dittmar, G., and Winklhofer, K.F. (2020). Linear Ubiquitin Chains: Cellular Functions and Strategies for Detection and Quantification. *Front. Chem. 7:915*.doi: 10.3389/fchem.2019.00915
- Dikic, I., Wakatsuki, S., and Walters, K.J. (2009). Ubiquitin-binding domains - from structures to functions. *Nature reviews Molecular cell biology 10*, 659-671.
- Dunker, A.K., Lawson, J.D., Brown, C.J., Williams, R.M., Romero, P., Oh, J.S., Oldfield, C.J., Campen, A.M., Ratliff, C.M., Hipps, K.W., et al. (2001). Intrinsically disordered protein. *Journal of Molecular Graphics and Modelling 19*, 26-59.
- Eletr, Z.M., Huang, D.T., Duda, D.M., Schulman, B.A., and Kuhlman, B. (2005). E2 conjugating enzymes must disengage from their E1 enzymes before E3-dependent ubiquitin and ubiquitin-like transfer. *Nat Struct Mol Biol 12*, 933-934.
- Finley, D., Ozkaynak, E., and Varshavsky, A. (1987). The yeast polyubiquitin

## Bibliography

gene is essential for resistance to high temperatures, starvation, and other stresses. *Cell* 48, 1035-1046.

- Finley, D., Sadis, S., Monia, B.P., Boucher, P., Ecker, D.J., Crooke, S.T., and Chau, V. (1994). Inhibition of proteolysis and cell cycle progression in a multiubiquitination-deficient yeast mutant. *Mol Cell Biol* 14, 5501-5509.
- Fischer, D.F., De Vos, R.A., Van Dijk, R., De Vrij, F.M., Proper, E.A., Sonnemans, M.A., Verhage, M.C., Sluijs, J.A., Hobo, B., Zouambia, M., *et al.* (2003). Disease-specific accumulation of mutant ubiquitin as a marker for proteasomal dysfunction in the brain. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 17, 2014-2024.
- Gokhale, R.S., and Khosla, C. (2000). Role of linkers in communication between protein modules. *Current opinion in chemical biology* 4, 22-27.
- Gong, M., Wang, H., Chen, M., Bao, D., Zhu, Q., and Tan, Q. (2016). A newly discovered ubiquitin-conjugating enzyme E2 correlated with the cryogenic autolysis of *Volvariella volvacea*. *Gene* 583, 58-63.
- Haas, A.L., Bright, P.M., and Jackson, V.E. (1988). Functional diversity among putative E2 isozymes in the mechanism of ubiquitin-histone ligation. *The Journal of biological chemistry* 263, 13268-13275.
- Haldeman, M.T., Xia, G., Kasperek, E.M., and Pickart, C.M. (1997). Structure and function of ubiquitin conjugating enzyme E2-25K: the tail is a core-dependent activity element. *Biochemistry* 36, 10526-10537.
- Hao, Y., Sekine, K., Kawabata, A., Nakamura, H., Ishioka, T., Ohata, H., Katayama, R., Hashimoto, C., Zhang, X., Noda, T., *et al.* (2004). Apollon ubiquitinates SMAC and caspase-9, and has an essential cytoprotection function. *Nature Cell Biology* 6, 849-860.
- Harper, J.W., and King, R.W. (2011). Stuck in the middle: drugging the ubiquitin system at the E2 step. *Cell* 145, 1007-1009.
- Hawe, A., Sutter, M., and Jiskoot, W. (2008). Extrinsic fluorescent dyes as tools for protein characterization. *Pharm Res* 25, 1487-1499.
- Hershko, A., and Ciechanover, A. (1998). THE UBIQUITIN SYSTEM. *Annual Review of Biochemistry* 67, 425-479.

## Bibliography

- Hilbich, C., Monning, U., Grund, C., Masters, C.L., and Beyreuther, K. (1993). Amyloid-like properties of peptides flanking the epitope of amyloid precursor protein-specific monoclonal antibody 22C11. *The Journal of biological chemistry* 268, 26571-26577.
- Hochstrasser, M. (2002). New Structural Clues to Substrate Specificity in the “Ubiquitin System”. *Molecular cell* 9, 453-454.
- Hochstrasser, M. (2006). Lingering mysteries of ubiquitin-chain assembly. *Cell* 124, 27-34.
- Hodgins, R., Gwozd, C., Arnason, T., Cummings, M., and Ellison, M.J. (1996). The tail of a ubiquitin-conjugating enzyme redirects multi-ubiquitin chain synthesis from the lysine 48-linked configuration to a novel nonlysine-linked form. *The Journal of biological chemistry* 271, 28766-28771.
- Hofmann, R.M., and Pickart, C.M. (1999). Noncanonical MMS2-encoded ubiquitin-conjugating enzyme functions in assembly of novel polyubiquitin chains for DNA repair. *Cell* 96, 645-653.
- Hofmann, R.M., and Pickart, C.M. (2001). In vitro assembly and recognition of Lys-63 polyubiquitin chains. *The Journal of biological chemistry* 276, 27936-27943.
- Hopp, T.P., and Woods, K.R. (1981). Prediction of protein antigenic determinants from amino acid sequences. *Proc Natl Acad Sci U S A* 78, 3824-3828.
- Howard, R.A., Sharma, P., Hajjar, C., Caldwell, K.A., Caldwell, G.A., du Breuil, R., Moore, R., and Boyd, L. (2007). Ubiquitin conjugating enzymes participate in polyglutamine protein aggregation. *BMC Cell Biol* 8, 32.
- Huang, D.T., Hunt, H.W., Zhuang, M., Ohi, M.D., Holton, J.M., and Schulman, B.A. (2007). Basis for a ubiquitin-like protein thioester switch toggling E1-E2 affinity. *Nature* 445, 394-398.
- Jin, L., Williamson, A., Banerjee, S., Philipp, I., and Rape, M. (2008). Mechanism of Ubiquitin-Chain Formation by the Human Anaphase-Promoting Complex. *Cell* 133, 653-665.
- Karathia, H., Vilaprinyo, E., Sorribas, A., and Alves, R. (2011). *Saccharomyces cerevisiae* as a Model Organism: A Comparative Study. *PLoS Studies on Mutant Forms of The Ubiquitin Conjugating Enzymes UBC1 and UBC4*

## Bibliography

ONE 6, e16015.

- Kelley, L.A., Mezulis, S., Yates, C.M., Wass, M.N., and Sternberg, M.J.E. (2015). The Phyre2 web portal for protein modeling, prediction and analysis. *Nature Protocols* 10, 845-858.
- Klempner, N.S., Berleth, E.S., and Pickart, C.M. (1989). A novel, arsenite-sensitive E2 of the ubiquitin pathway: purification and properties. *Biochemistry* 28, 6035-6041.
- Ko, S., Kang, G.B., Song, S.M., Lee, J.G., Shin, D.Y., Yun, J.H., Sheng, Y., Cheong, C., Jeon, Y.H., Jung, Y.K., *et al.* (2010). Structural Basis of E2-25K/UBB+1 Interaction Leading to Proteasome Inhibition and Neurotoxicity. 285, 36070-36080.
- Komander, D., and Rape, M. (2012). The Ubiquitin Code. *Annual Review of Biochemistry* 81, 203-229.
- Lee, I., and Schindelin, H. (2008). Structural insights into E1-catalyzed ubiquitin activation and transfer to conjugating enzymes. *Cell* 134, 268-278.
- Li, X.-J., and Li, S. (2011). Proteasomal dysfunction in aging and Huntington disease. *Neurobiology of Disease* 43, 4-8.
- Lois, L.M., and Lima, C.D. (2005). Structures of the SUMO E1 provide mechanistic insights into SUMO activation and E2 recruitment to E1. *Embo j* 24, 439-451.
- Lu, X., Malley, K.R., Brenner, C.C., Koroleva, O., Korolev, S., and Downes, B.P. (2016). A MUB E2 structure reveals E1 selectivity between cognate ubiquitin E2s in eukaryotes. *Nat Commun* 7, 12580.
- Lv, Z., Williams, K.M., Yuan, L., Atkison, J.H., and Olsen, S.K. (2018). Crystal structure of a human ubiquitin E1-ubiquitin complex reveals conserved functional elements essential for activity. *The Journal of biological chemistry* 293, 18337-18352.
- Mastrandrea, L.D., Kasperek, E.M., Niles, E.G., and Pickart, C.M. (1998). Core domain mutation (S86Y) selectively inactivates polyubiquitin chain synthesis catalyzed by E2-25K. *Biochemistry* 37, 9784-9792.
- Merkley, N., Barber, K.R., and Shaw, G.S. (2005). Ubiquitin manipulation by an E2 conjugating enzyme using a novel covalent intermediate. *The Journal of biological chemistry* 280, 31732-31738.

## Bibliography

- Merkley, N., and Shaw, G.S. (2004). Solution structure of the flexible class II ubiquitin-conjugating enzyme Ubc1 provides insights for polyubiquitin chain assembly. *The Journal of biological chemistry* *279*, 47139-47147.
- Michelle, C., Vourc'h, P., Mignon, L., and Andres, C.R. (2009). What was the set of ubiquitin and ubiquitin-like conjugating enzymes in the eukaryote common ancestor? *J Mol Evol* *68*, 616-628.
- Micsonai, A., Wien, F., Bulyáki, É., Kun, J., Moussong, É., Lee, Y.-H., Goto, Y., Réfrégiers, M., and Kardos, J. (2018). BeStSel: a web server for accurate protein secondary structure prediction and fold recognition from the circular dichroism spectra. *Nucleic acids research* *46*, W315-W322.
- Mishra, P., Prabha, C.R., Rao Ch, M., and Volety, S. (2011). Q2N and S65D substitutions of ubiquitin unravel functional significance of the invariant residues Gln2 and Ser65. *Cell Biochem Biophys* *61*, 619-628.
- Mishra, P., Volety, S., Rao Ch, M., and Prabha, C.R. (2009). Glutamate64 to glycine substitution in G1 beta-bulge of ubiquitin impairs function and stabilizes structure of the protein. *J Biochem* *146*, 563-569.
- Miura, T., Klaus, W., Ross, A., Guntert, P., and Senn, H. (2002). The NMR structure of the class I human ubiquitin-conjugating enzyme 2b. *J Biomol NMR* *22*, 89-92.
- Oddo, S. (2008). The ubiquitin-proteasome system in Alzheimer's disease. *J Cell Mol Med* *12*, 363-373.
- Ohi, M.D., Vander Kooi, C.W., Rosenberg, J.A., Chazin, W.J., and Gould, K.L. (2003). Structural insights into the U-box, a domain associated with multi-ubiquitination. *Nature structural biology* *10*, 250-255.
- Olsen, S.K., Capili, A.D., Lu, X., Tan, D.S., and Lima, C.D. (2010). Active site remodelling accompanies thioester bond formation in the SUMO E1. *Nature* *463*, 906-912.
- Olsen, S.K., and Lima, C.D. (2013). Structure of a ubiquitin E1-E2 complex: insights to E1-E2 thioester transfer. *Molecular cell* *49*, 884-896.
- Oughtred, R., Chatr-aryamontri, A., Breitkreutz, B.J., Chang, C.S., Rust, J.M., Theesfeld, C.L., Heinicke, S., Breitkreutz, A., Chen, D., Hirschman, J., *et al.* (2016). BioGRID: A Resource for Studying Biological Interactions in Yeast. *Cold Spring Harb Protoc* *2016*, pdb.top080754.

## Bibliography

- Oughtred, R., Stark, C., Breitkreutz, B.J., Rust, J., Boucher, L., Chang, C., Kolas, N., O'Donnell, L., Leung, G., McAdam, R., *et al.* (2019). The BioGRID interaction database: 2019 update. *Nucleic Acids Res* *47*, D529-d541.
- Ozkan, E., Yu, H., and Deisenhofer, J. (2005). Mechanistic insight into the allosteric activation of a ubiquitin-conjugating enzyme by RING-type ubiquitin ligases. *Proceedings of the National Academy of Sciences* *102*, 18890-18895.
- Pichler, A., Knipscheer, P., Oberhofer, E., van Dijk, W.J., Korner, R., Olsen, J.V., Jentsch, S., Melchior, F., and Sixma, T.K. (2005). SUMO modification of the ubiquitin-conjugating enzyme E2-25K. *Nat Struct Mol Biol* *12*, 264-269.
- Pires, D.E., Ascher, D.B., and Blundell, T.L. (2014). mCSM: predicting the effects of mutations in proteins using graph-based signatures. *Bioinformatics* *30*, 335-342.
- Pohl, C., and Jentsch, S. (2008). Final stages of cytokinesis and midbody ring formation are controlled by BRUCE. *Cell* *132*, 832-845.
- Porath, J., Carlsson, J.A.N., Olsson, I., and Belfrage, G. (1975). Metal chelate affinity chromatography, a new approach to protein fractionation. *Nature* *258*, 598-599.
- Prabha, C.R., Mukherjee, S., Raman, R., and Kulkarni, S. (2012). The ends and means of artificially induced targeted protein degradation. *Applied Microbiology and Biotechnology* *96*, 1111-1123.
- Raimalani, V., Panchamia, B., and Prabha, C.R. (2019). Construction and Characterization of UBC4 Mutants with Single Residues Swapped from UBC5. *Cell Biochem Biophys*. <https://doi.org/10.1007/s12013-019-00894-1>.
- Rodrigo-Brenni, M.C., and Morgan, D.O. (2007). Sequential E2s drive polyubiquitin chain assembly on APC targets. *Cell* *130*, 127-139.
- Sambrook, J., and Russell, D.W. (2001). Molecular cloning: a laboratory manual (Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press).
- Sammond, D.W., Payne, C.M., Brunecky, R., Himmel, M.E., Crowley, M.F., and Beckham, G.T. (2012). Cellulase Linkers Are Optimized Based on Domain Type and Function: Insights from Sequence Analysis, Biophysical Studies on Mutant Forms of The Ubiquitin Conjugating Enzymes UBC1 and UBC4

## Bibliography

- Measurements, and Molecular Simulation. 7, e48615.
- Scheffner, M., Nuber, U., and Huibregtse, J.M. (1995). Protein ubiquitination involving an E1-E2-E3 enzyme ubiquitin thioester cascade. *Nature* 373, 81-83.
- Schulman, B.A., and Harper, J.W. (2009). Ubiquitin-like protein activation by E1 enzymes: the apex for downstream signalling pathways. *Nature reviews Molecular cell biology* 10, 319-331.
- Schumacher, F.-R., Wilson, G., and Day, C.L. (2013). The N-Terminal Extension of UBE2E Ubiquitin-Conjugating Enzymes Limits Chain Assembly. 425, 4099-4111.
- Seufert, W., and Jentsch, S. (1990). Ubiquitin-conjugating enzymes UBC4 and UBC5 mediate selective degradation of short-lived and abnormal proteins. *EMBO J* 9, 543-550.
- Seufert, W., McGrath, J.P., and Jentsch, S. (1990). UBC1 encodes a novel member of an essential subfamily of yeast ubiquitin-conjugating enzymes involved in protein degradation. *EMBO J* 9, 4535-4541.
- Shabek, N., and Ciechanover, A. (2010). Degradation of ubiquitin: the fate of the cellular reaper. *Cell cycle (Georgetown, Tex)* 9, 523-530.
- Sharma, M., and Prabha, C.R. (2011). Construction and functional characterization of double and triple mutants of parallel beta-bulge of ubiquitin. *Indian J Exp Biol* 49, 919-924.
- Sharma, M., and Prabha, C.R. (2015). Q2N and E64G double mutation of ubiquitin confers a stress sensitive phenotype on *Saccharomyces cerevisiae*. *Indian J Exp Biol* 53, 617-620.
- Sheng, Y., Hong, J.H., Doherty, R., Srikumar, T., Shloush, J., Avvakumov, G.V., Walker, J.R., Xue, S., Neculai, D., Wan, J.W., et al. (2012). A human ubiquitin conjugating enzyme (E2)-HECT E3 ligase structure-function screen. *Mol Cell Proteomics* 11, 329-341.
- Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., et al. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. In *Molecular systems biology*, pp. 539.
- Skaar, J.R., and Pagano, M. (2009). Control of cell growth by the SCF and APC/C ubiquitin ligases. *Curr Opin Cell Biol* 21, 816-824.

## Bibliography

- Sluimer, J., and Distel, B. (2018). Regulating the human HECT E3 ligases. *Cellular and Molecular Life Sciences* 75, 3121-3141.
- Small, D.H., Mok, S.S., and Bornstein, J.C. (2001). Alzheimer's disease and Abeta toxicity: from top to bottom. *Nat Rev Neurosci* 2, 595-598.
- Sonan, G.K., Receveur-Brechot, V., Duez, C., Aghajari, N., Czjzek, M., Haser, R., and Gerday, C. (2007). The linker region plays a key role in the adaptation to cold of the cellulase from an Antarctic bacterium. *Biochem J* 407, 293-302.
- Song, S., Kim, S.Y., Hong, Y.M., Jo, D.G., Lee, J.Y., Shim, S.M., Chung, C.W., Seo, S.J., Yoo, Y.J., Koh, J.Y., *et al.* (2003). Essential role of E2-25K/Hip-2 in mediating amyloid-beta neurotoxicity. *Molecular cell* 12, 553-563.
- Srisodsuk, M., Reinikainen, T., Penttila, M., and Teeri, T.T. (1993). Role of the interdomain linker peptide of *Trichoderma reesei* cellobiohydrolase I in its interaction with crystalline cellulose. *The Journal of biological chemistry* 268, 20756-20761.
- Stewart, M.D., Ritterhoff, T., Klevit, R.E., and Brzovic, P.S. (2016). E2 enzymes: more than just middle men. *Cell Research* 26, 423-440.
- Sung, P., Berleth, E., Pickart, C., Prakash, S., and Prakash, L. (1991). Yeast RAD6 encoded ubiquitin conjugating enzyme mediates protein degradation dependent on the N-end-recognizing E3 enzyme. *The EMBO journal* 10, 2187-2193.
- Szklarczyk, D., Morris, J.H., Cook, H., Kuhn, M., Wyder, S., Simonovic, M., Santos, A., Doncheva, N.T., Roth, A., Bork, P., *et al.* (2017). The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. *Nucleic acids research* 45, D362-D368.
- Tongaonkar, P., Chen, L., Lambertson, D., Ko, B., and Madura, K. (2000). Evidence for an interaction between ubiquitin-conjugating enzymes and the 26S proteasome. *Mol Cell Biol* 20, 4691-4698.
- van Leeuwen, F.W., de Kleijn, D.P., van den Hurk, H.H., Neubauer, A., Sonnemans, M.A., Sluijs, J.A., Koycu, S., Ramdjielal, R.D., Salehi, A., Martens, G.J., *et al.* (1998). Frameshift mutants of beta amyloid precursor protein and ubiquitin-B in Alzheimer's and Down patients. *Science* 279, 242-

## Bibliography

247.

- van Leeuwen, F.W., Hol, E.M., and Fischer, D.F. (2006). Frameshift proteins in Alzheimer's disease and in other conformational disorders: time for the ubiquitin-proteasome system. *J Alzheimers Dis* 9, 319-325.
- van Nocker, S., and Vierstra, R.D. (1993). Multiubiquitin chains linked through lysine 48 are abundant in vivo and are competent intermediates in the ubiquitin proteolytic pathway. *The Journal of biological chemistry* 268, 24766-24773.
- van Tijn, P., de Vrij, F.M., Schuurman, K.G., Dantuma, N.P., Fischer, D.F., van Leeuwen, F.W., and Hol, E.M. (2007). Dose-dependent inhibition of proteasome activity by a mutant ubiquitin associated with neurodegenerative disease. *J Cell Sci* 120, 1615-1623.
- van Wijk, S.J., de Vries, S.J., Kemmeren, P., Huang, A., Boelens, R., Bonvin, A.M., and Timmers, H.T. (2009). A comprehensive framework of E2-RING E3 interactions of the human ubiquitin-proteasome system. *Mol Syst Biol* 5, 295.
- van Wijk, S.J., and Timmers, H.T. (2010). The family of ubiquitin-conjugating enzymes (E2s): deciding between life and death of proteins. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 24, 981-993.
- van Wijk, S.J.L., Melquiond, A.S.J., de Vries, S.J., Timmers, H.T.M., and Bonvin, A.M.J.J. (2012). Dynamic control of selectivity in the ubiquitination pathway revealed by an ASP to GLU substitution in an intra-molecular salt-bridge network. *PLoS Comput Biol* 8, e1002754-e1002754.
- Vandemark, A.P., Hofmann, R.M., Tsui, C., Pickart, C.M., and Wolberger, C. (2001). Molecular Insights into Polyubiquitin Chain Assembly. *Cell* 105, 711-720.
- Verdecia, M.A., Joazeiro, C.A., Wells, N.J., Ferrer, J.L., Bowman, M.E., Hunter, T., and Noel, J.P. (2003). Conformational flexibility underlies ubiquitin ligation mediated by the WWP1 HECT domain E3 ligase. *Molecular cell* 11, 249-259.
- Verma, R., Feldman, R.M., and Deshaies, R.J. (1997). SIC1 is ubiquitinated in vitro by a pathway that requires CDC4, CDC34, and cyclin/CDK activities.

## Bibliography

- Mol Biol Cell 8, 1427-1437.
- Walden, H., Podgorski, M.S., and Schulman, B.A. (2003). Insights into the ubiquitin transfer cascade from the structure of the activating enzyme for NEDD8. *Nature* 422, 330-334.
- Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F.T., de Beer, T.A P., Rempfer, C., Bordoli, L., *et al.* (2018). SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Research* 46, W296-W303.
- Wilson, R.C., Edmondson, S.P., Flatt, J.W., Helms, K., and Twigg, P.D. (2011). The E2-25K ubiquitin-associated (van Leeuwen et al.) domain aids in polyubiquitin chain synthesis and linkage specificity. *Biochem Biophys Res Commun* 405, 662-666.
- Wilson, R.C., Hughes, R.C., Flatt, J.W., Meehan, E.J., Ng, J.D., and Twigg, P.D. (2009). Structure of full-length ubiquitin-conjugating enzyme E2-25K (huntingtin-interacting protein 2). *Acta Crystallogr Sect F Struct Biol Cryst Commun* 65, 440-444.
- Winn, P.J., Religa, T.L., Battey, J.N.D., Banerjee, A., and Wade, R.C. (2004). Determinants of Functionality in the Ubiquitin Conjugating Enzyme Family. *J. Biomol Screen* 12, 1563-1574.
- Worthy lake, D.K., Prakash, S., Prakash, L., and Hill, C.P. (1998). Crystal structure of the *Saccharomyces cerevisiae* ubiquitin-conjugating enzyme Rad6 at 2.6 Å resolution. *The Journal of biological chemistry* 273, 6271-6276.
- Ye, Y., and Rape, M. (2009). Building ubiquitin chains: E2 enzymes at work. *Nature reviews Molecular cell biology* 10, 755-764.
- Yin, Q., Lin, S.C., Lamothe, B., Lu, M., Lo, Y.C., Hura, G., Zheng, L., Rich, R.L., Campos, A.D., Myszka, D.G., *et al.* (2009). E2 interaction and dimerization in the crystal structure of TRAF6. *Nat Struct Mol Biol* 16, 658-666.
- Zhang, M., Windheim, M., Roe, S.M., Peggie, M., Cohen, P., Prodromou, C., and Pearl, L.H. (2005). Chaperoned ubiquitylation--crystal structures of the CHIP U box E3 ubiquitin ligase and a CHIP-Ubc13-Uev1a complex. *Molecular cell* 20, 525-538.
- Zheng, Q., Huang, T., Zhang, L., Zhou, Y., Luo, H., Xu, H., and Wang, X.

## *Bibliography*

(2016). Dysregulation of Ubiquitin-Proteasome System in Neurodegenerative Diseases. *Frontiers in Aging Neuroscience* 8.