## LIST OF SCHEMES AND FIGURES

Figure No.	Title	Page
Scheme 2.1:	Short-course Chemotherapy Treatment for New	37
	Tuberculosis Cases, Adults >50 kg	
Scheme 4.1:	TFH and REV process stages in the preparation of INH	108
	liposomes	
Scheme 4.2:	TFH and REV process stages in the preparation of RFP	112
	liposomes	
Figure 2.1:	Mechanisms of particle deposition in the airways	23
Figure 2.2:	The structure of multilamellar vesicles showing the	41
	organization of phospholipid bilayers and the	
	encapsulation of lipophilic and hydrophilic compounds	
Figure 4.1:	DSC thermograph a) INH; b) EDTA; c) INH: EDTA	126
	complex arrow showing the absence of EDTA peak	
Figure 4.2:	IR spectrum a) INH; b) EDTA; c) INH: EDTA complex	127
Figure 4.3:	NMR spectrum a) INH; b) EDTA; c) INH: EDTA	128
	complex	
Figure 4.4:	Photomicrographs showing liposomal vesicles under	133
	normal light for INH17	
Figure 4.5:	Photomicrographs showing liposomal vesicles under dark	133
	background and polarized lens for RFP17	
Figure 5.1:	Effect of cryoprotectant type/concentration on mean	161
	particle size of rehydrated INH LDP	
Figure 5.2:	Effect of cryoprotectant type/concentration on mean	161
	particle size of rehydrated RFP LDP	
Figure 5.3:	Photomicrograph of liposomal dry powder formulation	163
	under polarized light	
Figure 5.4:	Photomicrograph of liposomal vesicles after hydration of	163
	liposomal dry powder formulation under polarized light	
Figure 5 5:	SEM photomicrographs of INH70	164
Figure 5.6:	SEM photomicrographs of RFP70	164
Figure 5.7:	In vitro deposition of INH and RFP LDPI formulations in	166
	NGI	

Figure 5.8.	INH liposomal dry powder moisture sorption	168
Figure 5.9:	RFP liposomal dry powder moisture sorption	168
Figure 6.1:	DSC thermograph of RFP LDPI formulation	183
Figure 6.2:	Stability profiles - INH70 A) PDR, B) Mean size and C)	185
	FPF vs. time in months	
Figure 6.3:	Stability profiles - INH71 A) PDR, B) Mean size and C)	186
	FPF vs. time in months	
Figure 6.4:	Stability profiles – RFP70 A) PDR, B) Mean size and C)	187
	FPF vs. time in months	
Figure 6.5:	Stability profiles – RFP71 A) PDR, B) Mean size and C)	188
	FPF vs. time in months	
Figure 7.1:	In vitro diffusion profile of INH plain drug and LDPI	199
	formulations	
Figure 7 2:	In vitro diffusion profile of RFP plain drug and LDPI	199
	formulations	
Figure 7.3:	Photomicrograph of alveolar macrophage (A) before and	200
	(B) after liposomal uptake	
Figure 8 1.	Drug level in biological samples following instillation of	209
	INH plain drug and liposomal	
Figure 8.2:	Drug level in biological samples following instillation of	211
	RFP plain drug and liposomal DPI	