

## List of Figures

List of Figures		
Chapter		Page No.
<b>1</b>	<b>Review of Literature and Introduction</b>	
<b>Figure1.1</b>	Molecular determinants of host specificity during nitrogen-fixing symbiosis	3
<b>Figure 1.2</b>	Rhizobia interacting with legumes (Haag et al., 2012). (WT: wild-type).	5
<b>Figure1.3</b>	Nodulation involves the coordinated development of bacterial infection and nodule organogenesis	5
<b>Figure1.4</b>	A model for the growth of rhizobial infection threads (ITs) in root hairs	8
<b>Figure1.5</b>	The different N <sub>2</sub> -fixing organisms and symbioses found in agricultural and terrestrial natural ecosystems	9
<b>Figure 1.6</b>	Mechanism of growth promotion by phosphate solubilizing bacteria	12
<b>Figure 1.7</b>	Schematic diagram of soil phosphorus mobilization and immobilization by bacteria	22
<b>Figure1.8</b>	Simplified cycle of phosphorus in agriculture	25
<b>Figure 1.9</b>	P dynamics in the soil/rhizosphere-plant continuum.	26
<b>Figure 1.10</b>	Pathways and enzymes involved in organic acid biosynthesis by rhizobacteria	27
<b>Figure1.11</b>	Mechanism of P solubilization by phosphate solubilizing bacteria	29
<b>Figure1.12</b>	Master network reaction that was used as the basis for net-flux analysis	31
<b>Figure1.13</b>	Simplified representation of the central carbon metabolism of <i>E. coli</i> .	32
<b>Figure1.14</b>	Carbohydrate metabolism in pseudomonads. Key to the pathway	34
<b>Figure1.15</b>	Glucose Metabolism in <i>Bacillus subtilis</i>	35

<b>Figure 1.16</b>	Pathways of glucose and fructose catabolism available to <i>R. trifolii</i> strain 7000	36
<b>Figure1.17</b>	Outline of Possible pathways of Carbohydrate metabolism in <i>R. meliloti</i>	37
<b>Figure1.18</b>	Reactions of the TCA cycle.	37
<b>Figure 1.19</b>	Possible integration of anaplerotic and bypass pathways with the TCA cycle.	38
<b>Figure1.20</b>	Carbohydrate metabolism in <i>S. meliloti</i>	39
<b>Figure1.21</b>	<i>In vivo</i> carbon flux distribution in <i>S. meliloti</i>	40
<b>Figure 1.22</b>	Metabolic pathways for <i>Rhizobium etli</i>	40
<b>Figure 1.23</b>	Schematic representation of major differences between (a) free living <i>B. japonicum</i> USDA110 and (b) bacteroids	41
<b>Figure 1.24</b>	The enzymes at the PEP–pyruvate–oxaloacetate node in aerobic bacteria.	42
<b>Figure 1.25</b>	The PEP–Pyruvate–OAA node in aerobic <i>E. coli</i> .	46
<b>Figure 1.26</b>	The PEP–pyruvate–OAA node in (a) <i>B. subtilis</i> and (b) <i>C. glutamicum</i> .	47
<b>Figure 1.27</b>	<i>In vivo</i> carbon flux distribution in <i>E. coli</i> , <i>B. subtilis</i> P. <i>fluorescens</i> and <i>S. meliloti</i> .	50
<b>Figure1.28</b>	Design and engineering of pathways for microbial chemical factories (MCFs)	52
<b>Figure1.29</b>	A holistic view of metabolic and cellular engineering	54
<b>Figure 1.30</b>	Model for improved plant nutrient use efficiency with inoculants	61
<b>Figure 1.31</b>	Metabolic basis for designing the genetic modifications in <i>Rhizobium</i> strains	63
<b>Chapter 2</b>	<b>Materials and Methods</b>	<b>Page No.</b>
<b>Figure 2.1</b>	Restriction maps of the plasmids used in this study	69

<b>Chapter 3</b>	<b>Effect of constitutive overexpression of <i>ppc</i> gene of <i>Synechococcus elongatus</i> PCC 6301 on production of organic acid in <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF030669</b>	<b>Page No.</b>
<b>Figure 3.1</b>	Summary of the effects of overexpression of TCA cycle genes in <i>E. coli</i> .	97
<b>Figure 3.2</b>	Glucose metabolism of <i>P. fluorescens</i> and media dependant alterations due to <i>S. elongatus</i> PCC 6301 <i>ppc</i> overexpression.	99
<b>Figure 3.3</b>	Restriction digestion pattern of plasmids containing <i>ppc</i> gene isolated from transformants of <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669 MAFF030669	104
<b>Figure 3.4</b>	MPS phenotype of <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669 MAFF030669 strains harboring pAB3 plasmid.	105
<b>Figure 3.5</b>	Effect of <i>ppc</i> overexpression on extracellular pH and growth	107
<b>Figure 3.6</b>	P solubilization by (A) <i>B. japonicum</i> USDA110 and (B) <i>M. loti</i> MAFF0300669 MAFF030669 transformants on TRP medium.	109
<b>Figure 3.7</b>	Organic acid production from <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669 MAFF030669 <i>ppc</i> gene: A and C : Gluconic, 2-keto gluconic, acetic and citric acids levels and B and D: Organic acid Yields	111
<b>Figure 3.8</b>	Activities of enzymes PPC, PYC, GDH, G-6-PDH, ICDH and ICL in (A) <i>B. japonicum</i> USDA110 and (B) <i>M. loti</i> MAFF0300669 MAFF030669 <i>ppc</i> transformant.	112
<b>Figure 3.9</b>	Key metabolic fluctuations in <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669 MAFF030669 overexpressing <i>ppc</i> gene.	118

<b>Chapter 4</b>	<b>Effect of overexpression of <i>E. coli cs</i> gene on production of organic acid in <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669</b>	<b>Page No.</b>
<b>Figure 4.1</b>	Effect of <i>E. coli gltA</i> gene overexpression on organic acid secretion by <i>P. fluorescens</i> ATCC 13525.	123
<b>Figure 4.2</b>	Activities of key carbon utilization enzymes in <i>P. fluorescens</i> ATCC 13525 overexpressing <i>E. coli gltA</i> gene.	123
<b>Figure 4.3</b>	Activities of key enzymes of carbon utilization in <i>P. fluorescens</i> 13525 co-expressing the <i>ppc</i> and <i>gltA</i> genes.	124
<b>Figure 4.4</b>	Restriction maps of the plasmids used in this chapter .	127
<b>Figure 4.5</b>	Restriction Digestion pattern of plasmids containing <i>cs</i> gene isolated from transformants of <i>B. japonicum</i> USDA110 and <i>M. loti</i> . MAFF030669.	128
<b>Figure 4.6</b>	Restriction digestion pattern of plasmids containing <i>cs</i> gene isolated from transformants of analysis of <i>M. loti</i> MAFF0300669.	129
<b>Figure 4.7</b>	MPS phenotype of (A),(C) <i>B. japonicum</i> USDA110 and (B),(D) <i>M. loti</i> MAFF0300669 harboring pAB7 plasmid expressing <i>E. coli cs</i> gene.	130
<b>Figure 4.8</b>	Effect of <i>cs</i> gene overexpression on extracellular pH and growth profile of (A) <i>B. japonicum</i> USDA110 and (B) <i>M. loti</i> MAFF0300669 .	132
<b>Figure 4.9</b>	P solubilization by (A) <i>B. japonicum</i> USDA110 (B) <i>M. loti</i> MAFF030669 transformants on TRP medium.	135
<b>Figure 4.10</b>	Organic acid production from <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF030669 <i>cs</i> gene (A), (C) organic acids in mM (Gluconic, 2-keto gluconic, acetic and citric acids); (B), (D) Organic acid Yields .	136
<b>Figure 4.11</b>	Activities of enzymes. A- <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF030669 PPC, PYC, GDH, G-6-PDH, ICDH and ICL in and <i>cs</i> transformant. (A): <i>B. japonicum</i> USDA110 and	138

	(B): <i>M. loti</i> MAFF0300669.	
<b>Figure 4.12</b>	Key metabolic fluctuations in <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF030669 overexpressing NADH insensitive <i>E. coli cs</i> gene.	143
<b>Chapter 5</b>	<b>Effect of overexpression of <i>E. coli</i> NADH insensitive Y145F <i>cs</i> gene on production of organic acid in <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF030669</b>	<b>Page No.</b>
<b>152Figure 5.1</b>	<i>E. coli</i> CS protein sequence showing the regulatory variant	146
<b>Figure 5.2</b>	Route of communication between NADH binding sites and active sites in wild type hexameric <i>E. coli</i> CS	146
<b>Figure 5.3</b>	Partial sequence of <i>E. coli</i> NADH insensitive <i>cs</i> gene	150
<b>Figure 5.4</b>	NCBI BLAST analysis of partial <i>cs</i> sequence.	151
<b>Figure 5.5</b>	EBI pair wise alignment of NADH insensitive and wild type <i>cs</i> gene showing the position of mutation.	152
<b>Figure 5.6</b>	Citric acid levels and yields in <i>P. fluorescens</i> PfO-1 wild type and plasmid bearing strains Km, AB7 and YF.	153
<b>Figure 5.7</b>	Key metabolic fluctuations in <i>P. fluorescens</i> PfO-1 overexpressing NADH insensitive <i>E. coli</i> CS.	154
<b>Figure 5.8</b>	Restriction map of the plasmid used in this chapter	156
<b>Figure 5.9</b>	Restriction Digestion pattern of plasmids containing pJNK3 isolated from transformants of (A) <i>B. japonicum</i> USDA110 and (B) <i>M. loti</i> MAFF0300669	157
<b>Figure 5.10:</b>	MPS phenotype of <i>B. japonicum</i> (a),(c) and <i>M. loti</i> MAFF0300669 (b),(d) harboring pJNK3 plasmid expressing <i>cs</i> gene	158
<b>Figure 5.11</b>	Effect of <i>E. coli</i> NADH insensitive <i>cs</i> gene overexpression on extracellular pH and growth profile of (A) <i>B. japonicum</i> USDA110 and (B) <i>M. loti</i> MAFF0300669 on TRP medium	160
<b>Figure 5.12</b>	P solubilization by (A) <i>B. japonicum</i> , (B) <i>M. loti</i>	163

	<i>MAFF0300669</i> <i>MAFF030669</i> transformants on TRP medium.	
<b>Figure 5.13</b>	Organic acid production {(a) and (c)} and Yield {(b) and (d)} from <i>B. japonicum</i> <i>USDA110</i> and <i>M. loti</i> <i>MAFF0300669</i> <i>cs</i> gene transformants, respectively.	164
<b>Figure 5.14</b>	Activities of enzymes PPC, PYC, GDH, G-6-PDH, ICDH and ICL in <i>B. japonicum</i> <i>USDA110</i> and <i>M. loti</i> <i>MAFF0300669</i> <i>ppc</i> transformants.	165
<b>Figure 5.15</b>	Key metabolic fluctuations in <i>B. japonicum</i> <i>USDA110</i> and <i>M. loti</i> <i>MAFF0300669</i> overexpressing NADH insensitive <i>E. coli</i> <i>cs</i> gene.	168
<b>Figure 5.16</b>	Key metabolic fluctuations in <i>Pseudomonas</i> overexpressing NADH insensitive <i>E. coli</i> <i>cs</i> gene.	169
<b>Chapter 6</b>	<b>Effect of overexpression of <i>E. coli</i> NADH insensitive Y145F <i>cs</i> and Na<sup>+</sup> dependant citrate transporter in <i>B. japonicum</i> <i>USDA110</i> and <i>M. loti</i> <i>MAFF030669</i>.</b>	<b>Page No.</b>
<b>Figure 6.1</b>	Basic metabolism and the electron transport chain (ETC) in <i>S. cerevisiae</i> cells	173
<b>Figure 6.2</b>	Oxidation of substrates of plasma membrane dicarboxylate transporter in <i>S. cerevisiae</i>	174
<b>Figure 6.3</b>	Structural model for 2HCT family transporters (Lolkema, 2006).	178
<b>Figure 6.4</b>	Na <sup>+</sup> efflux mechanisms in bacteria. (A). Aerobic and (B). Anaerobic conditions	179
<b>Figure 6.5</b>	Bacterial stress responses and Na <sup>+</sup> homeostasis (Storz et al., 1996).	180
<b>Figure 6.6</b>	Citric acid levels and yields in <i>P. fluorescens</i> PfO-1 overexpressing citrate transporter.	181
<b>Figure 6.7</b>	Activities of enzymes G-6-PDH, ICDH, ICL, PYC, and GDH in <i>P. fluorescens</i> PfO-1 overexpressing citrate transporter.	182

<b>Figure 6.8</b>	Schematic representation of citric acid secretion in <i>Citrobacter</i> sp. DHRSS containing artificial citrate operon (Yadav,2013).	183
<b>Figure 6.9</b>	Organic acid production from <i>H. seropedicae</i> Z67 transformants	184
<b>Figure 6.10</b>	Restriction map of the plasmid used in this chapter (Wagh et al., 2013).	186
<b>Figure 6.11</b>	Restriction digestion analysis of (A) <i>B. japonicum</i> USDA110 containing pJNK4 plasmid. (B) <i>M. loti</i> MAFF0300669 containing pJNK4 plasmid.	187
<b>Figure 6.12</b>	MPS phenotype of <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669 strains harboring pJNK4 plasmid.	188
<b>Figure 6.13</b>	Effect of <i>E. coli</i> NADH insensitive <i>cs</i> gene and citrate transporter <i>citC</i> gene of <i>S. typhimorium</i> gene overexpression on extracellular pH and growth profile of (A) <i>B. japonicum</i> USDA110 and (B) <i>M. loti</i> MAFF0300669, on TRP medium with 50 mM glucose.	190
<b>Figure 6.14:</b>	P release by (A) <i>B. japonicum</i> USDA110 and (B) <i>M. loti</i> MAFF0300669 transformants on TRP medium.	192
<b>Figure 6.15</b>	Organic acid production {(a) and (c)} and Yield {(b) and (d)} from <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669 <i>cs</i> gene transformants, respectively.	194
<b>Figure 6.16</b>	Activities of enzymes PPC, PYC, GDH, G-6-PDH, ICDH and ICL in <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669 <i>ppc</i> transformant.	196
<b>Figure 6.17</b>	Key metabolic fluctuations in <i>B. japonicum</i> USDA110 and <i>M. loti</i> MAFF0300669 overexpressing NADH insensitive <i>E. coli</i> <i>cs</i> gene along with citrate transporter <i>citC</i> gene of <i>S. typhimorium</i> gene	198
<b>Chapter 7</b>	<b>Genomic integration of <i>E. coli</i> NADH insensitive <i>cs</i> along with Na<sup>+</sup> dependent citrate transporter <i>citC</i> gene of</b>	<b>Page No.</b>

	<i>Salmonella typhimurium</i> with <i>vgb</i> , <i>egfp</i> genes in <i>B. japonicum</i> USDA110 <i>M. loti</i> MAFF030669 and <i>S. fredii</i> NGR 234	
<b>Figure 7.1</b>	Strategy used for cloning of artificial citrate operon in pGRG36 containing <i>vhb</i> , <i>egfp</i> resulted in pJIYC.	205
<b>Figure 7.2</b>	PCR amplification of <i>YFcitC</i> with constitutive <i>lac</i> promoter	206
<b>Figure 7.3</b>	Restriction enzyme digestion pattern of pJIYC plasmid containing NADH insensitive Y145F and <i>S. typhimurium</i> sodium citrate transporter operon under <i>lac</i> promoter.	207
<b>Figure 7.4</b>	Confirmation of Genome integrants by PCR amplification of <i>YF-citC</i> from <i>B. japonicum</i> USDA110, <i>M. loti</i> MAFF0300669 MAFF030669 and <i>S. fredii</i> integrants:	207
<b>Figure 7.5</b>	MPS phenotype of <i>B. japonicum</i> USDA110 <i>M. loti</i> MAFF0300669 MAFF030669 and <i>S. fredii</i> integrants	209
<b>Figure 7.6</b>	Extracellular pH and growth profile on glucose 50 mM, Tris-Cl 50 mM rock phosphate medium of <i>B. japonicum</i> USDA110 integrant containing <i>YFcitC</i> , <i>vhb</i> , <i>egfp</i>	211
<b>Figure 7.7</b>	Extracellular pH and growth profile on glucose 50 mM , Tris-Cl 50 mM rock phosphate medium of <i>M. loti</i> MAFF0300669 MAFF030669 integrant containing <i>YFcitC</i> , <i>vgb</i> and <i>egfp</i> .	211
<b>Figure 7.8</b>	Extracellular pH and growth profile on glucose 50 mM , Tris-Cl 50 mM rock phosphate medium of <i>S. fredii</i> integrant containing <i>YFcitC</i> , <i>vgb</i> , <i>egfp</i> .	212
<b>Figure 7.9</b>	P Solubilization by different integrants.	214
<b>Figure 7.10</b>	Organic acid production by three integrants.	215
<b>Figure 7.11</b>	Alterations in enzyme activities in <i>B. japonicum</i> USDA110 integrants	217
<b>Figure 7.12</b>	Alterations in enzyme activities in <i>M. loti</i> MAFF0300669 MAFF030669 integrants.	217

<b>Figure 7.13</b>	Alterations in enzyme activities in <i>S. fredii</i> integrants	217
<b>Chapter 8</b>	<b>Effect of <i>Sinorhizobium fredii</i> NGR 234 genomic integrant containing <i>E. coli</i> NADH insensitive <i>cs</i> along with <i>S. typhimurium</i> <i>citC</i>, <i>vgb</i> and <i>egfp</i> gene cluster on growth promotion of Mung bean (<i>Vigna radiata</i>) plants</b>	<b>Page No.</b>
<b>Figure 8.1</b>	Effect of <i>P. fluorescens</i> genomic integrants on shoot length, weight and P content and root length, weight and P content of mung bean ( <i>Vigna radiata</i> ) at 45 Days after sowing	226
<b>Figure 8.2</b>	Effect of <i>P. fluorescens</i> genomic integrants on number of leaves, nodule number and weight of mung bean ( <i>Vigna radiata</i> ) at 45 Days after sowing	227
<b>Figure 8.3</b>	Effect of <i>P. fluorescens</i> genomic integrants on enzyme activities of mung bean ( <i>Vigna radiata</i> ) at 45 Days after sowing.	227
<b>Figure 8.4</b>	Effect of genomic integrant on bacteria of Rhizospheric soil and bacteroids of nodules.	231
<b>Figure 8.5</b>	Effect of <i>S. fredii</i> NGR 234 genomic integrant on nitrogenase activity of mung bean at 45 Days after sowing	232
<b>Figure 8.6</b>	Effect of <i>S. fredii</i> NGR 234 genomic integrant on shoot length and root length of mung bean at 20 Days after sowing	234
<b>Figure 8.7</b>	Effect of <i>S. fredii</i> NGR 234 genomic integrant on shoot length and root length of mung bean 45 Days after sowing	236
<b>Figure 8.8</b>	Effect of <i>S. fredii</i> NGR 234 genomic integrant on chlorophyll content of mung bean at 45 Days after sowing	236
<b>Figure 8.9</b>	Effect of <i>S. fredii</i> NGR 234 genomic integrant on enzyme activities of mung bean at 45 Days after sowing	242