	L: Drug emux phenomenon in multidrug resistant cells	8
Figure 2.	2: Structure of P-glycoproteins	9
Figure 2.3	3: Nanoparticle bound drug is retained in the cytoplasm	.18
Figure 2.4	4: Schematic illustration of the EPR effect principle. Angiogenesis and enhanced vascular permeability of tumour capillaries and impaired or missing lymphatic clearance of macromolecules result in accumulation of macromolecules (polymers tumour tissue.) in . 22
Figure 3.	1: FTIR spectra of paclitaxel	.33
Figure 3.	2: Regressed calibration curve of PTX in CHCl3: MeOH	.35
Figure 3.	3: Regressed calibration curve of PTX in ACN	.35
Figure 3.	4: Regressed calibration curve of PTX in pH 7.4 PBS containing 20% Ethanol	.36
Figure 3.	5: 1 st derivative regressed calibration curve of PTX in ACN	.38
Figure 3.	6: Regressed calibration curve of PTX in ACN: water	.39
Figure 3.	7: Regressed calibration curve of PTX in rat plasma	.41
Figure 3.	8: Overlay chromatogram of blank plasma and plasma spiked with PTX	.42
Figure 3.	9: Overlay chromatogram of blank liver homogenate and liver homogenate spiked w PTX	ith .44
Figure 3.	10: Overlay chromatogram of blank spleen homogenate and spleen homogenate spiked with PTX	.44
Figure 3.	11: Overlay chromatogram of blank kidney homogenate and kidney homogenate spiked with PTX	.45
Figure 3.	12: Overlay chromatogram of blank lung homogenate and lung homogenate spiked with PTX	. 45
Figure 3.	13: Overlay chromatogram of blank heart homogenate and heart homogenate spike with PTX	d . 46
Figure 3.	14: Overlay chromatogram of blank tumor homogenate and tumor homogenate spil with PTX	ed . 46
Figure 3.	15: Regressed calibration curve of coumarin-6 in ACN	.48
Figure 3.	16: Emission peaks of coumarin-6	.49
Figure 3.	17: Regressed calibration curve of coumarin-6 in ACN	.50
Figure 3.	18: Regressed calibration curve for determination of residual PVA	.52
Figure 3.	19: Regressed calibration curve of Tf by BCA method	.53
Figure 4.	1: Schematic presentation of formation of nanoparticles by nanoprecipitation technique	. 60
Figure 4.	2: Contour plots of effect of PLGA concentration and Organic to aqueous phase volu	me

Figure 4.	3: Contour plot of effect of PVA concentration and Organic to aqueous phase volume ratio on EE(a) -1 level of X ₂ (b) 0 level of X ₂ (c) +1 level of X ₂	69
Figure 4.	4: Contour plots of effect of concentration of lecithin and poloxamer on PS (a) -1 leve of X_2 (b) 0 level of X_2 (c) +1 level of X_2	el 84
Figure 4. !	5: Contour plots of effect of concentration of lecithin and poloxamer on PS (a) -1 leve of X_2 (b) 0 level of X_2 (c) +1 level of X_2	ا 86
Figure 4.	6: NMR spectra of Tf conjugated to PLGA NPs	98
Figure 4.	7: Schematic diagram of conjugation of Tf on PLGA NP surface	98
Figure 4.	8: In-vitro release of PTX in pH 7.4 PBS containing 20% ethanol1	.04
Figure 4.	9: DSC thermograms	.05
Figure 4.	10: TEM images of PLGA NPs (a) unconjugated (b) Tf- conjugated1	.06
Figure 4. :	11: (A) Alkyl cyanoacrylate molecule (B) Anionic polymerization process of alkyl cyanoacryaltes	.08
Figure 4. :	12: Effect of monomer concentration on PS and EE of PBCA NPs prepared by EPand D technique	P .10
Figure 4.	13: Effect of surfactant concentration on PS and EE of PBCA NPs prepared by EPand E technique)P .10
Figure 4.	14: Effect of surfactant type on PS and EE of PBCA NPs prepared by EPand DP technique	.14
Figure 4. :	15: Effect of temperature on PS and EE of PBCA NPs prepared by EPand DP technique	.15
Figure 4.	16: Effect of PTX concentration on PS and EE of PBCA NPs prepared by EPand DP technique	.19
Figure 4.	17: Effect of pH on PS and EE of PBCA NPs 1	.19
Figure 4.	18: Turbidimetric measurement for particle formation by EP and DP technique1	.20
Figure 4.	19: In-vitro release of PTX in pH 7.4 PBS containing 20% ethanol 1	.23
Figure 4.	20: FTIR spectra of (a) BCA monomer and (b) PBCA NPs 1	.24
Figure 4.	21: DSC thermograms of (a) PTX (b) PBCA NPs and (c) PTX loaded PBCA NPs 1	.25
Figure 4.	22: GPC chromatogram 1	.26
Figure 4.	23: NMR spectra of PBCA 1	.27
Figure 4. 2	24: TEM image of PBCA NPs (a) NPs prepared by DP technique and (b) NPs prepared EP technique	by .28
Figure 4.	25: In-vitro release of PTX in pH 7.4 PBS containing 20% ethanol 1	.35
Figure 4.	26: DSC thermograms (a) PTX (b) GTS and (c) PTX GTS SLN	.36
Figure 4.	27: X-ray diffraction pattern of (a) PTX, (b) GTS and (c) PTX GTS SLN	.38
Figure 4.	28: TEM image of PTX GTS SLN 1	.38
Figure 4.	29: Steric stability test of PTX loaded GTS SLN prepared with 1% poloxamer 188 concentration1	.39
Figure 4.	30: In-vitro release of PTX after storage for 6 months at $5^{\circ}C \pm 3^{\circ}C$.45

Figure 4. 31: In-vitro release of PTX after storage for 6 months at $5^{\circ}C \pm 3^{\circ}C$ 145
Figure 4. 32: In-vitro release of PTX after storage for 3 months at 25 \pm 2°C/60 \pm 5% RH146
Figure 4. 33: In-vitro release of PTX after storage for 3 months at $25 \pm 2^{\circ}C/60 \pm 5\%$ RH146
Figure 5. 1: Fluorescent images of intracellular uptake of coumarin-6 loaded nanoparticle formulations after 4hrs incubation161
Figure 5. 2: Uptake efficiency of coumarin-6 loaded PLGA NPs
Figure 5. 3: Uptake efficiency of coumarin-6 loaded Pluronic®P85 coated PLGA NPs163
Figure 5. 4: Uptake efficiency of coumarin-6 loaded Tf PLGA NPs
Figure 5. 5: Uptake efficiency of coumarin-6 loaded Tf PLGA NPs in presence of free Tf164
Figure 5. 6: Uptake efficiency of coumarin-6 loaded PBCA NPs
Figure 5. 7: Uptake efficiency of coumarin-6 loaded GTS SLN165
Figure 5. 8: Cell viability at 24h170
Figure 5. 9: Cell viability at 48h171
Figure 5. 10: Cell viability at 72h172
Figure 6. 1: (A) Normal rat (B) Rat with a subcutaneous C6 glioma
Figure 6. 2: Plasma PTX concentration Vs time profiles for PTX solution and NPs
Figure 6. 3: Liver PTX concentrations after i.v. administration in rats at dose of 20mg/kg184
Figure 6. 4: Spleen PTX concentrations after i.v. administration in rats at dose of 20mg/kg184
Figure 6. 5: Lung PTX concentrations after i.v. administration in rats at dose of 20mg/kg185
Figure 6. 6: Kidney PTX concentrations after i.v. administration in rats at dose of 20mg/kg185
Figure 6. 7: Heart PTX concentrations after i.v. administration in rats at dose of 20mg/kg186
Figure 6. 8: Tumor PTX concentrations after i.v. administration in rats at dose of 20mg/kg186

.

.