

III. LIST OF FIGURES

Figure No.	Title	Page No.
2.1	Structure of the oral mucosa	9
2.2	Time required for the onset of action through oral, dermal and sublingual route	13
2.3	The percent absorption of the drug through oral, dermal and sublingual route	14
2.4	Modified disintegration apparatus	25
2.5	Franz diffusion cell	28
3.1	Structure formula of salbutamol sulphate	53
3.2	UV Spectrum of salbutamol sulphate	57
3.3	Infra-red spectrum of salbutamol sulphate	57
3.4	Mass spectrum of salbutamol sulphate	58
3.5	Structure formula of ondansetron hydrochloride	59
3.6	UV spectrum of ondansetron hydrochloride	63
3.7	Mass spectrum of ondansetron hydrochloride	64
3.8	Structural formula of lamotrigine	65
3.9	UV spectrum of lamotrigine	69
3.10	Infra-red spectrum of lamotrigine	69
3.11	Mass spectrum of lamotrigine	70
4.1	Calibration curve of salbutamol sulphate in distilled water	82
4.2	Calibration curve of salbutamol sulphate in 0.1 N HCl	83
4.3	Calibration curve of salbutamol sulphate in simulated saliva (pH 6.8)	84
4.4	Calibration curve of salbutamol sulphate in simulated gastrointestinal fluid (pH 1.2)	85
4.5	Calibration curve of ondansetron hydrochloride in distilled water	86
4.6	Calibration curve of ondansetron hydrochloride in 0.1 N HCl	87

4.7	Calibration curve of ondansetron hydrochloride in simulated saliva (pH 6.8)	88
4.8	Calibration curve of ondansetron hydrochloride in gastrointestinal fluid (pH 1.2)	89
4.9	Calibration curve of lamotrigine in distilled water	90
4.10	Calibration curve of lamotrigine in 0.1N HCl	91
4.11	Calibration curve of lamotrigine in simulated saliva (pH 6.8)	92
4.12	Calibration curve of lamotrigine gastrointestinal fluid (pH 1.2)	93
4.13	Chemical structure of (A) salbutamol sulphate and (B) internal standard (chloramphenicol)	95
4.14	Representative chromatogram of blank rabbit plasma	99
4.15	Representative chromatogram of plasma spiked with 300 ng/ml salbutamol sulphate (peak 1) and 2000 ng/ml chloramphenicol (peak 2)	99
4.16	Chemical structures of ondansetron hydrochloride and chlorpromazine hydrochloride (Internal standard)	103
4.17	Chromatogram of blank rabbit plasma	107
4.18	Chromatogram of extracted rabbit blood spiked with 100 ng/ml ondansetron hydrochloride (peak 1) and 1000 ng/ml internal standard (peak 2)	107
4.19	Chemical structures of lamotrigine and phenobarbitone (internal standard)	111
4.20	Chromatogram of blank rabbit plasma	115
4.21	Chromatogram of 1000 ng/ml lamotrigine (peak 1) and 10 µg/ml internal standard (peak 2)	115
5.1	Conceptual structure of a biological neuron	127
5.2	A common design of a node in an artificial neural network	128
5.3	A three layered artificial network	131
6.1	Permeation profiles of salbutamol sulphate through porcine buccal mucosa: (□) pH 4.0, (■) pH 6.0, (▲) pH 6.8, (○) pH 7.4, (Δ) pH	165

	8.0, (●) pH 9.0. Each point represents the mean \pm S. D. of five experiments	
6.2	Effect of donor concentration of salbutamol sulphate on steady state flux at pH 7.4. Each point represents the mean \pm S. D. of five experiments	167
6.3	Effect of pH on permeability coefficient and partition coefficient of salbutamol sulphate: (□) Permeability coefficient $\times 10^{-5}$, (▲) Partition coefficient, (●) Fraction of ionized species, (○) Fraction of unionized species. Each point represents the means \pm S. D. of five experiments	169
6.4	Correlation between observed permeability coefficient and calculated permeability coefficient. Each point represents the means \pm S. D. of five experiments	171
6.5	Permeation profiles of ondansetron hydrochloride through porcine buccal mucosa: (□) pH 4.0, (■) pH 6.0, (▲) pH 6.8, (○) pH 7.4, (Δ) pH 8.0, (●) pH 9.0. Each point represents the mean \pm S. D. of five experiments	174
6.6	Effect of donor concentration of ondansetron hydrochloride on steady state flux at pH 7.4. Each point represents the means \pm S. D. of five experiments	176
6.7	Effect of pH on permeability coefficient and partition coefficient of ondansetron hydrochloride: (□) Permeability coefficient $\times 10^{-5}$, (▲) Partition coefficient, (●) Fraction of ionized species, (○) Fraction of unionized species. Each point represents the means \pm S. D. of five experiments	177
6.8	Correlation between observed permeability coefficient and calculated permeability coefficient. Each point represents the means \pm S. D. of five experiments	179
6.9	Permeation profiles of lamotrigine through porcine buccal mucosa. (□) pH 4.0, (■) pH 6.0, (▲) pH 6.8, (○) pH 7.4, (Δ) pH 8.0, (●) pH 9.0. Each point represents the mean \pm S. D. of three experiments	181

6.10	Effect of donor concentration of lamotrigine on steady state flux at ph 7.4. Each point represents the mean \pm S. D. of three experiments	182
6.11	Effect of pH on permeability coefficient and partition coefficient of lamotrigine: (■) permeability coefficient, (●) partition coefficient. Each point represents the mean \pm S. D. of three experiments	184
6.12	Correlation of 1-octanol/buffer partition coefficient and permeability coefficient	185
6.13	Correlation between observed permeability coefficient and calculated permeability coefficient	186
6.14	Relationship between permeability coefficient and the fraction of different species of lamotrigine: (■) fraction of ionized species, (○) permeability coefficient, (▲) fraction of unionized species. Each point represents the mean \pm S. D. of three experiments	187
6.15	Relationship between partition coefficient and the fraction of different species of lamotrigine: (■) fraction of ionized species, (○) partition coefficient, (▲) fraction of unionized species. Each point represents the mean \pm S. D. of three experiments	188
6.16	Content of salbutamol sulphate (μ g) in mucosa after run at various pH. Each point represents the means \pm S. D. of five experiments	191
6.17	Content of ondansetron hydrochloride (μ g) in mucosa after run at various pH. . Each point represents the means \pm S. D. of five experiments	192
6.18	Content of lamotrigine (μ g) in mucosa after run at various pH. . Each point represents the means \pm S. D. of five experiments	193
7.1	PAMPA Log $P_c^{7,4}$ vs. Human Absorption Values !	201
7.2	UV spectra of the equilibrium samples in the 96-well plate	207
7.3	UV spectra of the samples incubated for 6 hrs and 24 hrs in the 96-well plate	212
8.1	Contour plot for tensile strength (----), % drug release at 2 minutes (.....) and over all desirability (—) keeping amount of	235

	mannitol at 0 level	
8.2	Contour plot for tensile strength (----), % drug release at 2 minutes (.....) and over all desirability (—) keeping amount of poly vinyl alcohol at 0 level	235
8.3	Contour plot for tensile strength (----), % drug release at 2 minutes (.....) and over all desirability (—) keeping amount of glycerol at 0 level	236
8.4	SEM micrograph of salbutamol sulphate film at 2000X magnification	239
8.5	SEM micrograph of salbutamol sulphate film at 4500X magnification	239
8.6	DSC of thermograms of salbutamol sulphate optimized film	240
8.7	Comparative dissolution profiles of salbutamol sulphate film in distilled water (▲), simulated saliva (pH 6.8) (■) and simulated gastric fluid (pH 1.2) (●)	241
8.8	Dissolution of the film of salbutamol sulphate in distilled water in beaker (A) film outside the beaker, (B) film in distilled water (C) film in distilled water after 30 seconds, (D) film distilled water after 60 seconds	242
8.9	Dissolution of the film of salbutamol sulphate in guinea pig buccal mucosa (A) buccal mucosa without film (B) film put over the buccal mucosa (C) film over the buccal mucosa after 30 seconds, (D) film over the buccal mucosa after 60 seconds	243
8.10	SEM micrographs of ondansetron hydrochloride film at (A) 1000X (B) 2000X magnification	246
8.11	DSC of thermograms ondansetron hydrochloride film	247
8.12	Comparative dissolution profiles of ondansetron hydrochloride in distilled water (▲), simulated saliva (pH 6.8) (■) and simulated gastric fluid (pH 1.2) (●)	249
8.13	SEM micrograph of lamotrigine film at (A)1000X and (B) 2000X	251

	magnification	
8.14	DSC of thermograms lamotrigine film	252
8.15	Comparative dissolution profiles of lamotrigine film in distilled water (▲), simulated saliva (pH 6.8) (■) and simulated gastric fluid (pH 1.2) (●)	254
9.1	Architecture of three-layer neural network	263
9.2	Phase solubility diagram of the lamotrigine- β -cyclodextrin system	272
9.3	Optimization of neural network: (A) optimization of step of hidden layer, and (B) optimization of step size of output layer	273
9.4	DSC thermograms of lamotrigine, β -cyclodextrin, physical mixture (LM: β -CD: 1:1), coprecipitation solid dispersions (LM: β -CD:transcutanol: 1:1:0.03)and cogrinding solid dispersions (LM: β -CD:transcutanol: 1:1:0.03)	276
9.5	Powder x- ray diffraction patterns of lamotrigine, β -cyclodextrin, physical mixture (LM: β -CD: 1:1), coprecipitation solid dispersions (LM: β -CD:transcutanol: 1:1:0.03)and cogrinding solid dispersions (LM: β -CD:transcutanol: 1:1:0.03)	277
9.6	Scanning electron photomicrographs of (a) lamotrigine, (b) β -cyclodextrin, (c) physical mixture (LM: β -CD: 1:1), (d) solid dispersions produced by coprecipitation method (LM: β -CD:transcutanol: 1:1:0.03), (e) solid dispersions produced by cogrinding method (LM: β -CD:transcutanol: 1:1:0.03)	279
9.7	(a) Dissolution of lamotrigine from solid dispersions. (b) Dissolution of lamotrigine from solid dispersions (LM: β -CD:transcutanol: 1:1:0.03) stored at room temperatures for 3 month. (c) Dissolution of lamotrigine from solid dispersions (LM: β -CD:transcutanol: 1:1:0.03) stored at 45 °C for 3 month	280
9.8	Percent lamotrigine dissolved from tablets made from the different solid dispersion versus time	281
10.1	Contour plot of tensile strength obtained from simplex lattice design	298

10.2	Contour plot of disintegration time obtained from simplex lattice design	299
10.3	Superimposed contour plots of tensile strength and disintegration time for optimum region	301
10.4	Comparative dissolution profile of tablets of ondansetron hydrochloride in distilled water (▲) and simulated saliva (pH 6.8) (■)	306
10.5	Disintegration of the fast dissolving tablet of ondansetron hydrochloride in distilled water in glass. (A) tablet put into the glass, (B) tablet at 10 seconds, (C) tablet after 15 seconds, (D) tablet after 20 seconds	307
10.6	Comparative dissolution profile of tablets of salbutamol sulphate in distilled water (▲) and simulated saliva (pH 6.8) (■)	313
10.7	Comparative dissolution profile of lamotrigine tablets in distilled water (▲) and simulated saliva (pH 6.8) (■)	319
11.1	Comparison of pharmacokinetic profiles of different formulations of sabutamol sulphate. (○) Conventional tablet, (■) fast dissolving tablet, (▲) fast dissolving film	335
11.2	Comparison of pharmacokinetic profiles of different formulations of ondansetron hydrochloride. (○) Conventional tablet, (■) fast dissolving tablet, (▲) fast dissolving film	338
11.3	Comparison of pharmacokinetic profiles of different formulations of lamotrigine. (○) Conventional tablet, (■) fast dissolving tablet, (▲) fast dissolving film	341