

Bibliography



Abott, R.T., 1954. *American Sea shells*. Dvan Nostrand Company Inc. Newyork. 541pp.

Achuthankutty, C.T., Nair, S.R.S., Devassy, V.P and Nair, V.R., 1981. Plankton composition in two estuaries of Konkan coast during pre monsoon season. *Mahasagar*. 14(1):55-60.

AjmalKhan, S., Murugesan, P., Lyla P.S. and Jaganathan, S., 2004. A new indicator macro invertebrate of pollution and utility of graphical tools and diversity indices in pollution monitoring studies. *Curr. Sci.* 87 (11): 1508-1510.

Alkawri, A.A.S. and Ramaiah, N., 2010. Spatio-temporal variability of dinoflagellate assemblages in different salinity regimes in the west coast of India. *Harmful Algae.* 9 (2): 153-162

Alongi, D.M., 1990. The ecology of soft-bottom benthic ecosystems, *Oceanogr Marine Biol. Annl. Review.* 28: 381–496.

Ananthan, G., 1995. Plankton ecology and heavy metal studies in the marine environments of Pondicherry. Ph.D. Thesis, Annamalai University, India. pp 225.

Ansari, Z.A. and Parulekar, A.H., 2001. Benthos, In: *The Indian Ocean: A perspective* (Ed) SenGupta, R. and Ehrlich D. 2: 451-466.

Ansari, Z.A., Ingole, B.S., Banerjee, G. and Parulekar, A.H., 1986. Spatial and temporal changes in benthic macrofauna from Mandovi and Zuari estuaries of Goa, West coast of India. *Ind. J. Marine Science*. 15: 223–229.

Anthony, K.R.N. and Fabricius, K.E., 2000. Shifting roles of heterotrophy and autotrophy in coral energetic under varying turbidity. *J. Exp. Mar. Biol. Ecol.* 252: 221-253.

APHA, 1998. Standard Method for the Examination of Water and Seawater American Public Health Association, New York. pp 769.

Apte, D., 1998. *The book of Indian shells*. Bombay Natural History Society, Oxford University press, Delhi. pp 115.

Araujo, H M P. Nascimento-Vieira, D., Neumann-Leitão, S., Schwamborn, R., Lucas, A P O. Alves, J P H. (2008) ;Zooplankton community dynamics in relation to the seasonal cycle and nutrient inputs in an urban tropical estuary in Brazil, *Braz. J. Biol.* 68(4) São Carlos Nov. 2008

Banse, K. and Mcclain, C.R., 1986. Winter blooms of phytoplankton in the Arabian sea as observed by Coastal Zone color scanner. *Marine ecology:Progress series*. 34:201-211.

Bhattathiri, P.M.A., 1992. Primary production of tropical marine ecosystems. *Tropical ecosystems: Ecology and Management*, NIO, Goa. pp 269-276.

Bhattathiri, P.M.A.Pant, A.Sawant, S.S.Gauns, M.Matondkar, S.G.P. Mahanraju, R.(1996); Phytoplankton production and chlorophyll distribution in the eastarn and central Arabian Sea in 1994-1995; *Current science*, 71(.11) 10 December 1996

Bhunia, A.B. and Choudhury, A., 1998. Studies on the seasonal abundance and biomass of Crustacean Zooplankton and Chaetognaths in relation to ecological parameter of a tidal creek (Mooriganga), of sagar Island (north), Sunderbans, West Bengal. *Ind. J. Mar. Sci.* 28: 193-198.

Bloom, S.A., Simon, J.L. and Hunter, V.D., 1972. Animal sediment relations and community analysis of Florida estuaries. *Mar. Biol.* 13: 43-56.

Boesch, D.F., Field, J.C. and Scavia, D. (ed.), 2000. The potential consequences of climate variability and change on coastal areas and marine resources: Report of the coastal areas and marine resources sector team, *Coastal ocean program decision analysis series* 21, U.S. Global Change Research Program, National Oceanic and Atmospheric Administration, Silver Springs, Maryland.

Bouillon, S., Koedam, N., Raman, A.V. and Dehairs, F., 2002. Primary producers sustaining macro-invertebrates communities in intertidal mangrove forest. *ecologia*. 130: 441-448.

Bray, J.R. and Curtis, J.T., 1957. An introduction of the upland forest communities of southern Wisconsin. *Ecol. Monogr.* 27: 325-349.

Chandran, R. and Ramamoorthi, K., 1984. Hydrological studies in the gradient zone of the velar estuary, I: Physicochemical parameters. *Mahasagar–Bull.Natn.Inst. Oceanography.* 17(2): 69-77.

Chandran, R. and Ramamurthi, K., 1984. Hydrobiological studies in the gradient zone of the velar estuary-nutrients. *Mahasagar Bull.Natl.Inst.Oceanogr*. 17:133-140.

Chandran, R., 1982. Hydrobiological studies in the gradient zone of the velar estuary. Ph.D. Thesis, Annamalai University, India. pp 198.

Chandran, R., 1987. Hydrobiological studies in the gradient zone of the velar estuary, IV, Benthic fauna. *Mahasagar Bull. Nat. Inst. Oceanography.* 20(1): 1–13.

Chandran, R., 1987. Hydrological studies in the gradient zone of the vellar estuary benthic fauna. *Mahasagar*. 24(2): 81-88.

Chapgar, B.F., 1957. Marine crabs of Bombay state. Taraporevala Marine Biological Station, Bombay. pp 88.

Chapman A.J. and Underwood, M.G. 1998. Spatial Analyses of Intertidal assemblages on sheltered rocky shores. Aust. J. Ecol. 23: 138-157.

Chavez, F.P., Barber, R.T., 1987. An estimate of new production in the equatorial pacific. Deep-Sea Re. 34:1229-4.

Clarke, K.R. and Green, R.H., 1988. Statistical design and analysis for a biological effects study. *Mar. Ecol. Prog. Ser.* 46: 316-226.

Clarke, K.R. and Warwick, R.M. 1994. Similarity-based testing for community pattern: The two-way layout with no replication. *Mar Biol.* 118 (1): 167 - 176.

Coastal Ocean Monitoring and Prediction System (COMAPS), 1999. Annual Report, Central Salt & Marine Chemicals Research Institute, Bhavnagar. pp 208.

Damodara, N.W., Santhanam, R., Krishnamurthy, K. and Natarajan, R., 1997. The species biomass and the seasonal composition of Tintinida (Protozoa: Ciliata). *Proc. Sym. Warm Water. Zoopl.* (Spl. Publ. NIO/UNESCO). pp 520-527.

Damotharan, P., Vengadesh Perumal, N, Arumugam, M (2010); Studies on zooplankton ecology from Kodaikkarai (Point Calimere) Coastal Waters (South East Coast of India); Research Journal of Biological Sciences 5(2): 187-198, 2010.

Das, A.K., 1995. Free living ciliates (Protozoa) of Chilika Lake, Orissa. Wetland Ecosystem Series I.

Davassy, V.P. and Goes, J.I., 1989. Seasonal patterns of phytoplankton biomass and productivity in a typical estuarine complex. *Proc Indian Acad. Sci.* 99(55): 485-501.

Dave, C.S. (2011) Ecological assessment of Narara reef with special reference to coral community. Ph.D. thesis submitted to The M. S. University of Baroda, Vadodara, India. 139 p.

De Souza, S.N. and Gupta, R.S., 1986. Variations of dissolved oxygen in Mandovi and Zuari estuaries. *Ind. J. Marine Science*. 15: 67–71.

Desa, E., Zingde, M.D., Vethamony, P., Babu, M.T., DeSousa, S.N. and Verlecar, X.N., 2005. Dissolved oxygen - A target indicator in determining use of the Gulf of Kachchh waters. *Marine pollut. bul.* 50: 73-79.

Devassy, V.P. and Goes J.I. 1988; Phytoplankton community structure and succession in a tropical estuarine complex (central west coast of India) Estuarine, *Coastal and Shelf Science* 27(6), 671-685.

Devassy, V.P., 1983. Plankton ecology of some estuarine and marine regions of the west coast of India, Ph.D. Thesis, University of Kerala.

Devi, L.P., 1994. Ecology of Coleroon estuary: Studies on benthic fauna. *J. Marine Biol. Assoc. India.* 36(1&2): 260–266.

Dhawan, R.M., 1970. Plankton and hydrological factors at Kandla in the Gulf of Kutch during 1960-1963. *Indian J. Fish.* 17: 122-131.

Dickey-Collas M, Gowen R J and Fox C J (1996); Distribution of Larval and Juvenile Fish in the Western Irish Sea: Relationship to Phytoplankton, Zooplankton Biomass and Reccurrent Physical Features *Marine and Freshwater Research* 47(2) 169 – 181.

Dolganova, T.Y. and Zuenko, I., 2001. Seasonal and interannual dynamic of mesoplankton in northwestern Japan. *Marine Ecol. Prog. Series*. 215: 213-224.

Duarte, P., Macedo, M.F. and Fonseca, L.C.D., 2006. The relationship between phytoplankton diversity and community function in a coastal lagoon. *Hydrobiologia*. 555: 3-18.

Edinger, E.N. and Risk, M.J., 2000. Reef classification by coral morphology predicts coral reef conservation value. *Biol. Conser.* 92: 1-13.

Ekwu, A.O., Sikoki, F.D., 2006. Phytoplankton Diversity in the Cross River Estuary of Nigeria. *Appl. Sci. Environ. Mgt.* 10 (1): 89-95.

Falkowski, P.G., 1994. The role of Phytoplankton Photosynthesis in Global Biogeochemical Cycles. *Photosynthesis and research*. 3:39.

Gaonkar CA · Krishnamurthy V · Arga C A (2010) Changes in the abundance and composition of zooplankton from the ports of Mumbai, India; *Environ Monit Assess* (2010) 168:179–194

Gerlach, S.A., 1978. Food chain relationships in subtidal silty sand marine sediments and the role of meiofauna in stimulating bacterial productivity. *Oecologia*. 33: 55-69.

Gibson, R.N., 1982. Recent studies on the biology of intertidal fishes. *Oceano. Marine Biol. Annual Review.* 20: 363–414.

Giere, O., 1993. Meiobenthology: *The microscopic fauna in aquatic sediments*. Springer Verlag, New York. pp 1 – 328.

Golterman, H.L., Clymo, R.S. and Ohnstad, M.A.M., 1978. *Methods for Physical and Chemical Analysis of Freshwaters* (Sec. Ed.) Blackwell Scientific Publications, Oxford. pp 213.

Gopalakrishnan, P., 1972. Studies on marine plankton diatoms off Port Okha in Gulf of kachchh. *Phykos*. 11 (1-2):37-49.

Goswami, S.C and Padmavathi, G., 1996. Zooplankton production, composition and diversity in the coastal water of Goa. *Indian J. Mar. Sci.* 25: 91-97.

Gray, J.S., Clarke, K.R., Warwick, R.M. and Hobbs, J., 1990. Detection of initial effect of pollution on marine benthos: An example from the Ekofish and Eldfish oilfields, North Sea. *Mar. Ecol. Prog. Ser.* 66: 285-299.

Gray, J.S., McIntyre, A.D. and Stirn, J., 1992. *Manual of Methods in aquatic environmental research*. Part II. Biological assessment of marine pollution with particular reference to benthos. FAO Fisheries Technical Paper. 324: 49.

Gray, J.S., Wu, R.S. and Or, Y.Y., 2002. Effects of hypoxia and organic enrichment on the coastal environment. *Mar.Ecol.Prog.Ser.* 238: 249-279.

Gujarat Institute of Desert Ecology (GUIDE), 2005. Marine Monitoring of Kharo creek. Report submitted to Sanghi Industries Ltd.

Harvey, H.W., 1960. The chemistry and fertility of seawater. Cambridge University press, London.

Holme, N.A., 1964. Methods of the sampling in benthos. *Advan. Marine. Biol.* 2: 171-260.

ICMAM, 2002. Geographical Information System for Gulf of Kachchh. Integrated Coastal and Marine Area Management Directorate, Chennai, Department of Ocean Development, Government of India.

ICMAM, 2004. Model Integrated Coastal and marine area management plan for Gulf of Kachchh. Department of Ocean Development, Ministry of Earth Sciences.

ICMAM Project Directorate, Chennai, Government of India.

Jayalakshmy, K.V., Kumaran, S., Vijayan, M., 1986. Phytoplankton distribution in Cochin backwaters - a seasonal study. *Mahasagar*. 19:29-37.

Jayatissa, L.P., Dahdouh-Guebas, F. and Koedam, N., 2002. A review of the floral composition and distribution of mangroves in Srilanka. *Bot. J. Linn. Soc.* 138: 29-43.

Jean Jose J., Udayakumar, P., Chandran, A., Narendra Babu, K., Sudhanandh V.S.; Zooplankton Diversity in Vallarpadam, India: Influence of Hydrochemistry, Season and Semi Diel Cycle; Asian Journal of Water, Environment and Pollution 103-108.

Jones, D., 1984. Crabs of the mangal ecosystem, *In*: Por, F., and Dor, I. (ed.). *Hydrobiology of the Mangal*, Junk Publishers, The Hague. pp.89-109.

Jyothibabu, R., Madhav, N.V., Maheswaran, P.A., Nair, K.C.C., Venugopal, P. and Balasubramaniam, T., 2003. Dominance of dianoflagellates in microzooplankton community in oceanic regions of the Bay of Bengal and the Andaman Sea. *Curr. Sci.* 84:9.

Kalyani, N., 1988. The Meiofauna in and around the Pulicat Lake. Ph.D. Thesis, University of Madras. pp198.

Kannan, R., 1992. Microphytoplankton of the Pichavaram mangroves, southeast coast of India, Species composition and population density. *Hydrobiol.* 247: 77-86.

Kardani, H.K., Thivakaran, G.A., Mankodi, P.C. (2011) Community structure of bivalve in intertidal region of Gulf of Kachchh *Electronic Journal of Environmental Sciences* Vol. 4, 129-134 (2011)

Kathiresan, K. and Bingham, B.L., 2001. Biology of Mangroves and mangrove ecosystem. *Adv. Mar. Bio.* 40:81-251.

Khuantrairong, T. and Trichaiyaporn, S., 2008. Diversity and Seasonal Succession of the Phytoplankton Community in Doi Tao Lake, Chiang Mai Province, Northern Thailand. *Nat. Hist. J. Chulalongkorn Univer.* 8: (2):143-156.

Kinne, O., 1963. The effects of temperature and salinity or marine and brackish water animals. *Oceanogr. Mar. Biol. Ann. Rev.* 1: 301-340.

Krishnamoorthy, P. and Subramanian, P., 1999. Organization of commercially supported Meroplankton in Palk bay and Gulf of Mannar Biosphere reserve area, Southeast coast of India. *Indian J. Mar. Sci.* 28:211-215.

Krishnamurthy, K. and Santhanam, R., 1975. Ecology of Tintinids (Protozoa: Ciliata) in Porto Novo region. *Indian J. Mar. Sci.*, 4: 181-184.

Krishnamurthy, K.N., 1971. Preliminary studies on the bottom biota of Pulicat Lake. J. Mar. Boil. Ass. India. 13(2): 264-269.

Kristensen, R.M., 1983. Loricifera. A new phylum with Aschelminthes characters from the meiobenthos. Z. Zool.Syst. Evolutionsforsch. 21: 161-180.

Kruskal, J.B., 1964. Multidimensional scaling by optimising goodness of fit to a non-metric hypothesis. *Pscyhometrika*. 29: 1-27.

Kruskal, J.B., 1964. Nonmetric multidimensional scaling: a numerical method, *Psychometrika*. 29: 115-129.

Kumar, K., 1993. Studies on Copepod occurring in coastal waters of Parangipettai Ph.D. thesis, Annamalai University, India. pp 166.

Kumar, P.S., and Prasad, T.G., 1996. Winter cooling in the northern Arabian Sea. *Curr. Sci.* 71(11): 834–841.

Kumar, R.S. and Antony, A., 1994. Preliminary studies on the polychaete fauna of the mangrove areas of Cochin. *In* Kerala, T. and Ravikumar R. (Ed.) Proceedings of the 6th Kerala science congress (pp 74–77). Thiruvananthapuram: State Committee on Science, Technology and Environment.

Kumar, R.S., 1995. Macro benthos in the mangrove ecosystem of Cochin backwaters, Kerala (Southwest coast of India). *Ind. J. Marine Sciences*. 24: 56–61.

Kumar, R.S., 2001. Intertidal zonation and seasonality of benthos in a tropical mangrove. *Int. J. Ecol. and Environ. Sci.* 27: 199–208.

Kumari, K.L. and John, J., 2003. Biomass and quantitative indices of phytoplankton in Mandovi – Zuari estuary. NIO, Goa.

Kundu, S., Mondal, N., Lyla, P.S. and AjmalKhan, S., 2009. Biodiversity and Seasonal variation of Macrobenthic in faunal community in the inshore waters of Parangipettai Coast. *Environ. Monitor. Assess.* 163: 67-79.

Lakkis, S. and Novel-Lakkis, V., 1981. Composition, annual cycle and species diversity of the phytoplankton in Lebanese coastal water. *J. Plankton Research*. 3:123-126.

Lambshead, P.J.D., Platt, H.M. and Shaw, K.M., 1983. The detection of differences among assemblages of marine benthic species based on an assessment of dominance and diversity. *J. Nat. Hist.* 17: 859-874.

Langreth, R., 1994. The world according to Jan Janzen. *Popular Science*. 78-82: 112, 114-115.

Lee, S.Y., 1997. Potential trophic importance of the faecal materials of the mangrove crab *Sesarma messa*. *Mar.Ecol.Prog.Ser.* 159: 275-284.

Lee, S.Y., 1998. Ecological role of grasped crabs in mangrove ecosystems: a review. *Mar. Freshw. Res.* 49: 335-343.

López-Archilla, A.I., Moreira, D., López-García, P. and Guerrero, C., 2004. Phytoplankton diversity and cyanobacterial dominance in a hypereutrophic shallow lake with biologically produced alkaline pH. *Epub.* 8(2):109-15.

Madhav, G.V. and Kondalarao, B., 2004. Distribution of Phytoplankton in coastal waters of east coast of India. *Ind. J.Marine Science*. 33(2):262-268.

Madhupratap, M., Haridas, P., Ramaiah, N. and Achuthankutty, C.T., 1992. Zooplankton of the southwest coast of India: abundance, composition, temporal and spatial variability in 1987. *Oceanography of the Indian Ocean*. (Ed.) Desai, B.N. pp 99-112.

Madhupratap, M., Nair, S.R., Haridas, P. and Padmavati, G., 1990. Response of zooplankton to physical changes in the environment: coastal upwelling along central west coast of India. *J. Coastal research*. 6: 413-426.

Madondkar, S.G.P., Gomes, H., Parab, S.G., Pednekar, S. and Goes, J.I., 2007. Phytoplankton diversity, biomass, and production. *In* The Mandovi and Zuari estuaries. (Ed.) Shetye, S.R., DileepKumar, M. and Shankar, D." pp 67-81.

Magni, P. and Montani, S., 2000. Water chemistry variability in the lower intertidal zone of an estuary in the Seto Inland Sea, Japan: seasonal patterns of nutrients and particulate compounds. *Hydrobiologia* 432: 9–23.

Magurran, A.E., 1988. Ecological Diversity and its measurement. Princeton University press, Princeton, New Jercy.

Mahyavanshi, I.N., 1975. Further studies on marine planktonic diatoms in coastal waters Saurasthra. *Phykos.* 14(1-2): 99-111.

McIntyre, A.D., Elliot, J.M. and Ellis, D.V., 1984. Introduction: Design of sampling programmes. *In*: Methods for the Study of Marine Benthos (Ed.) Holme N.A. and McIntyre, A.D. Blackwell Scientific, Oxford. pp 1 - 26.

McKinnon, A.D. and Klumpp, D.W., 1998. Mangrove zooplankton in northern Queensland, I. Plankton community structure and environment. *Hydrobiologia*, 362:127-143.

Melluish, R.A.S., 1965. More Notes from Madras. *News Letter to Birdwatchers*. 5(10): 8.65.

Meybeck, M., Kimstach, V. and Helmer, R., 1996. Strategies for water quality assessment. *In*: [Ed.] Chapman, D. *Water quality Assessment: A Guide to the Use of Biota, Sediments and water in Environmental Monitoring*. (II Ed). Chapman and hall, London. pp 23-57.

Micheli, F., Gheradi, F. and Vannini, M., 1991. Feeding and burrowing ecology of two East African mangrove crabs. *Ma. Biol.* 111: 247-254.

Mishra, S. and Panigrahy, R.C., 1999. Zooplankton ecology of Bahuda estuary (Orrisa), East coast of India. *Indian J. Mar. Sci.* 28:297-301.

Misra, A., 1995. Anthozoa: Coelenterata. Wetland ecosystem series 1: Fauna of Chilika lake. Zoological Survey of India, Calcutta. pp 227-233.

Misra, S. and Kundu, R., 2005. Seasonal variations in population dynamics of key intertidal molluscs at two contrasting locations. *Aquatic Ecology*.29 (3): 315-324.

Mitra, A., Patra, K.C. and Panigrahy, R.C., 1990. Seasonal variations of some hydrographical parameters in tidal creek opening into the Bay of Bengal. *Mahasagar – Bull .Natn .Ins .Oceanogr.* 23(1): 55-62.

Mohamed M.A., Shriadah and Al-Ghais, S.M., 1999. Environmental characteristics of the United Arab Emirates waters along the Arabian Gulf: hydro chemical survey and nutrient salts. *Ind. J. Mar. Sci.* 28: 225-232.

Mohammed, S.Z., 1995. Observation on the benthic macrofauna of the soft sediment on western side of the Arabian Gulf (ROPME sea area) with respect to 1991 Gulf war oil pill. *Ind. J. Mar. Sci.* 24: 147–152.

Murphy, J. and Riley, J.P., 1962. A modified single solution method for the determination of phosphate in natural waters. *Analytica Chimica Acta*. 27:31-36.

Nair, V.R., 2002. Status of Fauna and Flora in Gulf of Kutch. National Institute of Oceanography, Regional Centre, Dr.Salim Ali Road, Kochi. pp 144.

Nandan, S.B. and Abdul Azis, P.K., 1993. Fish and shell fish fauna of retting and non-retting zones of the Kadinamkulam Kayal, Kerala. *J. Indian Fish. Ass.* 23: 35-43.

Niebauer, H.J., Smith, W.O. Jr., 1989. A numerical model of mesoscale physical-biological interactions in the fram strait marginal ice zone. *J.geophys.* res. 94(C11):61-75.

NIO (National Institute of Oceanography), 1998. Environmental studies for proposed BPCL jetty and associated facilities at Kandla. Part-I. Rapid marine EIA, May1998. NIO, R. C. 7 Bangalows, Mumbai.

Odum, H.T. and Odum, E.P., 1959. Fundamentals of Ecology. W.B. Saunders, Philadelphia, PA. pp 536.

Okemwa, E.N., 1990. A study of pelagic copepods in tropical marine creek, Tudor, Mombasa, Kenya with special reference to their community structure, Biomass and productivity. Ph.D thesis, Vrije Universiteit, Brussel. pp 225.

Padmavati, G. and Goswami, S.C., 1996. Zooplankton distribution in neuston and water column along west coast of India from Goa to Gujarat. *Ind. J. Mar. Sci.* 25: 85-90

Paulinose, V.T., Devi, C.B.L., Nair, V.R., Ramaiah. and Gajbhiye, S.N.N., 1998. Zooplankton standing and diversity in the Gulf of Kachchh with special reference to larvae of decapoda and pisces. *Ind. J. Mar. Sci.* 27:340-345.

Pearson, T.H. and Rosenberg, R., 1978. Macrobenthos secession in relation to organic enrichment and pollution of the marine environment. *Oceanog. Marine. Biol. Annual Review.* 16: 229–234.

Pereira, C., Rao, C.V. and Krishnan, S., 2002. Study of the dominant fauna of mangroves at seven bungalows beach, Versova, Mumbai –A preliminary study. *In* Proceedings of the National seminar on creeks, estuaries and mangroves – pollution and conservation, November. pp 196-200.

Pielou, E.C., 1966. The measurement of diversity in different types of biological collections. *J. Theoret. Biol.* 13: 144.

Plaziat, J.C., 1984. Mollusk distribution in the Mangal. *In*: Por, F.D, Dor, I (ed.), *Hydrobiology of the mangal*, Jumk, The Hague.

Prasad, M.N. and Mansuri, A.P., 1982. Population density of the Limpet *Cellana radiata* (Born) in polluted waters of Porbandar, West Coast of India. *J. Mar. Sci.* 11: 180-181.

Prasad, N.V., 2003. Diversity and richness of zooplankton in Coringa mangrove habitat. East coast. *J. Marine boil. Associ. India.* 45(1).

Quadros, G. and Athalye, R.P., 2002. Intertidal macrobenthos from Ulhas river estuary and Thane creek, *In*: Proceedings of the National seminar on creeks, estuaries and mangroves – pollution and conservation, November. pp 216-222.

Raghunathan, C., Tewari, A., Joshi, H.V., SaravanKumar, V.G., Trivedi R.H. and Khambhaty, Y., 2003. Impact of turbidity on Intertidal Macrofauna at Gopnath, Mahuva and Veraval coasts (West coast of India). *Ind.J.Mar.Sci.* 32(3): 214-221.

Ragothaman, G., and Patil, T., 1995. Studies on the physicochemical parameters and phytoplankton of Narmada estuary. *Enviromedia*. 14(2): 221–226.

Rainer, D.J., 1981. Temporal pattern in the structure of macrobenthic communities of an Australian estuary, Estuar. *Coast. Mar. Sci.* pp 597-619.

Rajasegar, M., 2003. Physico-chemical characteristic of the velar estuary in relation to shrimp farming. *J. Envron. Biol.* 24:95-101.

Rajendran, A., 1974. Studies on the ecology of velar estuary – chemistry and microbiology of regeneration of nutrients. Ph.D. Thesis, Annamalai University, India, pp 198.

Ramaiah, N., Ramaiah, N., Chandramohan, D. and Nair, V.R., 1995. Autotrophic and heterotrophic characters in polluted esturine complex. *Estn Coast Shelf Sci.* 40(1): 45-55.

Ramamurthy, S (1965); Studies on the plankton of the North Kanara coast in relation to the pelagic fishery; *Journal of Marine Biological Association of India*, 7 (1). pp. 127-149.

Ramamurthy, S. and Dhawan, R.M., 1963. On the characteristics of the plankton at the kandla in the Gulf Kachchh during August 1958-July 1960. *Indian .J. Fish.* 10 (1): 94-101.

Ramaraju, V.S., Sarma, V.V., Rao, N., T.V. and Vijayakumar, 1987. Variation of physico-chemical characteristics with tide in Visakhapatnam harbour waters, East Coast of India. *Indian J. Mar . Sci.* 16: 218-222.

Ramaswamy, V., Nath, B.N., Vethamony, P. and Ilangovan, D., 2007. Source and dispersal of suspended sediment in the macro-tidal Gulf of Kachchh. *Marine poll. Bullet.* 54:708-719.

Ramesh A. and Ramachandran, S., 2005. Factors influencing flamingo (*Phoenicopterus roseuis*) distribution in the Pulicat Lagoon ecosystem, India. Wetlands Ecology and Management. 3 (1):69-72.

Ramiah, N., Ramiah and Nair, V.R., 1998. Phytoplankton characteristics in a polluted Bombay harbor-Thana- Bassein creek estuarine complex. *Indian J. Mar.Sci.* 27: 281-285.

Rao, S., 1995. Fauna of Chilka Lake. Zoological Survey of India, Calcutta.

Redding, J.M. and Cory, R.L., 1975. *Macroscopic benthic fauna of three tidal creeks adjoining the Rhode river, Maryland*. Water Resources Investigation Report, USA. pp 39–75.

Ridd, P.V., 1996. Flow through animal burrows in mangrove creek. *Estuar. Coast. Shelf Sci.* 43: 617-625.

Roberston, A.I. and Blaber, S.J.M., 1992. Plankton, epibenthos and fish communities. *In* Roberston, A.I. and Alongi, D.M. (ed.) *Tropical Mangrove ecosystems*. American Geophysical Union, Washington. pp 173-224.

Roberston, A.I., Dixon, P. and Daniel, P.A., 1988. Zooplankton dynamics in mangroves and other near shore habitats in tropical Australia. *Mar. Ecol. Prog. Ser.* 43:139-150.

Robertson, A.I. 1986. Leaf – burying crabs: their influence on energy flow and export from a mangrove forest (*Rhizophora* spp.) in North eastern Australia, *J. Exp. Mar. Biol. Ecol.* 102: 231-248.

Robertson, D.I. and Daniel, P.A., 1989. The influence of crabs on leaf litter processing in a high intertidal mangrove forest in tropical Australia. *Oecologia*. 78: 191-198.

Roden, C. Roadhouse, M.P.G. and Hendry, M.P., 1987. Hydrography and the distribution of phytoplankton in Killar harbor. A Fjord in Western Ireland. *J. Mar. Biol. Ass. U.K.* 67: 359-371.

Sampathkumar, P., 1992. *Investigation on plankton in relation to hydrobiology and heavy metals in the Tranquebar-Nagapattinam coast, India*. Ph. D Thesis, Annamalai University, India. pp 254.

Sanders, H.L., 1968. Marine benthic diversity: A comparative study. *American Naturalist*. 102: 243–282.

Santhanam, P. and Perumel, P., 2003. Diversity of Zooplankton in Prangipettai coastal water, south east of India. *Indian Marine Ass.* 42(20):144-151.

Saraswathy., 1994. Hydrobiology of two estuarine systems (Arasalar and Cauvery) of South East Coast of India with special reference to plankton. Ph. D Thesis, Annamalai University. pp 144-155.

Saravanakumar A., Rajkumar, M., Thivakaran, G.A., and Sesh Serebiah J.; (2008); Abundance and seasonal variations of phytoplankton in the creek waters of western mangrove of Kachchh-Gujarat; *Journal of Enviornmental biology*, 29(2) 271-274.

Saravanakumar, A., Rajkumar, M., Thivakaran, G.A. and Serebiah, J.S., 2008. Abundance and Seasonal Variation of Phytoplankton in the creek water of western mangroves of Kachchh-Gujarat. *J. Environ. Biol.* 29(2): 271-274.

Saravankumar, A, Rajmumar, M, Jun Sun, Serebiah J. S and Thivakaran G. A (2009); Forest structure of arid zone mangroves in relation to their physical and chemical environment in the western Gulf of Kachchh, Gujarat, Northwest coast of India. *Journal of coastal conservation* 13(4), 217-234

Sarkar, Ş.R.D., 1995. *Nematoda*: Wetland Ecosystem Series 1: Fauna of Chilika Lake. Zoological Survey of India, Calcutta. pp 279-318.

Sarma, A.L.N. and Wilsanand, V., 1996. Meiofauna of the outer channel of Chilka lagoon, Bay of Bengal. *Indian J. Mar. Sci.* 25: 302-306.

Sasekumar, A., 1974. Distribution of macrofauna on a Malayan mangrove shore. *J. Anim. Ecol.* 43: 51-69.

Sawant, S.S. Madhupratap, M (1996) Seasonality and composition of phytoplankton in the Arabian Sea; *Current Science*, Vol.71; 869-873p.

Selvam, V., Azariah, J. and Azariah, H., 1992. Diurnal Variation in Physio-Chemical Properties and Primary Production in the interconnected marine, mangrove and freshwater biotopes of Kakinada coast, Andhra Pradesh, India. *Hydrobiogia*. 247:181-186.

Selvaraj G.S.D, Thomas V.J. and Khambadkar, L.R., 2003. Seasonal variation of phytoplankton productivity in the surf zone and back water at Cochin. *J. Mar. Biol. Assoc. India.* 45(1):9-19.

Senthilkumar, R and Sivakumar, K., 2008. Phytoplankton diversity in Veeranam Lake in response to abiotic factors in Cuddalore district of Tamilnadu. *J. Environ. Biol.* 29(5):747-752.

Seshappa, G., 1953. Observations on the physical and biological features of the sea bottom along the Malabar Coast. *Proceed. Nation. Inst. Sciences of India.* 19: 257–279.

Shepard, R.N., 1962. The analysis of proximities: Multidimensional scaling with an unknown distance function. *Phychometrika*. 27: 125-140.

Singh, H.S., 2000. Mangroves in Gujarat: Current Status and Strategy for Conservation. GEER Foundation, Gandhinagar, Gujarat.

Singh, S.N.S.B.M. and Chaudhry, A., 1986. Zooplankton Production in Coastal Waters. *Proc. Indian. Asso. Sci (Animal Sci)*. 95:125.

Sithik, M.A., Thirumurugan, A., Arumugam, G., Kannan, R.R.R. and Anatharam, P., 2009. Studies of Phytoplankton diversity from agnitheertham and Kothandaramar Koil Coastal waters; South coast of India. *Global J. Environ. Research.* 3(2):118-125.

Sivasamy, S.N., 1990. Plankton in relation to Coastal Pollution at Ennore, Madras Coast. *Ind. J. Mar. Sci.*19: 115-119.

Smith, T.J, Botto, K.G., Rusher, S.D. and Giddens, R.L., 1991. Keystone species and mangrove forest dynamics: the influence of burrowing by crabs on soil nutrient status and forest productivity. *Estuar. Coast. Shelf Sci.* 33(5): 419-432.

Sreekumaran NS R.; Devassy V.P.; Madhupratap M.; Blooms of phytoplankton along the west coast of India associated with nutrient enrichment and the response of zooplankton *Science of the total environment* ISSN 0048-9697

Stieglitz, T., Ridd, P.and Muller, P., 2000. Passive migration and function morphology of crustacean burrows in a tropical mangrove swamp. *Hydrobiologia*. 421: 69-76.

Strickland J.D.H., Parsons T.R. (1972) A practical hand book of sea water analysis. *Bull.Fish.Res.Board Can.* 167. 310 p.

Subrahmanyan R. (1959); Studies on the phytoplankton of the west coast of India; *Proceedings: Plant Sciences* 50(3), 113-187.

SunderRaj, S.K. and Sanjeeva Raj, P.J., 1984. Polychaeta of Pulicat Lake, (TamilNadu). J. Bom. Nat. Hist. Soc. 84: 84-104.

Surekha Sawant and Madhupratap, M., 1996. Seasonality and composition of phytoplankton in the Arabian Sea. *current science*. 71(11): 869-873

Sverdrup, H.W., Johnson, M.W. and Fleming, R.H., 1942. The oceans, their physics, chemistry and general biology. New York: Prentice – Hall, inc.

Swami, B.S., Suryawanshi, U.G. and Karande, A.A., 2000. Water quality status of Mumbai harbour – an update. *Ind. J. Mar. Sci.* 29:111-115.

Thangavelu, R. and SanjeevaRaj, P.J., 1985a. Fishery and Biology of the clam Meretrix casta (Chemnitz) in the Pulicat Lake. *J. Mar. Biol. Ass. India.* 27 (1 & 2): 75-83.

Thangavelu, R. and SanjeevaRaj, P.J., 1985b. Exploitation of Shell-Deposits in Pulicat Lake. *J. Mar. Boil. Ass. India.* 27(1 & 2): 124-128.

Thangavelu, R., Thangavelu, R. and SanjeevaRaj, P.J., 1988. Distribution of Molluscan Fauna in Pulicat Lake. *In* National Seminar on Shellfish Resources and Farming. Aug. 1988. CMFRI Bulletin, 42, Part I. pp 58-67.

Tilman, D., Kilham, S.S and Kilham, D., 1982. Phytoplankton community ecology. The role of limiting nutrients. *Ann. Rev. Ecolog. Syst.* 13: 349-372.

Tiwari, L.R., and Nair, V.R., 1998. Ecology of phytoplankton from Dharmatar Creek, West coast of India. *Ind. J. Mar. Sci.* 27:302-309

Trivedy, R.K. and Goel, P.K., 1984. Physicochemical analysis of water. *In: Chemical and biological methods for water pollution studies*. Environmental Publications, Karad, India. pp 35-96.

Upadhyaya, S.; 1988. Physico-chemical characteristics of the Mahanadi estuarine ecosystem, East coast of India. *Indian J. mar. sci.* 17: 19-23.

Valdes, D. S and Real, E., 1998. Variations and relationships of salinity, nutrients and suspended solids in Chelem coastal lagoon at Yucatan, Mexico. *Indian. J. Mar. Sci.* 27: 149-156.

Valiela, I., 1995. Marine ecology. Springer-Verlag, New York.

Vareethiah, K and Haniffa, M.A., 1997. Quantitative indices of phytoplankton in bar-built estuary. *J. Bio. Ass. India.* 39 (1-2): 155-159.

Venkatachalam, S. and Kale, P.G., 2002. Benthic Macrofauna of Thane creek: Seasonal variations, *In*: Proceedings of the National seminar on creeks, estuaries and mangroves – pollution and conservation, November. pp 186-195.

Venkatraman, K. and Wafar, M., 2005. Coastal and Marine Biodiversity of India. *Indian J.Mar.Sci.* 34(1): 57-75.

Vethamony, Babu, P., Reddy, M.T., Sudheesh, G.S., Desa, K. and Zingde, E., 2007. Estimation of carrying capacity of the Gulf of Kachchh, west coast of India in relation to petroleum hydrocarbon through oil spill modelling. *In* Proceedings of the International Maritime-Port Technology and Development Conference. MTEC 2007, Maritime and Port Authority of Singapore. pp 505-511.

Viswanathan P K., Pathak K.D. and Mehta I.; Socio economic and ecological benefits of mangrove plantation: a study of community base mangrove restoration activities in Gujarat. Gujarat Ecology Commission, Gandhinagar 128 p.

Walsh, J.J., 1975. A spatial simulation model of the Peru upwelling ecosystem. *Deep-sea Res.* 22:201-36.

Warren, J.H and Underwood, A.J., 1986. Effect of burrowing crabs on the topography of mangrove swamps in New South Wales. *J. Exp. Mar. Biol. Ecol.* 102: 223-236.

Bibliograph

Warwick, R.M. and Clarke, K. R., 2001. Practical measure of marine biodiversity based on relatedness of species. *Oceanography and Marine Biol: Annual Review.* 39: 207-231.

Warwick, R.M. and Clarke, K.R., 1991. A comparison of some methods for analyzing changes in benthic community structure. *J. Mar. Biol. Assoc. UK*. 71(1): 225 - 244.

Warwick, R.M., 1989. Role of meiofauna in the marine ecosystem: Evolutionary considerations. *Zool. J. Linn. Soc.* 96: 229-241.

Weaver, W. and Shannon, C.E., 1949. *The Mathematical Theory of Communication*. Urbana, University of Illinois, Illinois.

Wells, F.E., 1984. Comparative distribution of macro molluscs and macro crustaceans in a North-Western Australian Mangrove system. *Aust. J. Mar. Freshw. Res.* 35: 591-596.

Wooldridge, T.T., Erasmus T. and Estuar, J., 1980. Coast. Mar. Sci. pp 37-58.

Zacharias, M.A and Howes, D.E., 1998. An analysis of Marine protected area in British Columbia, Canada, using marine ecological classification. *Natural Area*. 18: 4-13.

Zalocar-de-Domitrovic, Y., Poi-de-Neiff, A.S.G. and Casco, S.L.,2007. Abundance and diversity of Phytoplankton in the Paraná River (Argentina) 220 km downstream of the Yacyretá reservoir. Braz. *J. Biol.* 67(1).

Zingde, M. D., Abidi, S.A.H., Sarma, P. and Rokade, M.A., 1987. Base water quality of Thailand I: Contributions in marine sciences. *Dr. S. Z. Qasim, Sixteenth Birthday Felicitation Volume*. pp 307–318.

Zingde, M.D., Sharma, P. and Sabnis, M.M., 1985. Physico-chemical investigation in Auranga river. *Mahasagar*. 18: 307-321.