

**REFERENCES**

- Agostini, N. (2002). La migrazione dei rapaci in Italia. Brichetti, P. and Gariboldi, A. [Eds]. *Manuale di Ornitologia*. Edagricole, Bologna, pp 157-182.
- Agostini, N. and Duchi, A. (1994). Water-crossing behavior of Black Kites (*Milvus migrans*) during migration. *Bird Behavior*, **10 (1.4)**:45-48.
- Agostini, N. and Panuccio, M. (2010). Western Marsh Harrier (*Circus aeruginosus*) migration through the Mediterranean Sea: a review. *Journal of Raptor Research*, **44(2)**: 136-142.
- Agostini, N., Lucia, G., Mellone, U., Ashton booth, J., Wilson, S., Chiatante, G. and Todisco, S. (2007). Le Serre Catanzaresi: Important Bird Area (IBA) per la migrazione autunnale dei rapaci. Page 68 in Proceedings of the XIV Italian Ornithological Congress.
- Alberti, M.A., Marzluff, J.M., Shulenberger, E., Bradley, G., Ryan, C. and Zumbrunnen, C. (2003). Integrating humans into ecology: opportunities and challenges for urban ecology. *BioScience*, **53**: 1169-1179.
- Ali, S. (1979). *The book of Indian Birds*. 11<sup>th</sup> Edition. Bombay Natural Society, Bombay.

- Altken, E. H. (1947). *The common birds of India*. Thacker and Co. Ltd, Bombay.
- Altmeyer, M., Dittman, J., Dmowski, K., Wagner, G. and Muller, P. (1991). Distribution of elements in flight feathers of a white tailed eagle. *Science of Total Environment*, **105**: 157-164.
- Anderson, D. L. (2001). Landscape heterogeneity and diurnal raptor diversity in honduras: the role of indigenous shifting cultivation. *Biotropica*, **33**: 511–519.
- Andersson, M. and Erlinge, S. (1977). Influence of predation on rodent populations. *Oikos*, **29**: 591-597.
- Bakaloudis, D. E. (2009). Implications for conservation of foraging sites selected by Short-toed Eagles (*Circaetus gallicus*) in Greece. *Ornis Fennica*, **86**: 89-96.
- Bakaloudis, D. E. (2010). Hunting strategies and foraging performance of the Shorttoed Eagle in the Dadia-Lefkimi-Soufli National Park, north-east Greece. *Journal of Zoology*, **281**: 168-174.
- Beale, C. M., and Monaghan, P. (2004). Behavioral responses to human disturbance: a matter of choice? *Animal Behaviour*, **68**:1065–1069.

- Begon, M., Harper, J. L. and Townsend, C. R. (1996). *Ecology - Individuals, Populations and Communities*. 3<sup>rd</sup> Edition. Blackwell, Oxford, UK.
- Begum, A. and Sehrin, S. (2013). Levels of Heavy Metals in Different Tissues of Pigeon (*Columba livia*) of Bangladesh for Safety Assessment for Human Consumption. *Bangladesh Pharmaceutical Journal*, **16(1)**: 81-87.
- Benson, W., Brooker, R. K., Dowsett, J. and Irwin M. P. (1971). *The birds of Zambia*. London, Collins Publishing Co. **65**: 148-164.
- Bervoets, L. and Blust, R., (2003). Metal concentrations in water, sediment and gudgeon (*Gobio gobio*) from a pollution gradient: relationship with fish condition factor. *Environmental Pollution*, **126**: 9-19.
- Bijlsma, R.G. (1997). Black Kite. W.J.M Hagemeijer and M.J. Blair [Eds], The EBCC atlas of European breeding birds, their distribution and abundance. T. and A.D. Poyser London, U.K. pp 132-133.
- Binkowski, Ł.J., Stawarz, R. M. and Zakrzewski, M. (2013). Concentration of cadmium, copper and zinc in tissue of mallard and coot from southern Poland. *Journal of environmental science and Health, Part:B*. **48**: 410-415.

- Bird Life International (2009) "*Milvus migrans*", IUCN Red List of Threatened Species. Version 2009.2. International Union for Conservation of Nature. Retrieved 2010-02-15.
- Bird, D. M., Varland, D. E. and Negro, J. J. (1996). *Raptors in Human Landscapes: Adaptations to Built and Cultivated Environments*. Academic press, London.
- Bird Life International. (2007). Bird Life's online world bird database. BirdLife International, Cambridge, United Kingdom. Available from <http://www.birdlife.org/datazone/index.html>.
- Birrer, S. (2010). Synthesis of 312 studies on the diet of the Long Eared Owl *Asio otus*. *Ardea*, **97**: 615- 624.
- Blair, R. B. (1996). Land use and avian species diversity along an urban gradient. *Ecological Applications*, **6**:506-519.
- Blake, J. G and Loiselle, B. A. (1991). Variation in resource abundance affects capture rates of birds in three lowland habitats in Costa Rica. *Auk*, **108**: 114-130.
- Blust, R., Van der Linden, A., Verheyen, E. and Decler, W. (1988). Evaluation of microwave heating digestion and graphite furnace atomic absorption spectrometry with continuum source background correction for the determination of Fe, Cu and Cd in brine shrimp. *Journal of Analytical Atomic Spectrometry*, **3**: 387-393.

- Boag, P. T and Grant, P. R. (1984). Darwin's finches (Geospiza) on Isla Daphne Major, Galapagos: Breeding and feeding ecology in a climatically variable environment. *Ecological Monograph*, **54**: 463- 489.
- Bolger, D. T., Alberts, A. C., Sauvajot, R. M., Potenza, P., McCalvin, C., Tran, D., Mazzoni, S., Soulé, M. E. (1997). Response of rodents to habitat fragmentation in coastal southern California. *Ecological Applications*, **7**: 552–563.
- Bontzorlos, V. A., Peris, S. J., Vlachos, C. G. and Bakaloudis, D. E. (2005). The diet of Barn Owl in agricultural landscapes of Central Greece. *Folia Zoologica*, **54**: 99-110.
- Boulinier, T. (1996). On breeding performance, colony growth and habitat selection in Buff Necked Ibis. *Condor*, **98**: 440-441.
- Bourquin, J. D. (1983). Mortalité des rapaces le long de l'autoroute Genève-Lausanne. *Nos Oiseaux*, **393**: 149-169.
- Bowman, R. and Woolfenden, G. E. (2001). *Nest success and timing of nest failure of Florida ScrubJays in suburban and wildland habitats*. Kluwer Academic Publishers, Boston, MA. pp 383- 402.
- Bradley, M., Johnstone, R., Court, G. and Duncan, T. (1997). Influence of weather on breeding success of Peregrine Falcons in the Arctic. *Auk*, **114**: 786–791

- Braune, B. M., and David E. G. (1987): A mercury budget for the Bonaparte's gull during autumn moult. *Ornis Scandinavica*, **18**: 244-250.
- Brown, L. R., Gardner, G. and Halweil, B. (1998). *Beyond Malthus: 16 dimensions of the population problem*. World watch Paper 143. Worldwatch Institute, Washington, DC.
- Brown, M. A. (1964). Food poisoning involving zinc contamination. *Archives of Environmental Health*. **8(5)**: 657-660.
- Brown, T., Dawson, C. and Miller, R. (1979). Interest and attitudes of metropolitan New York residents about wildlife. *North American Wildlife Nature Resource Conference*. **44**: 289-297.
- Bubb, J. M and Lester, J. N. (1991). The impact of heavy metal on lowland river and the implication for man and the environment. *The science of the total environment*, **100**: 207-233.
- Burger, J. (1978a). The pattern and mechanism of nesting in mixed species heronries. New York. pp 45-48.
- Burger, J. (1978b). Competition between Cattle Egret and native North American herons, egrets and ibises. *Condor*, **80**: 15-23.
- Burger, J. (1993). Metals in avian feathers: bioindications of environmental pollution. *Review of Environmental Contamination and Toxicology*, **5**: 203-311.

- Burger, J. (1995). A risk assessment for lead in birds. *Journal of Toxicology Environmental Health*, **45**:369- 396.
- Burger, J. and Gochfeld, M. (1988). Nest-site selection and temporal patterns in habitat use of Roseate and Common terns. *Auk*, **105**: 433-438.
- Burger, J. and Gochfeld, M. (1997). Age differences in metals in the blood of Herring gull (*Larus pipixcan*). *Archives of Environmental Contamination and Toxicology*, **18**: 673- 678.
- Burger, J. and Hahn, C. (1989). Crow predation on Black-crowned Night Heron eggs. *Wilson Bulletin*, **89**: 350-351.
- Burger, J., Woolfenden, G. E. and Gochfeld, M. (1999). Metal concentrations in the eggs of endangered Florida scrub jays from central Florida. *Archives of Environmental Contamination and Toxicology*, **37**: 385-388.
- Bustamante, J. (1994). Behaviour of colonial common kestels (Falcon tinunculus) during the post-fledging dependece period in southwestern Spain. *Journal of Raptor Research*, **28(2)**: 79-83.
- Cahil, T. M., Anderson, D. W., Elbert, R. A., Perley, B. P. and Johnson, D. R. (1998). Elemental profiles in feather samples from a mercury- contaminated lake in central California. *Archives of Environmental Contamination and Toxicology*, **35**: 75-81.

- Cameroon, G. N. (2003). Prey use by a pair of barn owl (*Tyto alba*) in Texas coastal habitat. *Texas Journal of Science*, **55**: 219–224.
- Carney, K.M., and Sydeman, W. J. (1999). A review of human disturbance effects on nesting colonial waterbirds. *Waterbirds*, **22**:68–79.
- Carrete, M. and Donázar, J. A. (2005). Application of central place foraging theory shows the importance of Mediterranean dehesas for the conservation of the cinereous vulture, *Aegypius monachus*. *Biological Conservation*, **126**: 582–590.
- Carrete, M.A., Jose, L., Tella, A., Blanco, B. and Beretellotti, C. (2009). A Effects of habitat degradation on the abundance, richness and diversity of raptors across Neotropical biomes. *Biological Conservation*, **142(10)**: 2002-2011.
- Catterall, C. P., Green, R. J., Jones, D. N. (1991). *Habitat use by birds across a forest-suburb interface in Brisbane: implications for corridors*. Saunders, D. A. and Hobbs, R. J. [Eds.], Nature Conservation 2: The Role of Corridors. Surrey Beatty and Sons, Chipping Norton, pp 247–258.
- Chace, J. F. and Walsh, J. J. (2006). Urban effects on native avifauna: a review. *Landscape Urban Planning*, **74**: 46–69.

- Chace, J. F., Walsh, J. J., Cruz, A., Prather, J. W. and Swanson, H. M. (2003). Spatial and temporal activity patterns of the brood parasitic Brown-headed Cowbird at an urban/wildland interface. *Landscape and Urban Planning*, **64**:179-190.
- Clark, A. J., and Scheuhammer, A. M. (2003). Lead poisoning in upland-foraging birds of prey in Canada. *Ecotoxicology*, **12**: 23-30.
- Clark, R. G., Hobson, K. A., Nichols, J. D. and Bearhop, S. (2004). Avian dispersal and demography: scaling up to the landscape and beyond. *Condor*, **106**:717–719.
- Clergeau, F. and Quenot, F. (2007). Roost selection flexibility of European starlings aids invasion of urban landscape. *Landscape and Urban Planning*, **80**: 56 -62.
- Clergeau, P., Jokimäki, J. and Savard, J. (2001). Are urban bird communities influenced by the bird diversity of adjacent landscapes? *Journal of Applied Ecology*, **38**: 1122–1134.
- Cohen, T., Hee, S. and Ambrose, R. (2001). Trace metals in fish and invertebrates of three California coastal wetlands. *Marine Pollution Bulletin*, **42**: 224-232.
- Collias, N.E. and Collias, E.C. (1984). *Nest building and bird behaviour*. Princeton University Press, Princeton, NJ.

- Collopy, M. W. (1983). A comparison of direct observations and collections of prey remains in determining the diet of Golden Eagles. *Journal of Wildlife Management*, **47**:360- 368.
- Cooper, D. S. (2002). Geographic associations of breeding bird distribution in an urban open space. *Biological Conservation*, **104**: 205– 210.
- Corlett, R. (1992b). The changing urban vegetation. Gupta, A.J., Pitts, J. [Eds.], *Physical Adjustments in a Changing Landscape: The Singapore Story*. Singapore University Press, Singapore, pp 190– 214.
- Coulson, J.C. (1968). Differences in the quality of birds nesting in the centre and on the edge of a colony. *Nature*, **217**: 478-479.
- Crooks, K. R., Suarez, A.V., Bolger, D. T. and Soule. M. E. (2001). Extinction and colonization of birds on habitat islands. *Conservation Biology*, **15**: 159-172.
- Dauwe, T., Bervoets, L., Janssens, E., Pinxten, R., Blust, R. and Eens, M. (2002). Great and blue tit feathers as biomonitors for heavy metal pollution. *Ecological Indicators*, **1**: 227-234.
- Davis, S. K., Brigham, R. M., Shaffer, T. L., and James, P. C. (2006). Mixed-grass prairie passerines exhibit weak and variable responses to patch size. *The Auk*, **123**(3): 807-821.

- DeForest, D. K., Schlekot, C. E., Brix, K. V. and Fairbrother, A. (2012). Secondary poisoning risk assessment of terrestrial birds and mammals exposed to Nickel. *Integrated environmental assessment and management*, **8(1)**:107-119.
- DeSante, D. F. and Geupel, G. R. (1987). Land bird productivity in central coastal California: The relationship to annual rainfall, and a reproductive failure in 1986. *Condor*, **89**: 636-653.
- Dhinsa, S.M., Peter, E.K. and David, A.B. (1989). Nest height of Black-Billed Magpies: Is it determined by human disturbance or habitat type? *Canadian Journal of Zoology*, **67**: 228-232.
- Dmowski, K. (1999). Birds as bioindicators of heavy metal pollution: review and examples concerning European species. *Acta Ornithologica*, **34(1)**: 1-25.
- Donazar, J.A., Ceballos, O., Travaini, A., Rodriguez, A., Funes, M., and Hiraldo, F. (1994). Breeding performance in relation to nest site substratum in a Buff-necked Ibis (*Theristicus caudatus*) population in Patagonia. *Condor*, **96**: 994- 1002.
- Donazar, J.A., Travaini, A., Rodriguez, A., Ceballos, O. and Hiraldo, F. (1996). Nesting association of Raptors and Buff-necked Ibis in the Argentinean Patagonia. *Colonial Waterbirds*, **19**: 11-115.

- Dusi, J. L. (1968). The competition between Cattle Egrets and Little Blue Herons. *Alabama Birdlife*, **16**: 4-6.
- Dżugan, M., Zielińska, S., Hęclik, J., Pieniążek, M. and Szostek, M. (2012). Evaluation of heavy metal environmental contamination based on their concentrations in tissues of wild Pheasant (*Phasianus colchicus*). *Journal of Microbiology, Biotechnology and Food Sciences*, **2(1)**: 238-245.
- Eens, M., Pinxten, R., Verheyen, R. F., Blust, R. and Bervoets, L. (1999). Great and blue tits as indicators of heavy metal contamination in terrestrial ecosystems. *Ecotoxicology and Environmental Safety*, **44**: 81-85.
- Eeva, T. and Lehikoinen, E. (2000). Recovery of breeding success in wild birds. *Nature*, **403**: 851-852.
- Eiserer, L. A. (1984). Communal roosting in Birds. *Bird Behaviour*. **5**: 61 - 80.
- Eisler, R. (1986). *Chromium hazards to fish, wildlife, and invertebrates: a synoptic review*. Patuxent Wildlife Research Center, Laurel, MD (USA).
- Elkin, N. (1988). *Weather and bird behaviour*, 2<sup>nd</sup> Edition. TandA D Poyser, Calton.

- Ellison, L. N., and Cleary, L. (1978). Effects of human disturbance on breeding of double crested cormorants. *The Auk*, **95**:510-517
- Emlen, J.T., (1974). An urban bird community in Tucson, Arizona: derivation, structure, regulation. *Condor*, **76**: 184–197.
- Erdoğan, Ö. and Erbilir, F. (2007). Heavy metal and trace elements in various fish samples from Sir Dam Lake, Kahramanmaraş, Turkey. *Environmental Monitoring and Assessment*, **130**: 373-379.
- Esselink, H., Van der Geld, F. M., Jager, L.P., Posthuma- Trupie, G.A., Zoun, P. and Baars, A. J. (1995). Biomonitoring heavy metals using the Barn Owl (*Tylo alba guttata*): sources of variation specially relating to body condition. *Archives of Environmental contamination and toxicology*, **28**: 471-486.
- Esselink, H., Van der Geld, F. M., Jager, L.P., Posthuma- Trupie, G.A., Zoun, P. and Baars, A.J. (1995). Biomonitoring heavy metals using the Barn Owl (*Tylo alba guttata*): sources of variation specially relating to body condition. *Archives of Environmental contamination and toxicology*. **28**: 471-486.
- Estrella, R. R., Donázar, J.A. and Hiraldo, F. (1998). Raptors as Indicators of Environmental Change in the Scrub Habitat of Baja California Sur, Mexico. *Conservation Biology*, **12**(4): 921–925.

- Everding, S. E. and Jones, D. N. (2006). Communal roosting in a suburban population of Torresian crows (*Corvus orru*). *Landscape and Urban Planning*, **74**: 21 -33.
- Faulkner, S. (2004). Urbanization impacts on the structure and function of forested wetlands. *Urban Ecosystems*, **7**: 89-106.
- Fernandez-Juricic, E. (2000). Avifaunal use of wooded streets in an urban landscape. *Conservation Biology*, **14**:513-521.
- Fernández-Juricic, E. and Tellería, J. L. (2000). Effects of human disturbance on Blackbird *Turdus merula* spatial and temporal feeding patterns in urban parks of Madrid, Spain. *Bird Study*, **47**: 13–21.
- Fetterolf, P. M. (1983). Effects of investigator activity on Ring-billed Gull behavior and reproductive performance. *Wilson Bulletin*, **95**:23-41.
- Finlayson, C. (1992). *Birds of the Strait of Gibraltar*. London, Poyser. Forsman.
- Fowler, G. S. (1999). Behavioral and hormonal responses of Magellanic penguins (*Spheniscus magellanicus*) to tourism and nest site visitation. *Biological Conservation*, **90**:143–149.
- Fretwell, S. D. (1972). *Populations in a seasonal environment*. Princeton University Press, Princeton, NJ.

- Furness, R. W. (1996) Cadmium in birds. Beyer, W. N., Heinz, G. H., Redmon-Norwood, A. W. [Eds.] *Environmental contaminants in wildlife*. CRC Press, Boca Raton. pp 389–404.
- Furness, R.W. (1993). *Birds as monitors of pollutants*. Furness, R. W., Greenwood, J. J. D. [Eds.] Chapman and Hall, New York, pp 86–143.
- Furness, R.W., Lewis, S.A. and Mills, J. A. (1990). Mercury Levels in the Plumage of Red Gulls *Larus novaehollandiae scopulinus* of known Sex and Age. *Environmental Pollution*, **63**: 33-39.
- Gad, S. C. (1989). Acute and chronic systemic chromium toxicity. *Science of the Total Environment*, **86** (1): 149-157.
- Galushin, V. M. (1971). A huge urban population of birds of prey in Delhi, India. *Ibis*, **113**: 552.
- Gamauf, A., Tebb, G. and Nemeth, E. (2013). Honey Buzzard *Pernis apivorus* nest-site selection in relation to habitat and the distribution of Goshawks *Accipiter gentilis*. *Ibis*, **155**: 258–270.
- Garcia- Fernandez, A.J, Motas-Guzman, M., Navas, I., Maria-Mojica, P., Luna, A. and Sanchez-Garcia, J.A. (1997). Environmental exposure and distribution of lead in four species of raptors in southern eastern Spain. *Achieves of Environmental Contamination and Toxicology*, **33**: 76-82.

- Gardner, J. E., Garner, J. D. and Hofmann, J. E. (1991). Summer roost selection and roosting behavior of *Myotis sodalis* (Indiana bat) in Illinois. Illinois Natural History Survey/Illinois Department of Conservation, Champaign, USA.
- Getz, L. (1961). Hunting areas of the long-eared owl. *The Wilson Bulletin*, 1:79-82.
- Giese, M. (1996). Effects of human activity on Adélie penguins *Pygoscelis adeliae* breeding success. *Biological Conservation*, 75:157-164.
- Gill, J. A., Norris, K. and Sutherland, W. J. (2001). The effects of disturbance on habitat use by black-tailed godwits, *Limosa limosa*. *Journal of Applied Ecology*, 38:846- 856.
- Gillett, W. H., Hayward, J. L. and Stout, J. F. (1975). Effects of human activity on egg and chick mortality in a Glaucous-winged Gull colony. *Condor*, 77: 492-495
- Gochfeld, M., Belant, J., Shulka, T., Benson, T. and Burger, J. (1996). Heavy metals in laughing gulls: gender, age and tissue differences. *Environmental Toxicology and Chemistry*, 15: 2275-2283.
- Götmark, F. (1992). Blue eggs do not reduce nest predation in the song thrush, *Turdus philomelos*. *Behavioral Ecology and Sociobiology*, 30:245-252

- Green, R.E., Newton, I., Shultz, S., Cunningham, A.A., Gilbert, M., Pain, D.J., Prakash, V. (2004). Diclofenac poisoning as a cause of vulture population declines across the Indian subcontinent. *Journal of Applied Ecology* **41**: 793–800.
- Greenberg, R. (1981). Frugivory in some migrant tropical forest wood warblers. *Biotropica*, **13**: 215- 223.
- Greenwood, P.J. and Harvey, P.H. (1976). The adaptive significance of variation in breeding Area fidelity in Black Bird (*Turdus merula*). *Journal of Animal Ecology*, **45**: 887-898.
- Gunn, S. A. and Gould, T. C. (1957). Selective accumulation of Cd115 by cortex of rat kidney. *Experimental Biology and Medicine*, **96(3)**: 820-823.
- Hanski, I., Hansson, L. and Henttonen, H. (1991). Specialist predators, generalist predators, and the microtine rodent cycle. *Journal of Animal Ecology*, **60**: 353-367.
- Heinz, G. (1976). Methylmercury: second-year feeding effects on mallard reproduction and duckling behavior. *Journal of Wildlife Management*, **40**: 82-90.
- Hernandez, M. (1988). Road mortality of the little owl (*Athene noctua*) in Spain. *Journal of Raptor Research*, **22**: 81-84.

- Holmes, R. T., Marra, P. P. and Sherry, T. W. (1996). Demographic responses of breeding Blackthroated Blue Warblers (*Dendroica caerulescens*) to a heterogeneous habitat: implications for population dynamics. *Journal of Animal Ecology*, **65**:183–195.
- Holmgren, J. (2004) Roosting in tree foliage by Common Swift *Apus apus*. *Ibis*, **146**: 404 -416.
- Hostetler M. (2001). The importance of multi-scale analyses in avian habitat selection studies in urban environments, In J. M. Marzluff, R. Bowman, and R. Donnelly [Eds.], *Avian ecology and conservation in an urbanizing world*. Kluwer Academic Publishers, Boston, MA. pp 139-154.
- Huang, K.Y., Lin, Y.S., Severinghaus, L.L., (2006). Comparison of three common methods for studying the diet of nestling of two Accipiter species. *Zoological Studies*, **(45)**: 234 - 243.
- Hui, A., Takekawa, J., Baranyuk, V. and Litvin, K. (1998). Trace element concentrations in two subpopulations of lesser snow geese from Wrangel Island, Russia. *Archives of Environmental Contamination and Toxicology*, **34**: 197-203.
- Hui, C. (1998). Elemental contaminants in the livers and ingest of four subpopulations of the American coot (*Fulica americana*): an

- herbivorous winter migrant in San Francisco Bay. *Environmental Pollution*, **101**: 321-329.
- Hutton, M. (1981). Accumulation of heavy metals and selenium in three seabird species from the United Kingdom. *Environmental Pollution*, **26** : 129-145.
- Illner, H. (1992). Road deaths of Westphalian owls: methodological problems, influence of road type and possible effects on population levels. In: Galbraith, C. A., Taylor, I. R., Percival, S. [Eds.], *The Ecology and Conservation of European Owls*. Joint Nature Conservation Committee, Peterborough UK, pp 94-100.
- Imhoff, M. L., Bounoua, L., Defries, R., Lawrence, W.T., Stutzer, D., Tucker, C.J. and Ricketts, T. (2004). The consequences of urban land transformation on net primary productivity in the United States. *Remote Sensing of Environment*, **89**:434-443.
- Immelmam, K. (1971). Ecological aspects of periodic reproduction. Farner, D. S. and King, J. R. [Eds.] *Avian biology*. Vol. 1. Academic Press, New York, pp 342-390.
- IUCN. (2002) 2002 IUCN red list threatened species. Available from: <http://www.redlist.org>.
- Iwata, H., Tanabe, S., Sakai, N. and Tatsukawa, R. (1993). Distribution of persistent organochlorines in oceanic air and surface seawater

and the role of ocean in their global transport and fate.

*Environmental Science and Technology*, **27**: 1080-1098.

Jackson, S. L., Hik, D. S. and Rockwell, R. F. (1988). The influence of nesting habitat on reproductive success of Lesser Snow Geese.

*Canadian Journal of Zoology*, **66**: 1699-1703.

Jacob, R. A., Skala, J. H., Omaye, S. T. and Turnlund, J. R. (1987). Effect of varying ascorbic acid intakes on copper absorption and ceruloplasmin levels in young men. *Journal of Nutrition*, **117**:2109-2115.

Janicke, T. and Chakarov, N. (2007). Effect of weather conditions on the communal roosting behaviour of common ravens *Corvus corax* with unlimited food resources. *Journal of Ethology*, **25**: 71-78.

Jayson, E.A. and Mathew, D.N. (1995) Roosting behaviour of common Indian Mayna (*Acridotheres tristis*) at Trichur, Kerala. *Pavo*, **33 (1 and 2)**: 41-46.

Jeejeebhoy, K. N. (1999). Chromium and parenteral nutrition. *The Journal of Trace Elements in Experimental Medicine*, **12(2)**: 85-89.

Jokimäki, J., Clergeau, P., Kaisanlahti-Jokimäki, M. L. (2002). Winter bird communities in urban habitats: a comparative study between central and northern Europe. *Journal of Biogeography*, **29**: 69–79.

- Jokimäki, J., Suhonen, J. (1998). Distribution and habitat selection of wintering birds in urban environments. *Landscape Urban Planning*, **39**: 253–263.
- Jumber, J. F. (1956) Roosting Behaviour of the Starling in Central Pennsylvania. *The Auk*, **73(3)**: 411-426.
- Kalisinska, E., Salicki, W., Myslek, P., Kavetska and K., Jackowski, A. (2004). Using the Mallard to biomonitor heavy metal contamination of wetlands in north-western Poland. *Science of the Total Environment*, **320(2)**: 145-161.
- Kang, N. and Yei, V.Y.Y. (1956) Roost site selection and waking and roosting behaviour of mynas in relation to light intensity. *Malayan Nature Journal*, **46**:255-263.
- Kark, S., Iwaniuk, A., Schalimtzek, A., Banker, E. (2007). Living in the city: can anyone become an “Urban exploiter?”. *Journal of Biogeographical research*, **34**: 638-651.
- Karr, J. R. (1982). Population variability and extinction in the avifauna of a tropical land bridge island. *Ecology*, **63**: 1975-1978.
- Karr, J. R. and Freemark, K. E. (1983). Habitat selection and environmental gradients: Dynamics in the “stable” tropics. *Ecology*, **64**: 1481- 1491.
- Karr, J. R., Schemskean, D. W., Brokaw, D. N. (1982). Temporal variation in the undergrowth bird community of a tropical forest.

- The ecology of a tropical forest* [Eds.] Leigh, E. G., Rand, A. S., and Windsor, D. M. Smithsonian Inst. Press, Washington, D.C. pp. 441-453
- Kelly, J. P. (1993). The effect of nest predation on habitat selection by dusky flycatchers in limber pine juniper woodland. *Condor*, **95**: 83- 93.
- Kim, E. Y., Ichihashi, H., Saeki, K., Atrashkevich, G., Tanabe, S., and Tatsukawa, R. (1996). Metal accumulation in tissues of seabirds from Chaun, Northeast Siberia, Russia. *Environmental Pollution*, **92**: 240-252.
- Klopfer, P. (1963). Behavioural aspects of habitat selection: the role of early experience. *Wilson Bulletin*, **75**: 15-22.
- Knight, R. L., and Cole, D. N. (1991). Effects of recreational activity on wildlife in wildlands. Transactions of the North American Wildlife and Natural Resources Conference **56**:238-247
- Korpimäki, E and Norrdahl, K. (1991). Numerical and functional responses of Kestrels, Short eared Owls, and Long-eared Owls to vole densities. *Ecology*, **72**: 814-82.
- Korpimäki, E. (1987). Dietary shifts, niche relationships and reproductive output of coexisting Kestrels and Long-eared Owls. *Oecologia*, **74**: 277-28.

- Korpimäki, E. and Norrdahl, K. (1989). Avian predation on mustelids in Europe 2: Impact on small mustelid and microtine dynamics - a hypothesis. *Oikos*, **55**: 273-276.
- Kostrzewa, A. and Kostrzewa, R. (1990). The relationship of spring and summer weather with density and breeding performance of the buzzard *Buteo buteo*, goshawk *Accipiter gentilis* and kestrel *Falco tinnunculus*. *Ibis*, **132**: 550-559.
- Kotler, B. P. (1985). Owl predation on desert rodents which differ in morphology and behavior. *Journal of Mammalogy*, **66**(4): 824-828.
- Kozulin, A. and Pavluschick, T. (1993). Content of heavy metals in tissues of mallards *Anas platyrhynchos* wintering in polluted and unpolluted habitats. *Acta Ornithologica*, **28**: 55-61.
- Kraaijeveld, K. (2008). Non-breeding habitat preference affects ecological speciation in migratory waders. *Naturwissenschaften*, **95**: 347-354.
- Kruess A, and Tschardtke, T. (1994). Habitat fragmentation, species loss, and biological control. *Science*, **264**: 1581-1584.
- Krüger, O. (2002). Analysis of nest occupancy and nest reproduction in two sympatric raptors: common buzzard *Buteo buteo* and Goshawk *Accipiter gentilis*. *Ecography*, **25**: 523-532.

- Kurta, A., Williams, K. J. and Mies. R. (1996). Ecological, behavioral, and thermal observations of a peripheral population of Indiana bats (*Myotis sodalis*). [Eds.] Barclay, R. M. R. and R. M. Brigham, *Bats and forests. British Columbia Ministry of Forests*. pp 102-117
- Kushlan, J. A. (1976). Site selection for nesting colonies by the American White Ibis *Eudocimus albus* in Florida. *Ibis*, **118**: 590-593.
- Lack, D. (1946). Competition for food in birds of prey. *Journal of Animal Ecology*, **15**: 123-129.
- Lack, D. (1954). *The natural regulation of animal numbers*. Oxford University Press, Oxford. pp 343.
- Lack, D. (1968). *Ecological adaptations for breeding in birds*. Methuen and Co., London. pp 409.
- Lambeck, R. J. (1997). Focal species: a multi-species umbrella for nature conservation. *Conservation Biology*, **11**: 849–856.
- Lancaster, R. K., Rees, W. E. (1979). Bird communities and the structure of urban habitats. *Canadian Journal of Zoology*, **57**: 2358–2368.
- Lande, R., Engen, S. and Saether, B.E. (2003). *Stochastic Population Dynamics in Ecology and Conservation*. Oxford University Press, Oxford.

- Levey, D. J. (1988). Spatial and temporal variation in Costa Rican fruit and fruit-eating bird abundance. *Ecological Monograph*, **58**: 251-269.
- Li, P. and Martin, T. E. (1991). Nest site selection and nesting success of cavity nesting birds in high elevation forest drainages. *Auk*, **108**: 405-418.
- Licata, P., Naccari, F., Lo Turco, V., Rando, R., Di Bella, G. and Dugo, G. (2010). Levels of Cd (II), Mn (II), Pb (II), Cu (II), and Zn (II) in Common Buzzard (*Buteo buteo*) from Sicily (Italy) by Derivative Stripping Potentiometry. *International Journal of Ecology*, **1**: 1-7.
- Limíñana, R., Javaloyes, T. and Urios, V. (2012). Diet of the Montagu's Harrier *Circus pygargus* nesting in natural habitat in Eastern Spain. *Ornis Fennica*, **89**: 74-80.
- Lundberg, P. (1988). The evolution of partial migration in birds. *Trends in Ecology and Evolution*, **3**: 172-175.
- Lyon, L.A. and Caccamise, D.F. (1981). Habitat selection by roosting blackbirds and starlings: management implications. *Journal of Wildlife Management*, **45**: 435 - 443.
- McGavin, G. C. (2000). *Insects, Spiders and other terrestrial arthropods*. A Dorling Kindersley Limited, 80 Strand, London.

- Mader, W. J. (1981). Notes on nesting raptors in the Llanos of Venezuela. *Condor*, **84**: 48-51.
- Mader, W. J. (1982). Ecology and breeding habits of the Savanna Hawk in the Llanos of Venezuela. *Condor*, **84**: 261-271.
- Maeda, T. (1998). Bird communities and habitat relationships in a residential area of Tokyo. *Journal of Yamashina Institute of Ornithology*, **30**: 83–100.
- Mahabal A. and Bastawade, D. B. (1987). Population ecology and communal roosting behavior of black kite *Milvus migrans govinda* in Pune (Maharashtra). *Journal of Bombay Natural History Society*. **82**: 337-346.
- Mahmood-ul-Hassan, M., Beg, M. A. and Ali, H. (2007). Seasonal variation in the diet of the barn owl *Tyto alba stertens* in central Punjab, Pakistan. *Acta Zoologica Sinica*, **53(3)**: 431–436.
- Makarieva, Anastassia M., Victor G. Gorshkov, and Bai-Lian Li. (2005). Why do population density and inverse home range scale differently with body size? Implications for ecosystem stability. *Ecological Complexity*, **2(3)**: 259-271.
- Mañosa, S. (1994). Goshawk diet in a Mediterranean area of Northeastern Spain. *Journal of Raptor Research*, **28**: 84-92.

- Mansouri, B., Babaei, H., Hoshyari, E., Khodaparast, S. H. and Mirzajani, A. (2012). Assessment of Trace-Metal Concentrations in Western Reef Heron (*Egretta gularis*) and Siberian Gull (*Larus heuglini*) From Southern Iran. *Archieves of Environmental Contamination and Toxicology*, **63**:280–287.
- Margules, C. R. and Pressey, R.L. (2000). Systematic conservation planning. *Nature*, **405**: 243–253.
- Marti, C. D. (1974). Feeding ecology of four sympatric owls. *Condor*, **76(1)**: 45-61.
- Marti, C. D., Bechard, M. and Jacksic, F. M., (2007). *Food habits*. Bird, D.M., Bildstein, K.L. [Eds.], Raptor Research and Management Techniques. Hancock House Publishers, Surrey, BC., pp 129-149.
- Marti, C.D. (1987). *Raptor food habit studies* National Wildlife Federation, Washington, D.C.
- Martin, T. E. (1988). Habitat and area effects on forest bird assemblages: is nest predation an influence? *Ecology*, **69(1)**:74-84.
- Martin, T. E. (1993). Nest predation among vegetation layers and habitat types: revising the dogmas. *American Naturalist*, **141**: 897–913.
- Martinez, J. A , Martinez, J. E. Zuberogotitia, I., Garcia, J. T., Caarbonell, R., de Lucas, M. and Diaz, M. (2003). Environmental impact

- assessment on raptor populations: difficulties in implementation and a search for solution. *Ardeola*, **50**: 85-102.
- Marzluff, J. M., Bowman, R., Donnelly, R. (2001). *Avian ecology and conservation in an urbanizing world*. Kluwer Academic Publ. Boston.
- Marzluff, J. M. (2001). Worldwide urbanisation and its effects on birds.
- Marzluff, J. M., Bowman, R., Donnelly, R. [Eds.], *Avian Ecology and Conservation in an Urbanizing World*. Kluwer Academic Publishers, Boston, pp 19–47.
- Marzluff, J. M. and Ewing, K. (2001). Restoration of fragmented landscapes for the conservation of birds: a general framework and specific recommendations for urbanizing landscapes. *Restoration Ecology*, **9**: 280–292.
- Marzluff, J. M. and Hamel, N. (2001). Land use issues. Levin, S. A. [Eds.], *Encyclopedia of biodiversity*. Academic Press, San Diego, CA. **3**: 659-673
- Matlack, G. R. (1993). Sociological edge effects: spatial distribution of human impact in suburban forest fragments. *Environmental Management*, **17**: 829-835.
- May, C. A., Petersburg, M. L. and Gutierrez, R. J. (2004) Mexican spotted owl nest and roost site habitat in northern Arizona. *Journal of Wildlife Management*, **68**: 1054- 1064.

- May, R. (1999). Unanswered questions in ecology. *Philosophical Transactions: Biological Sciences*, **354**: 1951–1959.
- Mearns, R. and Newton, I. (1988). Factors affecting breeding success of Peregrines in south Scotland. *Journal of Animal Ecology*, **57**: 903-916.
- Meffe, G. K. and Carroll, C. R. (1994). *Principles of Conservation Biology*. Sinauer Associates, Sunderland, Massachusetts.
- Mehra, R., Thakur, A. S. and Bhalla, S. (2010). Trace level analysis of chromium, manganese, cobalt and iron in human hair of people residing near heavy traffic area by biomonitoring. *International Journal of Pharma and Bio Science*, **1 (4)**: 57- 61.
- Melles , S., Glenn, S. and Martin, K. (2003). Urban bird diversity and landscape complexity: species-environment associations along a multi-scale habitat gradient. *Ecology and Society*, **7(1)**: 5-11.
- Mellone U., Agostini N., Panuccio M., Gustin M. (2007). Spring raptor migration over the Eolian archipelago. Proceeding -XIV Italian Ornithological Congress: **67**.
- Menge, B. A., Berlow, E. L., Blanchette, C. A., Navarrete, S. A. and Yamada. S. B. (1994). The keystone species concept: variation in interaction strength in a rocky intertidal habitat. *Ecological Monographs*, **64**:249–286

- Meretsky, V. J., Snyder, N. F., Beissinger, S. R., Clendenen, D. A. and Wiley, J. W., (2000). Demography of the California condor: implications for reestablishment. *Conservational Biology*, **14**: 957–967.
- Meyer, W. B and Turner, B.L. (1992). Human population growth and global land-use/cover change. *Annual Review of Ecology and Systematics*, **23**:39-61.
- Middleton J. (1994). Effects of urbanization on biodiversity in Canada. In: Biodiversity in Canada: A Science Assessment for Environment Canada. Environment Canada, Ottawa, pp. 15-20
- Miller, J. R., and Hobbs, R. J. (2002). Conservation where people live and work. *Conservational biology*. **16**: 330–337.
- Mills, G. S, Dunning, J. B. and Bates, J.M. (1989). Effects of urbanisation on breeding bird community structure in southwestern desert habitats. *Condor*, **91**: 416–429.
- Morneau, F., Décarie, R., Pelletier, R., Lambert, D., DesGranges, J. L., Savard, J.P. (1999). Changes in breeding bird richness and abundance in Montreal parks over a period of 15 years. *Landscape Urban Planning*, **44**: 111–121.
- Morse, D. H. (1980). *Behavioural mechanisms in ecology*. Harvard University Press, Cambridge, Massachusetts. pp 383.

- Mosher, J. A. and White, C. M. (1976). Directional exposure of golden eagle nests. *Canadian Field Naturalist*, **90**: 356-359.
- Munyenyembe, F., Harris, J. and Hone, J. (1989). Determinants of bird populations in an urban area. *Australian Journal of Ecology*, **14**: 549–557.
- Mushtaq-ul-Hassan, M., Raza, M. N., Shahzadi, B. and Ali, A. (2004). The diet of Barn Owl from canal bank, canal rest house and graveyard of Gogra. *Journal of Research (Science)*, **15(3)**: 291–296
- Naidu, B. (2008). *Diversity, Ecology and Conservation of Insect: A habitat Approach*. Ph.D. Thesis submitted to The M. S. University of Baroda. Vadodara, India.
- Naoroji, R. (2007). *Birds of prey of the India Subcontinent*. Om books international. Delhi.
- Natuhara, Y. and Imai, C. (1999). Prediction of species richness of breeding birds by landscape level factors of urban woods in Osaka Prefecture, Japan. *Biodiversity and Conservation*, **8**: 239–253.
- Newton, I. (1979). *Population ecology of Raptors*. Poyser, Berkhamsted. pp 399.

- Newton, I. (1982). Fidelity to breeding area and mate in Sparrow Hawks (*Accipiter nisus*). *Journal of Animal Ecology*, **51**: 327-341.
- Newton, I. (1991). Habitat variation and population regulation in Sparrow Hawks. *Ibis*, **133**: 76- 88.
- Newton, I. (2004). Population limitation immigrants. *Ibis*, **146**: 197–226.
- Nicholson, J. K. (1981). The comparative distribution of zinc, cadmium and mercury in selected tissues of the herring gull (*Larus argentatus*). *Comparative Biochemistry and Physiology Part C: Comparative Pharmacology*, **68 (1)**: 91-94.
- Niemelä, J. (1999a). Ecology and urban planning. *Biodiversity and Conservation*, **8**: 119–131.
- Nilsson, S. G. (1987). Limitation and regulation of population density in the Nuthatch *Sitta europaea* breeding in natural cavities. *Journal of Animal Ecology*, **56**: 921-937.
- Nordberg, G.F, Nogawa, K., Nordberg, M. and Friberg, L. (2007). *Cadmium. In: Handbook on The Toxicology of Metals* [Eds.] Nordberg, G. F., Fowler, B. F., Nordberg, M. and Friberg, L. Amsterdam: *Elsevier*, pp 445-448.
- Norris, D. R. and Marra, P. P. (2007). Seasonal interactions, habitat quality, and population dynamics in migratory birds. *Condor*, **109**: 535–547.

- O'Donoghue, M., Boutin, S., Murray, D. L., Krebs, C. J., Hofer, E. J. and Breitenmoser, U. L. (2001). Coyotes and lynx. C. J. Krebs, S. Boutin & R. Boonstra [Eds.], *Ecosystem dynamics of the boreal forest*. New York: Oxford University Press.
- Olga, C. and Jose, A. D. (1989). Factors influencing the breeding density and nest site selection of the Egyptian Vulture *Neophron percnopterus*. *Journal of ornithology*, **130(5)**: 353-359.
- Olojo, E. A. A., Olurin, K. B., Mbaka, G. and Oluwemimo, A. D. (2005). Histopathology of the gill and liver tissues of the African catfish *Clarias gariepinus* exposed to lead. *African Journal of Biotechnology*, **4**: 117-122.
- Olsen, P. D. and Olsen, J. (1989a). Breeding of the Peregrine Falcon *Falco peregrinus*. II. Weather, nest quality and the timing of egg laying. *Emu*, **89**: 1-5.
- Ontiveros, D. (1999). Selection of nest cliffs by bonelli's eagle (*hieraaetusfa sdatus*) in southeastern spain. *Raptor Research*, **33(2)**:110-116.
- Orth, P. B. and Kennedy, P. L. (2001). Do land-use patterns influence nest-site selection by Burrowing owls (*Athene cunicularia hypugaea*) in northeastern Colorado? *Canadian Journal of Zoology*, **79**: 1038-1045.

- Outridge, P. M., and Scheuhammer, A. M. (1993). Bioaccumulation and toxicology of chromium: implications for wildlife. *Reviews of environmental contamination and toxicology*, **130**: 31-77.
- Pain, D. J., Carter, I., Sainsbury, A.W., Shore, R.F., Eden, P., Taggert, M. A., Konstantions, S., Walker, L. A., Meharg, A. A. and Raab, A. A. (2007). Lead contamination and associated disease in captive and reintroduced red kite *Milvus milvus* in England. *Science of the Total Environment*, **376**: 116-127.
- Pain, D. J. (1992). Lead poisoning in waterfowl: a review. Pain D. J. [Eds.]. *Lead Poisoning in Waterfowl*. Proceedings of an IWRB Workshop, Brussels, Belgium, 13–15 June 1991. International Waterfowl and Wet- lands Research Bureau Special Publication 16, Slimbridge, UK. pp 7–13.
- Pain, D. J. (1996). Lead in waterfowl. Beyer, W. M., Heinz, G. H. and Redman-Norwood, A. W. [Eds.] *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*. Lewis Publishers, Boca Raton, Florida, USA. pp 251– 262.
- Pain, D. J., Carter, I., Sainsbury, A.W., Shore, R.F., Eden, P., Taggert, M. A., Konstantions, S., Walker, L. A., Meharg, A. A. and Raab, A. A. (2007). Lead contamination and associated disease in captive

- and reintroduced Red Kite *Milvus milvus* in England. *Science of the Total Environment*, **376**: 116-127.
- Paine, R. T. (1980). Food webs: road maps of interactions or grist for theoretical development? *Ecology*, **69**: 1648-1654.
- Paletto, A. and Tosi V. (2009). Forest canopy cover and canopy closure: comparison of assessment techniques *European Journal of Forest Research*, **128**:265–272.
- Palomino, D. and Carrascal, L. M. (2007). Threshold distances to nearby cities and roads influence the bird community of a mosaic landscape. *Biological Conservation*, **140**(1): 100-109.
- Panasci, T. A. and Whitacre, D. F. (2002). Roadside hawk breeding ecology in forest and farming landscapes. *Willson Bulletin*, **114**: 114–121.
- Panuccio M., Agostini N. and Massa B. (2004). Spring raptor migration at Ustica, southern Italy. *British Birds*, **97**: 400-403.
- Panuccio, M., Agostini, N. and Mellone, U. (2005). Autumn migration strategies of Honey Buzzards, Black Kites, Marsh and Montagu's Harriers over land and over water in the Central Mediterranean. *Avocetta*, **29**: 27-32.

- Panuccio, M., Agostini, N., D'Amicis, B., Canale, E., and Roccella, A. (2005). Sex and age ratios of Marsh Harriers *Circus aeruginosus* wintering in central-southern Italy. *Avocetta*, **29**:13–17
- Park, C. R and Lee, W. S. (2000). Relationship between species composition and area in breeding birds of urban woods in Seoul, Korea. *Landscape Urban Planning*, **51**:29–36.
- Park, C. R., Lee, W. S. (2000). Relationship between species composition and area in breeding birds of urban woods in Seoul, Korea. *Landscape Urban Plannning*, **51**: 29–36.
- Peh K. S. H. (2002) Roosting behaviour of House crow (*Corvus splendens*) in relation to envriornmental variables. *The Raffles Bulletin of Zoology*, **50(1)**: 257-262.
- Penland, S. (1987). Attitudes of urban residents toward avian species and species' attributes. Adams, L.W. and Leedy, D.L. [Eds.] *Integratineg Man and Nature in the Metropolitan Environment*, Chevy Chase, New York, pp 77-82.
- Pickett, S. T., Cadenasso, M. L., Grove, J. M., Nilon, C. H., Pouyat, R. V., Zipperer, W. C. and Rcostanza. A. (2001). Urban ecological systems: linking terrestrial ecological, physical, and socioeconomic components of metropolitan areas. *Annual Review of Ecology and Systematics*, **32**:127-157.

- Pimm, S. L. (1982). *Food Webs*. Chapman and Hall, London.
- Pimm, S. L., Gilpin, M. E. (1989). Theoretical issues in conservation biology. In: May, R.M., Levin, S.A., Roughgarden, J. [Eds.], *Perspective in Ecological Theory*. Princeton University Press, Princeton, New Jersey, pp 287-305.
- Polis, G. A. (1991). Complex desert food webs: an empirical critique of food web theory. *American Naturalist*, **138**:123-155.
- Porneluzi, P., Bednarz J. C., Goodrich L. J, Zawada N. and Hoover, J. (1993). Reproductive performance of territorial Ovenbirds occupying forest fragments and a contiguous forest in Pennsylvania. *Conservation Biology*, **7**:618–622.
- Porter, E. E., Forschner, B. R. and Blair, R. B. (2001). Woody vegetation and canopy fragmentation along a forest-to-urban gradient. *Urban Ecosystem*, **5**: 131–151.
- Powell, G. N. and Bjork, R. D. (1994). Implication of altitudinal migration for conservation strategies to protect tropical biodiversity: A case study of the Resplendent Quetzal *Pharomacrus mocinno* at Monteverde, Costa Rica. *Bird conservation international*, **4**: 161-174.
- Prakash V., Pain D. J., Cunningham A. A., Donald P. F., Prakah N., Verma A., Gargi, R., Shivakumar S. and Rahmani A. R. (2003).

- Catastrophic collapse of Indian White-backed *Gyps bengalensis* and Long-billed *Gyps indicus* vulture population. *Biological Conservation* **109**: 381-390.
- Probst, J. R., and Hayes, J. P. (1987). Pairing success of Kirkland's Warblers in marginal vs. suitable habitat. *Auk*, **104**:234–241.
- Pulido, F. (2007). The genetics and evolution of avian migration. *BioScience*, **57**: 165–174.
- Puschner, B., J., Leger, S. T. and Galey, F. D. (1999). Normal and toxic zinc concentrations in serum/plasma and liver of psittacines with respect to genus differences. *Journal of Veterinary Diagnostic Investigation*, **11** (6): 522–527.
- Pyle, G. G., Rajotte, J. W. and Couture, P. (2005). Effects of industrial metals on wild fish populations along a metal contamination gradient. *Ecotoxicology and Environment Safety*, **61**: 287-312.
- Rajkowska, M. and Protasowicki, M. (2013). Distribution of metals (Fe, Mn, Zn, Cu) in fish tissues in two lakes of different trophic in Northwestern Poland. *Environment Monitoring Assessment*, **185**: 3493–3502.
- Rathod, J. and Padate, G. (2004). *Feeding adaptation of Black kite (Milvus migrans govinda) in urban area of Vadodara*. Paper presented at International conference of Birds and Environment Organized by Gurukul Kangri University, Haridwar, India.

- Rathod, J. Y. (2009). *Avifauna of Urban areas: its significance and implication under various human disturbances*. Ph.D. Thesis submitted to The M. S. University of Baroda. Vadodara, India.
- Real, J. (1996). Biases in diet study methods in the Bonelli's Eagle. *Journal of Wildlife Management*, **60**: 632–638.
- Redpath, S. M., R. Clarke, M. Madders and S. J. Thirgood. (2001). Assessing raptor diet: comparing pellets, prey remains, and observational data at Hen Harrier nests. *Condor*, **103**: 184-188.
- Reebs, S. G. (1986). Influence of temperature and other factors on the daily roosting times of black-billed magpies. *Canadian Journal of Zoology*, **64**: 1614 -1619.
- Rendell, W. B. and Robertson, R. J. (1989). Nest site characteristics, reproductive success and cavity availability for tree swallows breeding in natural cavities. *Condor*, **91**: 875-885.
- Ricklefs, R. E. (1969). An analysis of nesting mortality in birds. *Smithsonian Contributions to Zoology*, **9**: 1-48.
- Rodríguez-Estrella, R., Donázar, J.A., Hiraldo, F., 1998. Raptors as indicators of environmental change in the scrub habitat of Baja California Sur, Mexico. *Conservation Biology*, **12**: 921–925.
- Root, T. (1988). Energy constraints on avian distributions and abundance. *Ecology*, **69**: 330-339.

- Rosenberg, K. V. and Cooper, R. J. (1990). Approaches to avian diet analysis. *Avian Biology*, **13**: 80- 90.
- Rosenzweig, M. L. (2003). *Win-win ecology*. Oxford University Press, Oxford, UK.
- Roshier, D. A., Robertson, A. I., Kingsford, R. T. (2002). Responses of waterbirds to flooding in an arid region of Australia and implications for conservation. *Biological Conservation*, **106**: 399–411
- Ruddock, M., Wells, J. H., Smith, J. P. and Montgomery, I. W. (2009). The biases in raptor diet studies; resolving conservation and sporting conflicts. Sielicki, J. and Mizera, T. [Eds.], *Peregrine Falcon Populations Status and Perspectives in the 21st Century*. Turul/Poznan University of Life Sciences Press, Warsaw Pozna. pp 567-580.
- Rutz, C. (2003). Assessing the breeding season diet of Goshawks *Accipiter gentilis*: biases of plucking analysis quantified by means of continuous radio-monitoring. *Journal of Zoology*, **259**: 209-217.
- Ryder, P. L. and Ryder, J. R. (1981). Reproductive performances of Ring billed Gulls in relation to nest location. *Condor*, **83**: 57-60.

- Sánchez-Zapata, J. and José, F. (1999). Raptor distribution in relation to landscape composition in semi-arid Mediterranean habitats. *Journal of Applied Ecology*, **36**(2):254-262
- Sánchez-Zapata, J.A., Carrete, M., Gravilov, A., Sklyarenko, S., Ceballos, O., Donazar, J.A., Hiraldo, F. (2003). Land use changes and raptor conservation in Steppe habitats of Eastern Kazakhstan. *Biological Conservation*, **111**: 71–77
- Sanderson, G. C. and Bellrose, F. C. (1986). A review of the problem of lead poisoning in waterfowl. *Linois Natural History Survey, Special Publication*, **4**:1–34.
- Sandstead H. H and Au, W. (2005). *Handbook on Toxicology of metals*. 3<sup>rd</sup> Edition. Academic Press, Denmark. pp 925 – 946.
- Sandvik, H. and Barrett. R. T. (2001) Effect of investigator disturbance on the breeding success of the black-legged kittiwake. *Journal of Field Ornithology*, **72**:30–42.
- Santhanakrishnan, R., Ali, M. S. and Anbarsan, U. (2010). Diet Variations of the Barn Owl *Tyto alba* in Madurai district, Tamilnadu, Southern India. *Podoces*, **5**(2):95-103.
- Sarà, M., and Vittorio, M. (2003). Factors influencing the distribution, abundance and nest-site selection of an endangered Egyptian

- Vulture (*Neophron percnopterus*) population in Sicily. *Animal Conservation*, **6(4)**: 317-328.
- Sarasola, J. H. and Negro, J. J. (2006) Role of exotic tree stands on the current distribution and social behaviour of Swainson's Hawk, *Buteo swainsoni* in the Argentine. *Journal of Biogeography*, **33**: (6): 1096-1101.
- Sauvajot, R. M., Buechner, M., Kamradt, D. A. and Schonewald, C.M. (1998). Patterns of human disturbance and response by small mammals and birds in chaparral near urban development. *Urban Ecosystems*, **2**: 279– 297.
- Savard, J. P., Clergeau, P. and Mennechez, G. (2000). Biodiversity concepts and urban ecosystems. *Landscape and Urban Planning*, **48**: 131-142.
- Scheuhammer, A. M., and Norris, S. L. (1996). The ecotoxicology of lead shot and lead fishing weights. *Ecotoxicology*, **5(5)**: 279-295.
- Schoener, T. W. (1989). Food webs from the small to the large. *Ecology*, **70**:1555-1559.
- Schreiber, R.W. (1967) Roosting Behavior of the Herring Gull in Central Maine. *Wilson Bulletin*, **79 (4)**: 421-431.

- Schwartz, K., and Mertz, W. (1959). Chromium (III) and glucose tolerance factor. *Archives of Biochemistry and Biophysics*, **85**: 292–295.
- Selås, V. (1993). Selection of avian prey by breeding Sparrowhawks *Accipiter nisus* in south Norway: the importance of size and foraging behaviour of prey. *Ornis Fennica*, **70**: 144-154.
- Selås, V., Tveiten, R. and Aanonsen, O. M. (2007). Diet of Common Buzzards (*Buteo buteo*) in southern Norway determined from prey remains and video recordings. *Ornis Fennica*, **84**: 97-104.
- Sergio, F., Caro, T., Brown, D., Clucas, B., Hunter, J., Ketchum, J., McHugh, K. and Hiraldo, F. (2008). Top predators as conservation tools: ecological rationale, assumptions and efficacy. *Annual Review of Ecology Evolution and Systematics*, **39**:1-19.
- Sergio, F., Newton, I., Marchesi, L. and Pedrini, P. (2006). Ecologically justified charisma: Preservation of top predators delivers biodiversity conservation. *Journal of applied ecology*, **43**:1049-1055
- Serrano, D., Forero, M.G., donázar, J.A. and Tella, J. L. (2004). Dispersal and social attraction affect colony selection and dynamics of lesser kestrels. *Ecology*, **85(12)**:3438–3447.
- Sevin, I. F. (1980). *Nickel- Metals in the environment*. HA Waldron. pp 263-291.

- Simberloff, D. (1988). The contribution of population and community biology to conservation science. – *Annual review of ecology, evolution and systematic*, **19**: 473-511.
- Simmons, R. and Smith, P.C. (1985). Do Northern Harriers (*Circus cyaneus*) choose nest sites adaptively? *Canadian Journal of Zoology*, **63**: 494-498.
- Simmons, R. E., Avery, D. M. and Avery, G. (1991). Biases in diets determined from pellets and remains: correction factors for a mammal and bird-eating raptor. *Journal of Raptor Research*, **25**(3): 63-67.
- Sinclair, K. E., Hess, G. R., Moorman, C. E. and Mason, J. M. (2005). Mammalian nest predators respond to greenway width, landscape context, and habitat structure. *Landscape and Urban Planning*, **71**:277-293.
- Singh, A. P. and Srivastava, P. C. (2007). Relationships of heavymetals in natural lake waters with physic chemical characteristics of waters and different chemical fractions of metals in sediments. *Water, Air, and Soil Pollution*, **188**: 181–193.
- Singh, R. K., Chavan, S. L. and Sapkale, P. H. (2007). Heavy metal concentrations in water, sediments and body tissues of red worm (*Tubifex* spp.) collected from natural habitats in Mumbai, India. *Environmental Monitoring and Assessment*, **129**: 471-481.

- Skutch, A.F. (1976). *Parent Birds and Their Young*. University of Texas.
- Sodhi, N. S., Briffet, C., Kong, L. and Yuen, B. (1999). Bird use of linear areas of a tropical city: implications for park connector design and management. *Landscape Urban Planning*, **45**:123–130.
- Soni, K. C., Sharma, A. N. and Soni, V. C. (2010). Nesting Ecology, Interspecific Interaction and Nesting Association of Indian Black Ibis (*Pseudibis papillosa*) Inhabiting the Arid Zone of Rajasthan. *Our Nature*, **8**: 12-25.
- Soriano-Santos, J., & Guerrero-Legarreta, I. (2010). Chemical Composition and Nutritional Content of Raw Poultry Meat. *Handbook of Poultry Science and Technology*, John Wiley and Son, Inc, Hoboken, New Jersey.
- Speiser, R. and Bosakowski, T. (1987). Nest site selection by northern goshawks in northern new Jersey and Southeastern New York. *Condor*, **89**:387-394.
- Stanevièius, V. (2004). Nest-site selection by Marsh Harrier (*Circus aeruginosus*) in the shore belt of helophytes On large lakes. *Acta Zoologica Lituanica*, **14**(3): 47-53.
- Steenhof, K., Kochert, M. N., Carpenter, L. B. and Lehman, R. N. (1999) Long term prairie falcon population changes in relation to prey

- abundance, weather, land uses, and habitat conditions. *Condor*, **101**: 28–41
- Storelli, M. M., Barone, G., Storelli, A. and Marcotrigiano, G. O. (2006). Trace metals in tissues of mugilids (*Mugil auratus*, *Mugil capito*, and *Mugil labrosus*) from the Mediterranean Sea. *Bulletin of Environmental Contamination and Toxicology*, **77**: 43-50.
- Swaileh, K. M and Sansur, R. (2006). Monitoring urban heavy metal pollution using the House Sparrow (*Passer domesticus*). *Journal of Environmental Monitoring*, **8**: 209–213.
- Swingland, I. R. (1976) The influence of Light intensity on the roosting times of the rook (*Corvus frugilegus*). *Animal Behaviour*, **24**: 154-158.
- Szefer, P., Czarnowski, W., Pempkowiak, J. and Holm, E. (1993). Mercury and major essential elements in seals, penguins, and other representative fauna of the Antarctic. *Archived Environmental Contamination Toxicology*, **25**: 422–427.
- Szymczyk, K. and Zalewski, K. (2003). Copper, zinc, lead, and cadmium content in liver and muscle of Mallards (*Anas platyrhynchos*) and other Hunting Fowl Species in Warmia and Mazury in 1999- 2000. *Polish Journal of Environmental Studies*. **12(3)**: pp 381-386.

- Technical Guidance document on Risk Assessment. (2003). European Commission Joint research Centre.
- Tella, J.L., Forero, M.G., Hiraldo, F., Donázar, J.A., (1998). Conflicts between lesser kestrel conservation and European agricultural policies as identified by habitat use analyses. *Conservation Biology*, **12**: 593–604.
- Tester, J.R. (1987) Changes in daily activity rhythms of some free-ranging animals in Minnesota. *Canadian Field Naturalist*, **101**: 13-21.
- Thiollay, J M. (1989). Area requirement for the conservation of rain forest raptors and game birds in French Guiana. *Conservation biology*, **3**: 128-137
- Thorington, K. K. and Bowman, R. (2003). Predation rates on artificial nests increases with human housing density in suburban habitats. *Ecography*, **26**:188-196.
- Tom, D., Bervoets, L., Blust, R. and Eens, M. (2002). Tissue levels of lead in experimentally exposed Zebra finches (*Taeniopygia guttata*) with particular attention on the use of feathers as biomonitor. *Archived Environmental Contamination Toxicology*, **42**: 88-92.

- Tomialojc, L. (1998). Breeding bird densities in some urban versus non-urban habitats: the Dijon case. *Acta Ornithologica*, **33**: 159–171.
- Toumenpuro, J. (1991). Effect of nest site on nest survival in the Dunnock, *Prunella modularis*. *Ornis Fennica*, **68**: 49-56.
- Toyne, E. P. (1998). Breeding season diet of the Goshawk *Accipiter gentilis* in Wales. *Ibis*, **140**: 569-579.
- Tree, A. J. (2001). Kittlitz's Plover as an intra-African migrant. *Honeyguide*, **47**: 10–16.
- Tree, A.J. (1973). Ageing the Kittlitz's Plover. *Safring News*, **2(2)**: 23–25.
- U. S. Fish and Wildlife Service (1996). Canadian Wildlife Service. 1986. *North American waterfowl management plan: a strategy for cooperation*. US Fish and Wildlife Service, Washington, DC.
- Underhill, L.G., Tree, A.J., Oschadleus, H.D., Parker, V. (1999). *Review of ring recoveries of Waterbirds in southern Africa*. Cape Town: Avian Demography Unit, University of Cape Town.
- Urban, E.K. (1986). "Charadriinae, plovers" *The birds of Africa*, **2**: 225-246.
- Wackernagel, M. and Rees, W. (1996). *Our Ecological Footprint: Reducing Human Impact on the Earth*. The New Catalyst Bioregional Series, New Society Publishers.

- Walsh, C. T., Sandstead, H. H., Prasad, A. S., Newberne, P. M. and Fraker, P. J. (1994) Zinc: health effects and research priorities for the 1990's. *Environment Health Perspecive*. **102(2)**: 5–46.
- Walters, J. R. (2000). Dispersal behavior: an ornithological frontier. *Condor*, **102**:479–481.
- Watling, J.I., Nowakowski, J.A., Donnelly, M.A. and Oroock, J. L. (2009). Meta-analysis reveals the importance of matrix composition for animals in fragmented habitat. *Global Ecology and Biogeography*, **20**: 209–217.
- Watson, J. (1992). Nesting ecology of the Seychelles Kestrel *Falco araea* on Mahe, Seychelles. *Ibis*, **134**: 259-267.
- Welty, J. C and Baptista, L. F. (1988). *The life of birds*, 4<sup>th</sup> Edition. WB Saunders, New York.
- Whellwright, N. T. (1983). Fruits and the ecology of resplendent Quetzal. *Auk*, **100**: 286-301.
- White, J. G., Amos, M. J., Fitzsimons, J. A. and Palmer, G. C. (2005). Non-uniform bird assemblages in urban environments: the influence of streetscape vegetation. *Landscape and Urban Planning*, **71**: 123-135.
- White, T.C.R. (2008). The role of food, weather and climate in limiting the abundance of animals. *Biological Reviews*, **83**: 227–248.

- Widen, P. (1994). Habitat quality for raptors: a field experiment. *Journal of Avian Biology*, **25**: 219–223.
- Wiemeyer and Stanley N. (1988). Environmental contaminants in California condors. *The Journal of Wildlife Management*, **52**: 238–247.
- Wiens, J. A. (1976). Population responses to patchy environments. *Annual Review of Ecology, Evolution and Systematic*, **7**: 81–120.
- Wild life Protection Act. (1972). Amendment, 2008.
- Wilson, M.W., Irwin, S., Norriss, D.W., Newton, S.F., Collins, K., Kelly, T.C. and O'Halloran, J. (2009). The importance of pre-thicket conifer plantations for nesting Hen Harriers *Circus cyaneus* in Ireland. *Ibis*, **151**: 332–343.
- Wilson, R. P. and Culik, B. (1995). Penguins disturbed by tourists. *Nature*, **376**:301–302.
- World Health Organization. (1991). *International programme on chemical safety*. Environmental health criteria: Nickel. WHO, O'Day, P. A., Chisholm Brause, C. J., Towle, S.N., Parks, G. A. and Geneva. G. E.
- World Resources Institute. (1996). World Resources 1996–1997. Oxford University Press, Oxford.

- Xu, Y, Ran. J, Zhou. X, Yang. N., Yue. B. and Wang. Y. (2008) The effect of temperature and other factors on roosting times of Szechenyi Monal Partridges *Tetraophasis szechenyii* during the Breeding season. *Ornis Fennica*, **85**: 126-134.
- Zammuto, R.M. and Franks, E.C. (1981) Environmental effects on roosting behaviour of Chimney Swifts. *Wilson Bulletin*, **93**: 77-84.
- Zhuang, P., Zou, H., and Shu, W. (2009). Biotransfer of heavy metals along a soil, plant, insect, chicken food chain: Field study. *Journal of Environmental Science.*, **21(6)**: 849–853.