

## FIGURES

Figure No	Title	Page No.
<b>Chapter 1</b>		
<b>Introduction</b>		
Figure 1.1	Global distribution of phosphorus in soil	2
Figure 1.2	The P cycle in soil	3
Figure 1.3	Plant- microbe mechanism to increase phosphorus availability in rhizosphere	6
Figure 1.4	Molecular mechanism of P solubilisation by direct oxidative pathway	12
Figure 1.5	Schematic representation of the physiological and chemical processes	15
Figure 1.6	Colonization of the root of <i>Arabidopsis thaliana</i> by <i>B. subtilis</i> biofilms limiting the outer space for <i>Pseudomonas syringae</i> to infection	30
Figure 1.7	Molecular Strategies for the development of phosphate biofertilizers	32
<b>Chapter 2</b>		
<b>Substrate specificity of Glucose dehydrogenase (GDH) of <i>Enterobacter asburiae</i> PSI3 and RP solubilisation with GDH substrates as C sources</b>		
Figure 2.1	RP solubilisation by <i>E. asburiae</i> PSI3 on buffered methyl red plates	43
Figure 2.2	Growth profile and acidification pattern of <i>E. asburiae</i> PSI3 on Tris RP minimal medium with 75mM and 50mM glucose concentration respectively	44
Figure 2.3	Growth profile and acidification pattern of <i>E. asburiae</i> PSI3 on Tris RP minimal medium with 15mM and 10mM sugar mixture respectively	44
Figure 2.4	Organic acid analysis of <i>E. asburiae</i> PSI3 on different GDH substrates	46
<b>Chapter 3</b>		
<b>Effect of the presence of plasmids on the P-solubilization by <i>E. asburiae</i> PSI3</b>		
Figure 3.1	Restriction maps of the plasmids used in this study	54
Figure 3.2	Schematic representation of construction of plasmid pBSK-Gm and pBR322-Gm	59

Figure No.	Title	Page No.
Figure 3.3	Restriction endonuclease analysis of pBSK-Gm and pBR322-Gm	60
Figure 3.4	Restriction map of (A) pBSK-Gm and (B) pBR322-Gm	60
Figure 3.5	Acidification of methyl red plates by <i>E. asburiae</i> PSI3 containing different plasmids on Tris-Cl containing minimal media with RP (A,C,E) and $\text{KH}_2\text{PO}_4$ (B,D,F) as a sole P source (1) <i>E. a</i> (pBSK-Gm) (2) <i>E. a</i> (pBR322-Gm) (3) <i>E. a</i> (pGM160) (4) <i>E. a</i> (pTOL) (5) <i>E. a</i> (pACYC184)	62
Figure 3.6	Growth profile (A) and acidification pattern (B) of <i>E. asburiae</i> PSI3 containing different plasmids on Tris RP minimal media	63
Figure 3.7	Growth profile (A) and acidification pattern (B) of <i>E. asburiae</i> PSI3 containing different plasmids on Tris minimal media containing 10mM of $\text{KH}_2\text{PO}_4$	64
Figure 3.8	Growth profile and acidification pattern of <i>E. asburiae</i> PSI3 containing different plasmids on Tris minimal media containing 100 $\mu\text{M}$ of $\text{KH}_2\text{PO}_4$	65
<b>Chapter 4</b>	<b>Overexpression of <i>Escherchia coli appA</i> gene in different rhizobacteria Effect of acidification on phytate utilization and growth promotion of maize</b>	
Figure 4.1	Model of plant growth promotion by rhizobia in legume–cereal cropping systems	73
Figure 4.2	Restriction map of pBBRMCS-2	80
Figure 4.3	Restriction map of pTZ57R/T	81
Figure 4.4	Schematic representation of construction of plasmid pVA1	88
Figure 4.5	Restriction endonuclease analysis of pVA1	88
Figure 4.6	Restriction map of pVA1 plasmid	89
Figure 4.7	Phytase activity of transconjugants	89
Figure 4.8	Acid phosphatase activity of transconjugants	90
Figure 4.9	Growth and acidification profile of <i>E. a</i> (pVA1) and <i>E. a</i> (pBBRMCS-2) on buffered (A) and unbuffered (B) minimal containing NaP and CaP	91
Figure 4.10	Growth and acidification profile of <i>P. p</i> (pVA1) and <i>P. p</i> (pBBRMCS-2) on buffered (A) and unbuffered (B) minimal containing NaP and CaP	92

Figure No.	Title	Page No.
Figure 4.11	Growth and acidification profile of <i>S. m</i> (pVA1) and <i>S. m</i> (pBBRMCS-2) on buffered (A) and unbuffered (B) minimal containing Nap and CaP	94
Figure 4.12	Western blot of AppA from Different strains harboring <i>E. coli appA</i>	96
Figure 4.13	Effect of <i>S. meliloti appA</i> transconjugant inoculation on maize ( <i>Zea mays</i> ) growth	98
<b>Chapter 5</b>	<b>Effect of heterologous overexpression of <i>cs</i> and <i>ppc</i> genes in <i>Enterobacter asburiae</i> PSI3</b>	
Figure 5.1	Restriction maps of various plasmids used in this study	110
Figure 5.2	Schematic representation of construction of plasmid pVSppc	114
Figure 5.3	Restriction digestion pattern of plasmid pVSppc	115
Figure 5.4	Restriction map of pVSppc	115
Figure 5.5	Phenotypic complementation of <i>E. coli</i> (JWK3928) by pVSppc plasmid	116
Figure 5.6	SDS-PAGE gel of expression of the <i>ppc</i> gene	117
Figure 5.7	Growth profile of <i>E. asburiae</i> PSI3 containing <i>cs</i> and <i>ppc</i> genes alone or in combination with their respective controls on glycerol as sole carbon source	120
Figure 5.8	Growth profile of <i>E. asburiae</i> PSI3 containing <i>cs</i> and <i>ppc</i> genes alone or in combination with their respective controls on fructose as a sole carbon source	121
Figure 5.9	P solubilisation of <i>E. asburiae</i> PSI3 transformants on Tris minimal medium with TCP as sole P source and glycerol and fructose as sole carbon source	123