

TABLES

Tables No	Title	Page No.
Chapter 1	Introduction	
Table 1.1	Nature of Organic acid secreted by PSMs	8
Table 1.1 A	Nature of organic acid secreted by PS fungi and Actinomycetes	8
Table 1.1 B	Nature of organic acids secreted by PS bacteria	9
Table 1.2	Cloning of genes involved in MPS phenotype	12
Table 1.3	List of soil bacteria capable of P release from lower forms of inositol phosphates	16
Table 1.4	List of soil bacteria capable of P release from phytase	18
Table 1.5	Single bacterial inoculation field studies on various crops	20
Table 1.6	<i>P. billai</i> inoculation field studies on various crops	21
Table 1.7	Co-inoculation studies of bacterial PSMs with fungal PSMs or multiple bacterial inoculants	22
Chapter 2	Substrate specificity of Glucose dehydrogenase (GDH) of <i>Enterobacter asburiae</i> PSI3 and RP solubilisation with GDH substrates as C sources	
Table 2.1	GDH activity of <i>E. asburiae</i> PSI3	41
Table 2.2	Solubilisation of RP by <i>E. asburiae</i> PSI3 on different carbon sources	45
Table 2.3	Comparison of the substrate specificity of GDH enzymes from various bacteria	47
Chapter 3	Effect of the presence of plasmids on the P-solubilization by <i>E. asburiae</i> PSI3	
Table 3.1	Bacterial strains and plasmids used in this study	52-53
Table 3.2	Plasmids used in this study and their relevant properties	58
Table 3.3	P solubilization by <i>E. asburiae</i> PSI3 containing various plasmids	61
Table 3.4	Physiological variables of <i>E. asburiae</i> PSI3 and its various transformants grown on Tris RP medium with 100μM of KH ₂ PO ₄ and 100mM glucose	66

Table No.	Title	Page No.
Table 3.5	Organic acid yield and phosphate release of <i>E. asburiae</i> PSI3 and its various transformants grown on Tris RP medium with 100 μ M of KH ₂ PO ₄ and 100mM glucose.	67
Table 3. 6	Stability of pACYC184 plasmid in <i>E. asburiae</i> PSI3 in absence of antibiotic selection	67
Chapter 4	Overexpression of <i>Escherchia coli appA</i> gene in different rhizobacteria - Effect of acidification on phytate utilization and growth promotion of maize	
Table 4.1	Rhizobial molecules with potential for plant growth promotion	75
Table 4.2	Transgenic plants express different phytase gene for utilizing phytate P	77
Table 4.3	Organisms producing phytase used for P availability to different plants from Phytate	78
Table 4.4	Bacterial strains and plasmids used in this study	79
Table 4.5	P-release by transconjugants on buffered and unbuffered NaP and CaP media	95
Table 4.6	Effect of <i>S. meliloti appA</i> transconjugant inoculation on maize (<i>Zea mays</i>) growth parameters	97
Chapter 5	Effect of heterologous overexpression of <i>cs</i> and <i>ppc</i> genes in <i>Enterobacter asburiae</i> PSI3 gene	
Table 5.1	Efficacy of different organic acids at P-solubilization from alfisol supplemented with RP (40 mg g ⁻¹ soil) and alkaline vertisol	103
Table 5.2	Bacterial strains and plasmids used in this study	107
Table 5.3	PPC activity of <i>E. asburiae</i> PSI3 transformants	118
Table 5.4	CS activity of <i>E. asburiae</i> PSI3 transformants	118
Table 5.5	Physiological variables of <i>E. asburiae</i> PSI3 and its various transformants grown on M9 medium with 100mM glycerol	119
Table 5.6	Physiological variables of <i>E. asburiae</i> PSI3 and its various transformants grown on M9 medium with 100mM Fructose	122
Table 5.7	pH profile of <i>E. asburiae</i> PSI3 transformants	122