

## LIST OF TABLES

### CHAPTER 1

Table 1.1:	Historical landmark in GM2 gangliosidosis
Table 1.2:	Review of molecularly proven cases of TSD
Table 1.3:	Review of molecularly proven cases of AB variant form of TSD (GM2 activator protein deficiency)

### CHAPTER 3

Table 3.1:	Buffers and reagents used
Table 3.2:	Primers used for Tay-Sach's common mutation
Table 3.3:	PCR conditions for common mutations
Table 3.4:	Primers for sequencing of <i>HEXA</i> Gene
Table 3.5:	Primers for sequencing of <i>GM2A</i> gene

### CHAPTER 4

Table 4.1:	Hexosaminidase-A (Hex-A) and total-Hexosaminidase (total-Hex) enzyme activity (nmol/h/mg protein) extracted from peripheral blood total leukocytes in healthy individual, unaffected and affected patients with TSD
Table 4.2:	Hexosaminidase-A (Hex-A) and total-hexosaminidase (total-Hex) enzyme activity (nmol/h/mg protein) extracted from peripheral blood total leukocytes in healthy individual, unaffected and affected patients with SD
Table 4.3:	Hexosaminidase-A% (%Hex-A) enzyme activity (nmol/h/mg protein) extracted from peripheral blood total leukocytes in healthy individual and TSD carriers
Table 4.4:	Hexosaminidase-A% (%Hex-A) enzyme activity (nmol/h/mg protein) extracted from cultured trophoblasts (CT cells) and cultured amniotic fluid cells (AF cells)
Table-4.5:	Demographic profile for TSD patients
Table 4.6:	Common mutations screening for TSD patients
Table 4.7:	List of all <i>HEXA</i> sequence variants identified in TSD patients in present study
Table 4.8:	List of all <i>HEXA</i> sequence variants identified in TSD carrier parents
Table 4.9:	MPLA analysis for deletion/duplication analysis of <i>HEXA</i> gene
Table 4.10:	Details of <i>HEXA</i> missense mutations detected in infantile TSD patients and Insilico analysis
Table 4.11:	Details of <i>HEXA</i> gene intronic variants/deletion/insertion predicted by NNSplice 0.9 and Mutation taster
Table 4.12:	Ligand binding energy details for wild type and mutant $\alpha$ -subunits
Table 4.13:	Hydrogen bond interaction for the wild type and Mutant complex
Table 4.14:	Haplotype analysis of founder mutation E462V in Gujarati patients with Tay-

## Sachs disease

- Table 4.15: Overall clinical, biochemical and molecular profile of Indian TSD patients
- Table 4.16: Genotype-phenotype correlation of classical and AB variant (GM2A) TSD patients based on type of sequence variant
- Table 4.17: Genotype-phenotype correlations of classical and AB variant (GM2A) TSD patients individually
- Table 4.18: Enzyme activity in AF/CT cells for prenatal diagnosis
- Table 4.19: List of all *HEXA* gene sequence variants identified in prenatal study.