

REFERENCES

- Abdul Azeez, K. A., Unsworth, M. J., Patro, P. K. Harinarayana, T., and Sastry, R. S. (2013). Resistivity structure of the Central Indian Tectonic Zone (CITZ) from multiple magnetotelluric (MT) profiles and tectonic implications. *Pure and Applied Geophysics*, 170, 2231–2256.
- Acharyya, S. K., and Roy, A. (2000). Tectonothermal history of the central Indian tectonic zone and reactivation of major faults/shear zones. *Journal of Geological Society of India*, 55, 239–256.
- Aitchison, J. C., Ali, J. R., and Davis, A. M. (2007). When and where did India and Asia collide? *Journal of Geophysical Research: Solid Earth*, 112, B05423.
- Akhtar, K., and Ahmad, A. M. (1991). Palaeocurrents and sediment dispersal patterns of Lower Cretaceous sandstones, northeastern Gujarat. *Indian Journal of Earth Sciences*, 18, 10–17.
- Akhtar, K., and Khan, D. A. (1997). A tidal island model for carbonate sedimentation: Karondia Limestone of Cretaceous Narmada basin, India. *Journal of the Geological Society of India*, 50, 481–489.
- Akolkar, G. N., and Limaye, M. A. (2020). Geochemistry of calc-silicate rocks around Lunavada region, NE Gujarat: Implications for their protolith, provenance and tectonic setting. *Journal of Earth System Science*, 129, 1–14.
- Alpert, S. P. (1973). *Bergaueria* Prantl (Cambrian and Ordovician), a probable actinian trace fossil. *Journal of Paleontology*, 47, 919–924.

- Alpert, S. P. (1974). *Trace fossils of the Precambrian-Cambrian succession, White Inyo Mountains, California*. (Unpublished Ph.D. thesis). University of California, Los Angeles.
- Alpert, S. P., and Moore, J. N. (1975). Lower Cambrian trace fossil evidence for predation on trilobites. *Lethaia*, 8, 223–230.
- Andreu, B., Colin, J. P., and Singh, J. (2007). Cretaceous (Albian to Coniacian) ostracodes from the subsurface of the Jaisalmer Basin, Rajasthan, India. *Micropaleontology*, 53, 345–370.
- Arthur, M. A., Dean, W. E., and Pratt, L. M. (1988). Geochemical and climatic effects of increased marine organic carbon burial at the Cenomanian/Turonian boundary. *Nature*, 335, 714–717.
- Aslam, M. (1991). Trace fossils from the Ranipat sediments (Early Cretaceous), Saurashtra Basin, Gujarat, Western India. *Current science*, 61, 403–405.
- Auden, J. B. (1949). Dykes of western India: A discussion of their relationships with the Deccan Traps: *Transactions of the National Institute of Science*, 3, 123–157.
- Badve, R. M. (1987). A reassessment of stratigraphy of Bagh Beds, Barwah area, Madhya Pradesh, with description of trace fossils. *Journal of the Geological Society of India*, 30, 106–120.
- Badve, R. M., and Ghare, M. A. (1978). Palaeoecological aspects of the Bagh Beds, India. *Recent Researches in Geology*, 4, 388–402.
- Badve, R. M., and Ghare, M. A. (1980). Ichnofauna of Bagh Beds from Deva River valley, south of Narmada. *Biovigyanam*, 6, 121–130.
- Badve, R. M., and Nayak, K. K. (1984). Some additional fossil algae from the Nimar Sandstone, Bagh Beds, Madhya Pradesh, India (pp. 185–196). *Proceedings of the 10th Indian Colloquium Micropalaeontology and Stratigraphy*.
- Bardhan, S., Gangopadhyay, T. K. and Mandal, U. (2002). How far did India drift during the Late Cretaceous? -*Placenticeraskaffrarium* Etheridge, 1904 (*Ammonoidea*) used as a measuring tape. *Sedimentary Geology*, 147, 193–217.

- Barrett, B. J., Hodgson, D. M., Jackson, C. A. L., Lloyd, C., Casagrande, J., and Collier, R. E. L. (2021). Quantitative analysis of a footwall-scarp degradation complex and syn-rift stratigraphic architecture, Exmouth Plateau, NW Shelf, offshore Australia. *Basin Research*, 33, 1135–1169.
- Basse, E. (1947). Les peuplements Malagaches de *Barroisiceras* (Révision du genre Barroisiceras de Gross). Paléontologie de Madagascar 26. *Annales de Paléontologie*, 33, 97–178.
- Beaumont, H., Clarke, S. M., Burley, S. D., Taylor, A. M., and Mohapatra, P. (2019). Sedimentology and the facies architecture of the Ghaggar-Hakra Formation, Barmer Basin, India: Implications for early Cretaceous deposition on the north-western Indian Plate margin. *The Depositional Record*, 5, 53–83.
- Bera, S. (1996). Remarks on Paleoenvironment from the trace fossil from subsurface and outcrop Tertiary–Quaternary sediments of the western part of Bengal basin, India. *Journal of Geography and Environment*, 1, 1–15.
- Berger, W. H. (1979). Preservation of foraminifera. In J. H. Lipps, W. H. Berger, M. A. Buzas, R. G. Douglas and C. A. Ross (Eds.), *Foraminiferal Ecology and Paleoecology; Society Of Economic Paleontologists and Mineralogists* (Vol. 6, pp. 105–155). Houston: Society of Economic Paleontologists and Mineralogists, Short Course.
- Besairie, H. (1972). Géologie de Madagascar. I. Les terrains sédimentaires. *Annales Geologique Madagascar*, 35, 1–463.
- Besse, J., and Courtillot, V. (1988). Paleogeographic maps of the continents bordering the Indian Ocean since the Early Jurassic. *Journal of Geophysical Research: Solid Earth*, 93, 11791–11808.
- Bhandari, A. (1999). Phanerozoic Stratigraphy of Western Rajasthan Basin: A Review. In P. Kataria (Ed.), *Proceedings of the A. B. Roy Felicitation Volume*, Geology Department, MLSU, Udaipur, pp. 126–174.
- Bhatt, N. Y., and Patel, S. J. (2017). Ichnology and Sequence Stratigraphy of Wave-Dominated Prograding Delta, Bhuj Formation (Cretaceous), Khatrod Hill Range, Mainland Kachchh, Western India. *Journal of Geosciences*, 2, 121–137.

- Bhatt, N. Y., Solanki, P. M., Prakash, N., and Das, N. (2016). Depositional environment of Himmantnagar Sandstone (Lower/Middle Cretaceous): A perspective. *The Palaeobotanist*, 56, 67–84.
- Bhattacharya, B., and Jha, S. (2014). Late Cretaceous diurnal tidal system: a study from Nimar Sandstone, Bagh Group, Narmada Valley, Central India. *Current Science*, 107, 1032–1037.
- Bhosle, B., Johnson, C., Vaghela, S., Schultz, D. J., and Dholakia, V. (2019). First Report: Trace Fossil Assemblage *Ptychoplasma* (*P. excelsum*, *P. vagans*), *Dendroidichnites* (*D. irregulare*), *Ctenopholeus* (? *C. kutcheri*) and *Bergaueria* (*B. hemispherica*) in the Cretaceous Rocks of Bagh Formation, Mainland Gujarat, India. *Ichnos*, 26, 256–265.
- Billings, E. (1862). New species of fossils from different parts of the Lower, Middle and upper Silurian rocks of Canada. – In *Palaeozoic Fossils* (Vol. 1, pp. 96–168). Montreal (Geological Survey of Canada, Dawson Brothers).
- Biswas, S. K. (1982). Rift basins in western margin of India and their hydrocarbon prospects with special reference to Kutch basin. *American Association of Petroleum Geologists Memoir*, 66, 1497–1513.
- Biswas, S. K. (1987). Regional tectonic framework, structure and evolution of the western marginal basins of India. *Tectonophysics*, 135, 307–327.
- Biswas, S. K. (1999) A review on the evolution of rift basins in India during Gondwana with special reference to western Indian basins and their hydrocarbon prospects. In A. Sahni and R. S. Loyal (Eds.), *Gondwana Assembly: New issues and perspectives*. Proceedings of the Indian National Science Academy Special Issue, 65, 261–283.
- Biswas, S. K. (2016). Mesozoic and tertiary stratigraphy of Kutch* (Kachchh) – A review. In M. G. Thakkar (Ed.), Recent studies on the geology of Kachchh (Vol. 6, pp. 1–24). Bengaluru, India: Geological Society of India, Special Publication.
- Biswas, S. K., and Deshpande, S. V. (1983). Geology and hydrocarbon prospects of Kutch, Saurashtra and Narmada Basins. *Petroleum Asia Journal*, 6, 111–126.
- Blanford, H. F. (1862). On the Cretaceous and other rocks of South Arcot and Trichinopoly districts. *Memoirs of the Geological Survey of India*, 4, 7–217.

- Blanford, W. T. (1869). On the geology of Tapti and Lower Narmada valley and some adjoining districts. *Memoirs of the Geological Survey of India*, 6, 163–221.
- Blissett, D. J., and Pickerill, R. K. (2004). Soft-sediment ichnotaxa from the Cenozoic White Limestone Group, Jamaica, West Indies. *Scripta Geologica*, 127, 341–378.
- Boggs, S. Jr. (1987). Principles of sedimentology and stratigraphy: Columbus, Ohio, Merrill Publishing Co., 784 p.
- Borkar, V. D. and Kulkarni, K. G. (1992). On the Occurrence of Planolites Nicholson from the Bbaduka Limestone of the Wadhwan Formation (Cretaceous), Kathiawar, Gujarat. *Journal of Geological Society of India*, 40, 468–473.
- Borkar, V. D., and Kulkarni, K. G. (2001). On the occurrence of Rhizocorallium Zenker from the Habur Formation (Aptian), Rajasthan. *Gondwana Geological Magazine*, 16, 15–20.
- Borkar, V. D., and Kulkarni, K. G. (2002). Trace fossils from Fatehgarh Formation (? Aptian), Rajasthan. *Gondwana Geological Magazine*, 17, 33–37.
- Borkar, V. D., and Kulkarni, K. G. (2021). Formalizing the New Designation ‘Sitapuri Bryozoan Limestone’, in lieu of Conventional Term ‘Coralline Limestone’, for the Youngest Member of the Bagh Formation (Turonian). *Journal of the Geological Society of India*, 97, 881–886.
- Borkar, V. D., and Phadke, A. V. (1974). Fossil flora from the Gardeshwar Formation – A new Formation of the Upper Gondwanas of India. *Earth Sciences Bulletin*, 57 –64.
- Bose, P. K., and Das, N. G. (1986). A transgressive storm-and fair-weather wave dominated shelf sequence: Cretaceous Nimar Formation, Chakrud, Madhya Pradesh, India. *Sedimentary Geology*, 46, 147–167.
- Bose, P. N. (1884). Geology of the Lower Narbada Valley between Nimawar and Kawant. *Memoirs of the Geological Survey of India*, 21, 1–72.
- Boucot, A. J., Xu, C., Scotese, C. R., and Morley, R. J. (2013). Phanerozoic Paleoclimate: An Atlas of Lithologic Indicators of Climate. In G. J. Nichols and B. Ricketts (Eds.), *Concepts in Sedimentology and Paleontology* (Vol. 11, pp. 1–18). Tulsa: SEPM Society for Sedimentary Geology.

- Boulila, S., Galbrun, B., Miller, K. G., Pekar, S. F., Browning, J. V., Laskar, J., and Wright, J. D. (2011). On the origin of Cenozoic and Mesozoic “third-order” eustatic sequences. *Earth-Science Reviews*, 109, 94–112.
- Bouma, A. H. (1962). *Sedimentology of some flysch deposits. A graphic approach to facies interpretation*. Elsevier, Amsterdam.
- Bradshaw, M. A. (1981). Palaeoenvironmental interpretations and systematic of Devonian trace fossils from the Taylor Group (Lower Beacon Supergroup), Antarctica. *New Zealand Journal of Geology and Geophysics*, 24, 615–652.
- Brady, L. F. (1949). *Oniscoidichnus*, new name for *Isopodichnus* Brady 1947 not Bornemann 1889. *Journal of Paleontology*, 23, 573.
- BRANSON, C. C. (1961). New records on the scyphomedusan *Conostichus*. *Oklahoma Geology Notes*, 21, 130–138.
- Bromley, R. G. (1990). Trace Fossils: Biology and Taphonomy. Unwin Hyman, London, 280p.
- Bromley, R. G., and Asgaard, U. (1975). Sediment structures produced by a spatangoid echinoid: a problem of preservation. *Bulletin of the Geological Society of Denmark*, 24, 261–281.
- Bromley, R. G. (1996) Trace fossil assemblages, diversity and facies. In R.G. Bromley (Ed.), *Trace Fossils-Biology, Taphonomy and Applications*. Chapman and Hall, London, pp. 235–240.
- Bromley, R. G. and Asgaard, U. (1979). Triassic freshwater ichnocoenoses from Carlsberg Fjord, East Greenland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 28, 39–80.
- Bromley, R. G. (1975). Trace fossils at omission surfaces. In R.W. Frey (Ed.), *The Study of Trace Fossils*. New York, Springer-Verlag, pp. 399–428.
- Bromley, R. G., and ekdale, A. A. (1984). Trace fossil preservation in flint in the European chalk. *Journal of Paleontology*, 58, 298–311.
- Bromley, R. G., Pemberton, S. G. and Rahmani, R. A. (1984). A Cretaceous woodground: the *Teredolites* ichnofacies. *Journal of Paleontology*, 58, 488–498.
- Brown, B., Müller, R. D., Struckmeyer, H. I. M., Gaina, C., Stagg, H., Symonds, P. (2003). Formation and evolution of Australian passive margins: implications for locating the

boundary between continental and oceanic crust. In R. R. Hillis and R. D. Müller (Eds.), *Evolution and Dynamics of the Australian Plate* (Vol. 372, pp. 223–243). Geological Society of Australia Special Publication.

Brown, L. F. JR. and Fisher, W. L. (1977). Seismic stratigraphic interpretation of depositional systems: examples from Brazilian rift and pull apart basins. In C. E. Payton (Ed.), *Seismic Stratigraphy—Applications to Hydrocarbon Exploration* (Vol. 26, pp. 213–248). American Association of Petroleum Geologists Memoir.

Buatois, L. A., and Mángano, M. G. (2013). Ichnodiversity and ichnodisparity: significance and caveats. *Lethaia*, 46, 281–292.

Buatois, L. A., and Mángano, M. G. (1993). Trace fossils from a Carboniferous turbiditic lake: implications for the recognition of additional nonmarine ichnofacies. *Ichnos: An International Journal of Plant and Animal*, 2, 237–258.

Buatois, L. A., and Mángano, M. G. (2011). *Ichnology: Organism-substrate interactions in space and time*. Cambridge University Press.

Buatois, L. A., and Mángano, M. G. (2012). An early Cambrian shallow-marine ichnofauna from the Puncoviscana Formation of northwest Argentina: the interplay between sophisticated feeding behaviors, matgrounds and sea-level changes. *Journal of Paleontology*, 86, 7–18.

Buatois, L. A., Mangano, M. G., Genise, J. F., and Taylor, T. N. (1998). The ichnologic record of the continental invertebrate invasion; evolutionary trends in environmental expansion, ecospace utilization, and behavioral complexity. *Palaios*, 13, 217–240.

Buatois, L. A., Wetzel, A., and Mángano, M. G. (2020). Trace-fossil suites and composite ichnofabrics from meandering fluvial systems: The Oligocene Lower Freshwater Molasse of Switzerland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 558, 109944.

Buatois, L. A., Wissak, M., Wilson, M. A., and Mángano, M. G. (2017). Categories of architectural designs in trace fossils: A measure of ichnodisparity. *Earth-Science Reviews*, 164, 102–181.

Buatois, L. A. and Mángano, M. G. (1995). The paleoenvironmental and paleoecological significance of the lactusrine *Mermia* ichnofacies: an archetypical subaqueous nonmarine trace fossil assemblage. *Ichnos*, 4, 151–161.

- Buatois, L. A., Jalfin, G., Acenolaza, F. G., (1997). Permian nonmarine invertebrate trace fossils from southern Patagonia, Argentina: ichnologic signatures of substrate consolidation and colonization sequences. *Journal of Paleontology*, 71, 324–336.
- Buckman, J. O. (1994). *Archaeonassa* Fenton and Fenton 1937 reviewed. *Ichnos: An International Journal of Plant and Animal*, 3, 185–192.
- Cabrera, M. I. L., Mángano, M. G., Buatois, L. A., Olivero, E. B., and Maples, C. G. (2019). Bivalves on the move: the interplay of extrinsic and intrinsic factors on the morphology of the trace fossil *Protovirgularia*. *Palaios*, 34, 349–363.
- Cadée, G. C., and Goldring, R. (2007). The Wadden Sea, cradle of invertebrate ichnology. In W. Miller III (Ed.), *Trace Fossils: Concepts, Problems, Prospects* (pp. 3–13). Elsevier, London, UK.
- Caldeira, K., and Rampino, M. R. (1990). Carbon dioxide emissions from Deccan volcanism and a K/T boundary greenhouse effect. *Geophysical Research Letters*, 17, 1299–1302.
- Carter, H. J. (1857). Neocomian fossils from Bagh and its neighbourhood presented by Lieut. R H Keatinge. On the contribution to the Geology of Central and Western India. *Journal of the Bombay Branch of the Royal Asiatic Society*, 5, 614–638.
- Casshyap, S. M., and Aslam, M. (1992). Deltaic and shoreline sedimentation in Saurastra Basin, Western India: an example of infilling in an Early Cretaceous failed rift. *Journal of Sedimentary Petrology*, 62, 972–991.
- Catuneanu, O. (2002). Sequence stratigraphy of clastic systems: concepts, merits, and pitfalls. *Journal of African Earth Sciences*, 35, 1–43.
- Catuneanu, O. (2006). Principles of Sequence Stratigraphy. Elsevier, Amsterdam, 386 p.
- Catuneanu, O. (2017). Sequence stratigraphy: Guidelines for a standard methodology. In M. Montenari (Ed.), *Stratigraphy and timescales* (Vol. 2, pp. 1–57). Academic Press, UK.
- Catuneanu, O. (2019). Scale in sequence stratigraphy. *Marine and Petroleum Geology*, 106, 128–159.

- Catuneanu, O., Abreu, V., Bhattacharya, J. P., Blum, M. D., Dalrymple, R. W., Eriksson, P. G., Fielding, C.R., Fisher, W.L., Galloway, W.E., and Winker, C. (2009). Towards the standardization of sequence stratigraphy. *Earth-Science Reviews*, 92, 1–33.
- Catuneanu, O., Abreu, V., Bhattacharya, J. P., Blum, M. D., Dalrymple, R. W., Eriksson, P. G., Fielding, C. R., Fisher, W. L., Galloway, W. E., Gibling, M. R., Giles, K. A., Holbrook, J. M., Jordan, R., Kendall, C. G., St, C., Macurda, B., Martinsen, O. J., Miall, A. D., Neal, J. E., Nummedal, D., Pomar, L., Posamentier, H. W., Pratt, B. R., Sarg, J. F., Shanley, K. W., Steel, R. J., Strasser, A., Tucker, M. E. and Winker, C. (2009). Toward the standardization of sequence stratigraphy. *Earth Science Reviews*, 92, 1–33.
- Catuneanu, O., Galloway, W. E., Kendall, C. G. S. C., Miall, A. D., Posamentier, H. W., Strasser, A., and Tucker, M. E. (2011). Sequence stratigraphy: methodology and nomenclature. *Newsletters on stratigraphy*, 44, 173–245.
- Chamberlain, C. K. (1977). Ordovician and Devonian trace fossils from Nevada: Nevada Bureau of Mines and Geology, Bulletin 90, 24 pp.
- Chamyal, L. S., Maurya, D. M., Bhandari, S., and Raj, R. (2002). Late Quaternary geomorphic evolution of the lower Narmada valley, Western India: implications for neotectonic activity along the Narmada–Son Fault. *Geomorphology*, 46, 177–202.
- Chancellor, G. R., Kennedy, W. J., and Hancock, J. M. (1994). Turonian ammonite faunas from central Tunisia. *Special Papers in Palaeontology*, 50, 1–118.
- Chaudhuri, A., Banerjee, S., and Chauhan, G. (2020). Compositional evolution of siliciclastic sediments recording the tectonic stability of a pericratonic rift: Mesozoic Kutch Basin, western India. *Marine and Petroleum Geology*, 111, 476–495.
- Chen, X., Wang, C., Wu, H., Kuhnt, W., Jia, J., Holbourn, A., Zhang, L. and Ma, C. (2015). Orbitally forced sea-level changes in the upper Turonian–lower Coniacian of the Tethyan Himalaya, southern Tibet. *Cretaceous Research*, 56, 691–701.
- Chiplonkar, G. W. (1982). Bagh beds- Their stratigraphy, fauna, age and affinities and environment of deposition- a review (pp. 1–22). Presidential address, 69th Session of the Indian Science Congress, Mysore, India, section of geology and geography.

- Chiplonkar, G. W. (1983). I often recollect. *Journal of the Palaeontological Society of India*, 28, 1–6.
- Chiplonkar, G. W. and Badve, R. M. (1970). Trace fossils from the Bagh Beds. *Journal of the Palaeontological Society of India*, 14, 1–10.
- Chiplonkar, G. W. and Badve, R. M. (1972). Trace fossils from the Bagh Beds. Pt.II, *Journal of the Palaeontological Society of India*, 15, 1–5.
- Chiplonkar, G. W., and Badve, R. M. (1976). Palaeontology of the Bagh Beds. Part V. *Ostreina* (I). *Proceedings of the Indian Academy of Sciences-Section B*, 83, 244–256.
- Chiplonkar, G. W., and Badve, R. M. (1978). Palaeontology of the Bagh Beds—Part V. *Ostreina* (II). *Proceedings of the Indian Academy of Sciences-Section B, Animal Sciences*, 87, 105–117.
- Chiplonkar, G. W., and Badve, R. M. (1980). Depositional conditions of the Bagh sediment as indicated by the oyster beds. *Biovigyanam*, 6, 43–49.
- Chiplonkar, G. W., and Badwe, R. M. (1969). A preliminary note on trace fossils from the Bagh beds of Nerbada Valley. *Current Science*, 38, 594–594.
- Chiplonkar, G. W., and Borkar, V. D. (1973). *Feddenichnus*, a new burrow fossil from the Wadhwan Formation, Saurashtra. *Current Science*, 42, 572–573.
- Chiplonkar, G. W., and Ghare, M. A. (1975). Occurrence of *Keckia annulata* Glocker, in the Bagh Beds of Narmada Valley, *Current Science*, 44, 583–584.
- Chiplonkar, G. W., and Ghare, M. A. (1975). Some additional trace fossils from the Bagh Beds. *Bulletin of the Indian Geological Association*, 8, 71–84.
- Chiplonkar, G. W., and Ghare, M. A. (1976). Palaeontology of the Bagh Beds-Part VII: Ammonoidea. *Bulletin of Earth Science*, 4, 1–10.
- Chiplonkar, G. W., and Ghare, M. A. (1977). Palaeontology of the Bagh beds: Part X- *Scaphitidae*. *Proceedings of the Indian Academy of Sciences-Section B*, 85, 67–76.
- Chiplonkar, G. W., Badve, R. M., and Ghare, M. A. (1977a). On the stratigraphy of Bagh Beds of the Lower Narmada valley. In B. S. Venkatachala and V. V. Sastri (Eds.), *Proceedings*

of 4th Indian colloquium micropaleontology and stratigraphy (pp. 209–216). Dehradun: Institute of Petroleum Exploration.

Chiplonkar, G. W., Ghare, M. A., and Badve, R. M. (1977b). Bagh beds – Their fauna, age and affinities: A retrospect and prospect. *Biovigyanam*, 3, 33–60.

Chiplonkar, G. W., and Tapaswi, P. M., (1972). A new trace fossil from the Upper Cretaceous of South India, *Current Science*, 41, 747pp.

Chongzhi, T., Guoping, B., Junlan, L., Chao, D., Xiaoxin, L., Houwu, L., Dapeng, W., Yuan, W. and Min, L. (2013). Mesozoic lithofacies palaeogeography and petroleum prospectivity in North Carnarvon Basin, Australia. *Journal of Palaeogeography*, 2, 81–92.

Choubey, V. D. (1971). Narmada–Son Lineament, India. *Nature Physical Science*, 232, 38–40.

Cobban, W. A., and Hook, S. C. (1980). The Upper Cretaceous (Turonian) ammonite family Coilopoceratidae Hyatt in the Western Interior of the United States. *United States Geological Survey, Professional Paper*, 1192, 1–28.

Coffin, M. F., Pringle, M. S., Duncan, R. A., Gladczenko, T. P., Storey, M., Müller, R. D., and Gahagan, L. A. (2002). Kerguelen hotspot magma output since 130 Ma. *Journal of Petrology*, 43, 1121–1137.

Collier, J. S., Sansom, V., Ishizuka, O., Taylor, R. N., Minshull, T. A., and Whitmarsh, R. B. (2008). Age of Seychelles–India break-up. *Earth and Planetary Science Letters*, 272, 264–277.

Collinson, J. D. (1969). The sedimentology of the Grindslow shales and the Kinderscout grit; a deltaic complex in the Namurian of northern England. *Journal of Sedimentary Research*, 39, 194–221.

Collinson, J. D. (1996). Alluvial sediments. In H. G. Reading (Ed.), *Sedimentary environments: Processes, Facies, and Stratigraphy* (pp. 37–82). Blackwell Inc., Oxford.

Collinson, J. D., Thompson, D. B. (1989). Sedimentary Structures, second edition. George Allen and Unwin, London, 207 pp.

Compton, P. M. (2009). The geology of the Barmer Basin, Rajasthan, India, and the origins of its major oil reservoir, the Fatehgarh Formation. *Petroleum Geoscience*, 15, 117–130.

- Courtillot, V., Besse, J., Vandamme, D., Montigny, R., Jaeger, J. J., and Cappetta, H. (1986). Deccan flood basalts at the Cretaceous/Tertiary boundary?. *Earth and Planetary Science Letters*, 80, 361–374.
- Crawford, A. J., and von Rad, U. (1994). The petrology, geochemistry and implications of basalts dredged from the Rowley Terrace-Scott Plateau and Exmouth Plateau margins, northwestern Australia. *Journal of Australian Geology and Geophysics*, 15, 43–54.
- Crimes, T. P., and Herdman, J. (1970). Trilobite tracks and other trace fossils from the Upper Cambrian of North Wales. *Geological Journal*, 7, 47–68.
- Crimes, T. P. (1977). Trace fossils of an Eocene deep-sea sand fan, northern Spain. In T. P. Crimes and J. C. Harper (Eds.), *Trace Fossils 2* (Vol. 9, pp. 71–90). Geological Journal, Special Issue.
- Crimes, T. P., Legg, I., Marcos, A., and Arboleya, M. L. (1977). Late precambrian-low Lower Cambrian trace fossils from Spain. In T. P. Crimes and J. C. Harper (Eds.), *Trace Fossils 2* (Vol. 9, pp. 91–138). Geological Journal, Special Issue.
- Cross, T. A. (1991). High-resolution stratigraphic correlation from the perspectives of base-level cycles and sediment accommodation. In J. Dolson (Ed.), *Unconformity Related Hydrocarbon Exploration and Accumulation in Clastic and Carbonate Settings* (pp. 28–41). Rocky Mountain Association of Geologists, Short Course Notes.
- Cuerda, A. J., Arondo, O. G., Morel, E., and Spalletti, L. A. (1990). Procesos de continentalización en el Devónico de la Precordillera. *Revista del Museo de La Plata*, 10, 185–195.
- Curran, H. A. and Frey, R. W. (1977). Pleistocene trace fossils from North Carolina (USA), and their Holocene analogues. *Geological Journal Special Issues*, 9, 139–62.
- D'Alessandro, A. R., Bromley, G. (1987). Meniscate trace fossils and the *Muensteria-Taenidium* problem. *Palaeontology*, 30, 743–763.
- Dalrymple, R. W. (2010). Tidal depositional systems. In N. P. James and R. W. Dalrymple (Eds.), *Facies Models 4* (pp. 201–232). Geological Association of Canada, St. John's.

- Dam, G. (1990). Paleoenvironmental significance of trace fossils from the shallow marine Lower Jurassic Neill Klinter Formation, East Greenland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 79, 221–248.
- Dangerfield, F. (1818). Some accounts of caves near Bagh called Panch Pandoo. *Transactions of the Literary Society of Bombay*, 2, 194–204.
- Darngawn, J. L., Patel, S. J., Joseph, J. K., and Shitole, A. D. (2018). Palaeoecological significance of trace fossils of Chorar Island, Eastern Kachchh Basin, Western India. *Journal of the Palaeontological Society of India*, 63, 169–180.
- Das Gupta, S. K. (1977). Stratigraphy of western Rajasthan shelf. *Proceedings of 4th Indian Colloquium on Micropalaeontology and Stratigraphy* (pp. 219–233). Dehradun, India.
- Das, S., Singh, P. K., and Sikarni, C. (2009). A preliminary study of thermal metamorphism in the Champaner Group of rocks in Panchmahals and Vadodara districts of Gujarat. *Indian Journal of Geosciences*, 63, 373–382.
- Dassarma, D. C., and Sinha, N. K. (1966). On the occurrence of shark teeth and other marine fossils in the Nimar Sandstone horizon of the Bagh Beds. *Indian Minerals*, 20, 1–105.
- Dassarma, D. C., and Sinha, N. K. (1975). Marine Cretaceous formations of Narmada Valley (Bagh Beds), Madhya Pradesh and Gujarat. *Paleontologica Indica New Series*, 42, 1–106.
- De Raaf, J. F. M., Reading, H. G., and Walker, R. G. (1965). Cyclic sedimentation in the Lower Westphalian of north Devon. *England: Sedimentology*, 4, 1–52.
- Desai, B. G., and Saklani, R. D. (2014). Ichnofabric analysis of the Tithonian shallow marine sediments (Bhadasar Formation) Jaisalmer Basin, India. *Journal of Earth System Science*, 123, 1413–1431.
- Desai, B. G., and Saklani, R. D. (2015). Palaeocommunity dynamics and behavioral analysis of *Conichnus*: Bhuj Formation (Lower Cretaceous), Kachchh-India. *Ichnos*, 22, 43–55.
- Desai, B. G., Patel, S. J., Shukla, R., and Surve, D. (2008). Analysis of ichnoguilds and their significance in interpreting ichnological events: A study from Jhuran Formation (Upper

Jurassic), western Kachchh, India. *Journal of the Geological Society of India*, 72, 458–466.

Díez-Canseco, D., Buatois, L. A., Mángano, M. G., Díaz-Molina, M., and Benito, M. I. (2016). Ichnofauna from coastal meandering channel systems (Upper Cretaceous Tremp Formation, South-Central Pyrenees, Spain): delineating the fluvial-tidal transition. *Journal of Paleontology*, 90, 250–268.

Ding, Y., Liu, J., and Chen, F. (2020). Ichnology, palaeoenvironment, and ecosystem dynamics of the early Cambrian (stage 4, series 2) Guanshan Biota, South China. *Geological Journal*, 55, 77–94.

Dolson, J., Burley, S. D., Sunder, V. R., Kothari, V., Naidu, B., Whiteley, N. P., Farrimond, P., Taylor, A., Direen, N. and Ananthakrishnan, B. (2015). The discovery of the Barmer Basin, Rajasthan, India, and its petroleum geology. *American Association of Petroleum Geologists Memoir*, 99, 433–465.

Donovan, S. K., (2018). A new ichnogenus for *Teredoliteslongissimus* Kelly and Bromley. *Swiss Journal of Palaeontology*, 137, 95–98.

Donovan, S. K., and Ewin, T. A. (2018). Substrate is a poor ichnotaxobase: a new demonstration. *Swiss Journal of Palaeontology*, 137, 103–107.

Dott, R. H. (1964). Wacke, graywacke and matrix; What approach to immature sandstone classification? *Journal of Sedimentary Research*, 34, 625–632.

Dunham, R. J. (1962). Classification of carbonate rocks according to depositional texture. In W. E. Ham (Ed.), *Classification of carbonate rocks* (pp. 108–121). Tulsa, US: American Association of Petroleum Geologists Memoir.

Dwivedi, G. N., Mohabey, D. M., and Bandyopadhyay, S. (1982). On the discovery of vertebrate fossils from infratrappean Lameta beds, Kheda District, Gujarat. *Current Trends in Geology*, 7, 79–87.

Dymant, J. (1991). *Structure et évolution de la lithosphère océanique dans l'océan Indien: apport des anomalies magnétiques*, (Unpublished Ph.D. thesis), 374pp. Université Louis Pasteur de Strasbourg, France.

- Eagles, G., and König, M. (2008). A model of plate kinematics in Gondwana breakup. *Geophysical Journal International*, 173, 703–717.
- Ehrenberg, K. (1944). Ergänzende Bemerkungen zu den seinerzeit aus dem Miozan von Burgschleinitz beschriebenen Gangkernen und Bauten dekapoder Krebse. *Paläontologische Zeitschrift*, 23, 345–359.
- Ekdale, A. A., (1992). Muckraking and mudslinging: the joys of deposit-feeding. Short Courses in Paleontology, *The Paleontological Society*, 5, 145–171.
- Ekdale, A. A., Bromley, R. G., (2001). A day and a night in the life of a cleft-foot clam: *Protovirgularia-Lockeia-Lophoctenium*. *Lethaia*, 34, 119–124.
- Ekdale, A.A., Bromley, R.G., Pemberton, G.S. (1984). Ichnology: the use of trace fos sils in sedimentology and stratigraphy. SEPM Short Course, 15, 1–317.
- El-Sabbagh, A., El-Hedeny, M., and Al Farraj, S. (2017). Thalassinoides in the middle Miocene succession at Siwa Oasis, northwestern Egypt. *Proceedings of the Geologists' Association*, 128, 222–233.
- Emery, D., and Myers, K. J. (1996). *Sequence Stratigraphy*. Oxford, U. K., Blackwell, p. 297.
- Emmons, E. (1844). The Taconic System: Based on Observations in New York, Massachusetts, Maine, Vermont, and Rhode Island. Caroll and Cook, Albany, 68 pp.
- Fambrini, G. L., Neumann, V. H. M., Menezes-Filho, J. A. B., Silva-Filho, D., Wellington, F., and De Oliveira, É. V. (2017). Facies architecture of the fluvial Missao Velha formation (late Jurassic–early cretaceous), Araripe basin, northeast Brazil: paleogeographic and tectonic implications. *Acta Geologica Polonica*, 67, 515–545.
- Fenton, C. L., and Fenton, M. A. (1937). Archaeonassa, Cambrian snail trails and burrows. *American Midland Naturalist*, 18, 454–456.
- Fernández, M.S., Khosla, A. (2015). Parataxonomic review of the Upper Cretaceous dinosaur eggshells belonging to the oofamily *Megaloolithidae* from India and Argentina. *Historical Biology*, 27, 158–180.

- Fillion, D. and Pickerill, R. K. (1990). Ichnology of the Upper Cambrian? to Lower Ordovician Bell Island and Wabana groups of eastern Newfoundland, Canada. *Palaeontographica Canadiana*, 7, 119p.
- Fitch, A. (1850). A historical, topographical and agricultural survey of the Country of Washington. Part 2-5. *Transactions of the New York Agricultural Society*, 9, 753–944.
- Flögel, S., Wallmann, K., and Kuhnt, W. (2011). Cool episodes in the Cretaceous-Exploring the effects of physical forcings on Antarctic snow accumulation. *Earth and Planetary Science Letters*, 307, 279–288.
- Flügel, E. (2010). Microfacies of carbonate rocks: analysis, interpretation and application (2nd edition). New York, London, Springer, 984p.
- Foote, R. B. (1898). Geology of Baroda State (Baroda State: Govt Press).
- Frazier, D.E. (1974). Depositional Episodes: Their Relationship to the Quaternary Stratigraphic Framework in the Northwestern Portion of the Gulf Basin. Geological Circular. Austin, Texas: Bureau of Economic Geology.
- Freeman, K. H., and Hayes, J. M. (1992). Fractionation of carbon isotopes by phytoplankton and estimates of ancient CO₂ levels. *Global Biogeochemical Cycles*, 6, 185–198.
- Frey, F. A., McNaughton, N. J., Nelson, D. R., deLaeter, J. R., and Duncan, R. A. (1996). Petrogenesis of the Bunbury Basalt, Western Australia: interaction between the Kerguelen plume and Gondwana lithosphere?. *Earth and Planetary Science Letters*, 144, 163–183.
- Frey, R. W. (Ed.). (2012). *The study of trace fossils: a synthesis of principles, problems, and procedures in ichnology*. Springer-Verlag, Berlin, 562pp.
- Frey, R. W. and Howard, L. D. (1981). *Conichnus* and *Schaubcylindrichnus*: redefined trace fossils from the Upper Cretaceous of the Western Interior. *Journal of Paleontology* 55, 800–804.
- Frey, R. W., and Chowns, T. M. (1972). Trace fossils from the Ringgold Road cut (Ordovician and Silurian), Georgia. *Sedimentary environments in the Paleozoic rocks of northwest Georgia*, 11, 25–55.

- Frey, R. W., Curran, H. A. and Pemberton, G. S. (1984). Tracemaking activities of crabs and their environmental significance: the ichnogenus *Psilonichnus*. *Journal of Paleontology*, 58, 511–528.
- Frey, R. W. and Pemberton, S. G. (1987). The *Psilonichnus* ichnocoenose, and its relationship to adjacent marine and nonmarine ichnocoenoses along the Georgia coast. *Bulletin of Canadian Petroleum Geology*, 35, 333–357.
- Frey, R. W. and Seilacher, A. (1980). Uniformity in marine invertebrate ichnology. *Lethaia*, 13, 183–207.
- Fürsich, F. T (1974). Ichnogenus *Rhizocorallium* *Paläontologische Zeitschrift*, 48, 16–28.
- Fürsich, F. T., and Oschmann, W. (1993). Shell beds as tools in basin analysis: the Jurassic of Kachchh, western India. *Journal of the Geological Society*, 150, 169–185.
- Gaigalas, A., and Uchman, A. (2004). Trace fossils from Upper Pleistocene varved clays S of Kaunas, Lithuania. *Geologija*, 45, 16–26.
- Gaina, C., Müller, R. D., Brown, B., Ishihara, T., and Ivanov, S. (2007). Breakup and early seafloor spreading between India and Antarctica. *Geophysical Journal International*, 170, 151–169.
- Gaina, C., van Hinsbergen, D. J. J., and Spakman, W. (2015). Tectonic interactions between India and Arabia since the Jurassic reconstructed from marine geophysics, ophiolite geology, and seismic tomography. *Tectonics*, 34, 875–906.
- Gaina, C., Torsvik, T. H., vanHinsbergen, D. J. J., Medvedev, S., Werner, S.C., and Labails, C. (2013). The African Plate: A history of oceanic crust accretion and subduction since the Jurassic. *Tectonophysics*, 604, 4–25.
- Gale, A. S., J. Hardenbol, B. Hathway, W. J. Kennedy, J. R. Young, and Phansalkar. V. (2002). Global correlation of Cenomanian (upper cretaceous) sequences: Evidence for Milankovitch control on sea level. *Geology*, 30, 291–294.
- Galloway, W. E. (1989). Genetic stratigraphic sequences in basin analysis, I. Architecture and genesis of flooding-surface bounded depositional units. *American Association of Petroleum Geologists Bulletin*, 73, 125–142.

- Gangopadhyay, T. K., and Bardhan, S. (1998). Apertural modifications and jaw structures of placenticeratid ammonites from the Upper Cretaceous Bagh Group, central India. *Neues Jahrbuch für Geologie und Paläontologie-Monatshefte*, 1998, 193–202.
- Gangopadhyay, T. K., and Maiti, M. (2012). Geological implication of a turreted gastropod and astartid pelecypod bearing horizon in the nodular limestone of Sukar Nala Section near Zirabad of Bagh, Dhar District, MP, India. *Journal of Science and Technology Maharaja Sayajirao University*, 31, 45–50.
- Ganguly, T., and Bardhan, S. (1993). Dimorphism in *Placenticeras mintoi* from the Upper cretaceous Bagh Beds, central India. *Cretaceous Research*, 14, 747–756.
- Genise, J. F., Mángano, M. G., Buatois, L. A., Laza, J.H. and Verde, M. (2000). Insect trace fossil associations in paleosols: the *Coprinisphaera* ichnofacies. *Palaios*, 15, 49–64.
- Geoscience Australia (2013). [Web Page] National Petroleum Wells Database. Available at: http://dbforms.ga.gov.au/www/npm.well_search.
- Gevers, T. W. (1973). A new name for the ichnogenus *Arthropodichnus* Gevers, 1971. *Journal of Paleontology*, 47, 1002pp.
- Gevers, T. W., Frakes, L. A., Edwards, L. N., and Marzolf, J. E. (1971). Trace fossils in the Lower Beacon sediments (Devonian), Darwin Mountains, southern Victoria Land, Antarctica. *Journal of Paleontology*, 45, 81–94.
- Ghare, M. A., and Badve, R. M. (1980). Significance of trace fossils in sedimentation of the Bagh Beds. *Proceedings of 3rd Indian Geological Congress* (pp. 375–387). Pune: University of Poona.
- Ghibaudo, G., Grandesso, P., Massari, F., and Uchman, A. (1996). Use of trace fossils in delineating sequence stratigraphic surfaces (Tertiary Venetian Basin, northeastern Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 120, 261–279.
- Giannetti, A., Monaco, P., Caracuel Martín, J. E., Soria Mingorance, J. M., and Yébenes Simón, A. (2007). Functional morphology and ethology of decapod crustaceans gathered by *Thalassinoides* branched burrows in Mesozoic shallow water environments. *3rd Symposium on Mesozoic and Cenozoic Decapod Crustaceans*, 35, 48–52.

- Gibbons, A. D., Barckhausen, U., Van Den Bogaard, P., Hoernle, K., Werner, R., Whittaker, J. M., and Müller, R. D. (2012). Constraining the Jurassic extent of Greater India: tectonic evolution of the West Australian margin. *Geochemistry Geophysics Geosystems*, 13, 1–25.
- Gibbons, A. D., Whittaker, J. M., and Müller, R. D. (2013). The breakup of East Gondwana: Assimilating constraints from Cretaceous ocean basins around India into a best-fit tectonic model. *Journal of Geophysical Research: Solid Earth*, 118, 808–822.
- Gingras, M. K., MacEachern, J. A., Pickerill, R. K., (2004). Modern perspectives on the Teredolites ichnofacies: Observations from Willapa Bay, Washington. *Palaios*, 19, 79–88.
- Gingras, M. K., Pemberton, S. G., and Saunders, T. (2001). Bathymetry, sediment texture, and substrate cohesiveness; their impact on modern Glossifungites trace assemblages at Willapa Bay, Washington. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 169, 1–21.
- Glaessner, M. F. (1969). Trace fossils from the Precambrian and basal Cambrian. *Lethaia*, 2, 369–393.
- Glocker, F. E. (1841). Über die kalkführende Sandsteinformation auf beiden Seiten der mittleren March, in der Gegend zwischen Kwassitz und Kremsier. *Academia Caesarea Leopoldino–Carolina Germanica Naturae Curiosorum*, 19(Supplement 2), 309–334.
- Godbole, S. M., Rana, R. S., and Natu, S. R. (1996). Lava stratigraphy of Deccan basalts of western Maharashtra. *Gondwana Geological Magazine Special Volume*, 2, e134.
- Goldring, R., Pollard, J. E., and Radley, J. D. (2005). Trace fossils and pseudofossils from the Wealden strata (non-marine Lower Cretaceous) of southern England. *Cretaceous Research*, 26, 665–685.
- Gopalan, K., Trivedi, J. R., Merh, S. S., Patel, P. P., and Patel, S. G. (1979). Rb-Sr age of Godhra and related granites, Gujarat, India. *Proceedings of the Indian Academy of Sciences-Section A, Part 2, Earth and Planetary Sciences*, 88, 7–17.

- Ghosh, D. B. (1976). The nature of Narmada-Son Lineament, In Seminar volume on tectonics and Metallurgy of South and East Asia, *Geological Survey of India Miscellaneous Publication*, 34, 119–132.
- Gressly, A. (1838). *Observations géologiques sur le Jura soleurois*. Nouveaux mémoires de la Société Helvétique des Sciences Naturelles, Neuchâtel, 2, 349pp.
- Guha, A. K. (1976). Lithostratigraphic classification of Bagh Group (Beds), Madhya Pradesh. *Proceedings of the 4th Indian Colloquium on Micropaleontology and Stratigraphy* (pp. 209–216). Dehradun, India.
- Gupta, S. N., Arora, Y. K., Mathur, R. K., Iqbaluddin, B. P., Sahai, T. N., Sharma, S. B., and Murthy, M. V. N. (1980). Lithostratigraphic map of Aravalli region, southern Rajasthan and northeastern Gujarat. Geological Survey of India, Kolkata
- Gupta, S. N., Mathur, R. K., and Arora, Y. K. (1992). Lithostratigraphy of Proterozoic rocks of Rajasthan and Gujarat—a review. *Records of the Geological Survey of India*, 115, 63–85.
- Gupta, V. J. (1975). *Indian Mesozoic Stratigraphy* (p. 267). Delhi: Hindustan Publishing Corporation.
- Guthörl L, P. (1934). Die Arthropoden aus dem Carbon und Perm des Saar-Nahe-Pfalz-Gebietes. *Abhandlungen der Preußischen Geologischen Landesanstalt*, 164, 1–219.
- Gutiérrez-Marco, J. C. (1984). Una interesante señal de actividad biológica en el Ordovícico de los Montes de Toledo COL-PA, 39, 17–25.
- Gwalani, L. G., Rock, N. M. S., Chang, W. J., Fernandez, S., Allegre, C. J., and Prinzhofer, A. (1993). Alkaline rocks and carbonatites of Amba Dongar and adjacent areas, Deccan Igneous Province, Gujarat, India: 1. Geology, petrography and petrochemistry. *Mineralogy and Petrology*, 47, 219–253.
- Hakes, W. G. (1977). Trace fossils in Late Pennsylvanian cyclothsems, Kansas. In T. P. Crimes and J. C. Harger (Eds.), *Trace Fossils 2*. Geological Journal Special Issue, (vol. 9, pp. 209–226). Liverpool: Seel House Press.
- Haldeman, S. S. (1840). Supplement to number one of “A monography of the Limniades, and other freshwater bivalve shells of the apparently new animals in different classes, and

names and characters of the subgenera in *Paludina* and *Anculosa*: J. Dobson, Philadelphia. 3pp.

- Hall, J. (1847). Palaeontology of New York, Vol. 1. C. Van Benthuyzen, Albany, 362p.
- Han, Y., and Pickerill, R. (1994). Phycodes templus isp. nov. from the Lower Devonian of northwestern New Brunswick, eastern Canada. *Atlantic Geology*, 30, 37–46.
- Han, Y., and Pickerill, R. K. (1995). Taxonomic review of the ichnogenus *Helminthopsis* Heer 1877 with a statistical analysis of selected ichnospecies. *Ichnos: An International Journal of Plant and Animal*, 4, 83–118.
- Häntzschel, W. (1962). Trace fossils and problematica, In: R. C. Moore (Ed.), *Treatise on Invertebrate Paleontology*, Part W, Miscellanea. Geological Society of America and University. Kansas Press, Lawrence, pp. 177–245.
- Häntzschel, W. (1975) Trace fossils and problematica, In I. C. Teichert (Ed.), *Treatise on invertebrate paleontology* (2nd ed.), Part W, Miscellanea, Suppl. I. Geological Society of America and University. Kansas Press, Lawrence, 269p.
- Haq, B. U. (2014). Cretaceous eustasy revisited. *Global and Planetary change*, 113, 44–58.
- Haq, B. U., and Huber, B. T. (2017). Anatomy of a eustatic event during the Turonian (Late Cretaceous) hot greenhouse climate. *Science China Earth Sciences*, 60, 20–29.
- Haq, B. U., Hardenbol, J. A. N., and Vail, P. R. (1987). Chronology of fluctuating sea levels since the Triassic. *Science*, 235, 1156–1167.
- Harrington, H. J., and Moore, R. C. (1955). Fossil jellyfishes from Kansas Pennsylvanian rocks and elsewhere. *Bulletin of the Kansas University Geological Survey*, 114, 153–162.
- Hart, M. B., and Ball, K. C. (1986). Late Cretaceous anoxic events, sea-level changes and the evolution of the planktonic foraminifera. *Geological Society, London, Special Publications*, 21, 67–78.
- Hart, M. B., and Leary, P. N. (1989). The stratigraphic and palaeogeographic setting of the late Cenomanian ‘anoxic’ event. *Journal of the Geological Society*, 146, 305–310.

- Hasegawa, T. (1997). Cenomanian-Turonian carbon isotope events recorded in terrestrial organic matter from northern Japan. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 130, 251–273.
- Hay, W. W. (2011). Can humans force a return to a ‘Cretaceous’ climate?. *Sedimentary Geology*, 235, 5–26.
- Hay, W. W., DeConto, R., Wold, C. N., Wilson, K. M., Voigt, S., Schulz, M., Wold, A. R., Dullo, W. C., Ronov, A.B., Balukhovsky, A.N. and Söding, E. (1999). Alternative global Cretaceous paleogeography. Pages 1-47 In E. Barrera and C. Johnson (Eds.) *Evolution of Cretaceous ocean-climate systems*, Special paper 332. The Geological Society of America, Boulder, Colorado.
- Hays, J. D., and Pitman, W. C. (1973). Lithospheric plate motion, sea level changes and climatic and ecological consequences. *Nature*, 246, 18–22.
- Heer, O. (1877). Flora Fossilis Helvetiae. Die vorweltliche Flora der Schweiz. J. Wurster and Co. (Zurich), 182pp.
- Heinberg, C. (1974). A dynamic model for a meniscus filled tunnel (Ancorichnus n. ichnogen.) from the Jurassic Pecten Sandstone of Milne Land, East Greenland. *Grønlands Geologiske Undersøgelse, Raport*, 62, 1–20.
- Hewaidy, A. A., Awad, M. H., Farouk, S., and El-Balkeimy, A. (2012). Cenomanian biostratigraphy and sequence stratigraphy of southern Galala Plateau, North Eastern Desert, Egypt. *Egyptian Journal of Paleontology*, 12, 143–172.
- Hillgaertner, H. (1998). Discontinuity surfaces on a shallow-marine carbonate platform (Berriasian, Valanginian, France and Switzerland). *Journal of Sedimentary Research*, 68, 1093–1108.
- Hocking, R.M. (1990) Carnarvon Basin. In J. S. Meyers and R. M. Hocking (Eds.), *Geology and mineral resources of Western Australia: Western Australian geological survey, Memoir* (Vol. 3, pp. 457–460). Perth: State Printing Division.
- Hocking, R. M., Moors, H. T. and Van De Graaff, J. E. (1987). Geology of the Carnarvon Basin. *Bulletin of the Geological Survey of Western Australia*, 133, 1–289.

- Hofmann, H. J. (1990). Computer simulation of trace fossils with random patterns, and the use of goniograms. *Ichnos: An International Journal of Plant and Animal*, 1, 15–22.
- Hofmann, H. J., and Patel, I. M. (1989). Trace fossils from the type ‘Etcheminian Series’ (Lower Cambrian Ratcliffe Brook Formation), Saint John area, New Brunswick, Canada. *Geological Magazine*, 126, 139–157.
- Hofmann, H. J., Cecile, M. P., and Lane, L. S. (1994). New occurrences of Oldhamia and other trace fossils in the Cambrian of the Yukon and Ellesmere Island, Arctic Canada. *Canadian Journal of Earth Sciences*, 31, 767–782.
- Hofmann, R., Mángano, M. G., Elicki, O., and Shinaq, R. (2012). Paleoecologic and biostratigraphic significance of trace fossils from shallow-to marginal-marine environments from the Middle Cambrian (Stage 5) of Jordan. *Journal of Paleontology*, 86, 931–955.
- Holz, M., Vilas-Boas, D. B., Troccoli, F. B., Santana, V. C., and Vidigal-Souza, P. A. (2017). Conceptual models for sequence stratigraphy of continental rift successions. *Stratigraphy and Timescales*, 2, 119–186.
- Howard, J. D., and Frey, R. W. (1984). Characteristic trace fossils in nearshore to offshore sequences, Upper Cretaceous of east-central Utah. *Canadian Journal of Earth Sciences*, 21, 200–219.
- Howell, B. F. (1957). Stipsellus annulatus, a Skolithos-like Cambrian fossil from Arizona. *Bulletin of the Wagner Free Institute of Sciences of Philadelphia*, 32, 7–20.
- Hu, X., Jansa, L., Chen, L., Griffin, W. L., O'Reilly, S. Y., and Wang, J. (2010). Provenance of Lower Cretaceous Wulong volcaniclastics in the Tibetan Tethyan Himalaya: Implications for the final breakup of eastern Gondwana. *Sedimentary Geology*, 223, 193–205.
- Hunt, D. and Tucker, M. E. (1992). Stranded parasequences and the forced regressive wedge systems tract: deposition during base level fall. *Sedimentary Geology*, 81, 1–9.
- Hunter, S. J., Valdes, P. J., Haywood, A. M., and Markwick, P. J. (2008). Modelling Maastrichtian climate: investigating the role of geography, atmospheric CO₂ and vegetation. *Climate of the past discussions*, 4, 981–1019.

- Hyatt, A. (1903). *Pseudoceratites* of the Cretaceous. In T. W. Stanton (Ed.), *Monographs of the United States Geological Survey* (Vol. 44, pp. 1–351). Washington, US: Government printing office.
- Impey, E. (1856). Description of the caves of Bagh in Rath. *Journal of the Bombay Branch of the Royal Asiatic Society*, 5, 543–573.
- Ismail, A. A., Hussein-Kamel, Y. F., Boukhary, M., and Ghandour, A. A. (2009). Late Cenomanian-Early Turonian foraminifera from Eastern Desert, Egypt. *Micropaleontology*, 55, 396–412.
- ISSC (International Subcommission on Stratigraphic Classification) of IUGS International Commission on Stratigraphy (1976). In H. D. Hedberg (Ed.), *International stratigraphic guide* [1st edition]. New York: John Wiley and Sons, 200p.
- ISSC (International Subcommission on Stratigraphic Classification) of IUGS International Commission on Stratigraphy (1994). In A. Salvador (Ed.), *International stratigraphic guide* 2nd edition (214pp). Trondheim: International Union of Geological Sciences, and Boulder, Colorado, Geological Society of America.
- Jafar, S. A. (1982). Nannoplankton evidence of Turonian transgression along Narmada valley, India and Turonian-Coniacian boundary problem. *Journal of the Palaeontological Society of India*, 27, 17–30.
- Jain, S. C., Nair, K. K. K., and Yedekar, D. B. (1995). Geology of the Son-Narmada-Tapti lineament zone in central India. *Geological Survey of India, Special Publication*, 10, 1–154.
- Jain, S. L., and Bandyopadhyay, S. (1997). New titanosaurid (Dinosauria: sauropoda) from the Late Cretaceous of central India. *Journal of Vertebrate Paleontology*, 17, 114–136.
- Jain, S. P. (1971). *Contribution to the study of marine Cretaceous ostracodes of the Peninsular India* (Unpublished PhD thesis). Panjab University, Chandigarh.
- Jaityl, A. K., and Ajane, R. (2013). Comments on *Placenticeras mintoi* (Vredenburg, 1906) from the Bagh Beds (Late Cretaceous), Central India with special reference to Turonian Nodular Limestone horizon. *Journal of the Geological Society of India*, 81, 565–574.

- Jambusaria, B. B. (1970). *Geology of the area around Shrivardhanpur, Dist. Panchmahals, Gujarat with special reference to the stratigraphy, structure and metamorphism*. (Unpublished Ph.D. thesis). The M. S. University of Baroda, Vadodara.
- Jambusaria, B. B., and Merh, S. S. (1967). Deformed greywacke conglomerates of Jaban near Shrivardhanpur, Panchmahals district, Gujarat. *Indian Miner*, 8, 6–10.
- James, U. P. (1879). Description of new species of fossils and remarks on some others, from the Lower and Upper Silurian rocks of Ohio. *The Paleontologist*, 3, 17–24.
- Jana, B. N., King, S. C., and Hilton, J. (2013). Revision of the Cretaceous fossil plant-assemblage from Gardeshwar (Gujarat, India): A conifer dominated floral association from an Upper Gondwana sequence on the West Coast of India. *Journal of Asian Earth Sciences*, 73, 128–138.
- Jenkyns, H. C. (1980). Cretaceous anoxic events: from continents to oceans. *Journal of the Geological Society*, 137, 171–188.
- Jenkyns, H. C. (2010). Geochemistry of oceanic anoxic events. *Geochemistry Geophysics Geosystems*, 11, Q03004.
- Jenkyns, H. C., Gale, A. S., and Corfield R. M. (1994). Carbon- and oxygen-isotope stratigraphy of the English Chalk and Italian Scaglia and its palaeoclimatic significance. *Geological Magazine*, 131, 1–34.
- Jensen, S. (2003). The Proterozoic and earliest Cambrian trace fossil record; patterns, problems and perspectives. *Integrative and Comparative Biology*, 43, 219–228.
- Jervey, M. T. (1988). Quantitative geological modeling of siliciclastic rock sequences and their seismic expression. In C. K. Wilgus, B. S. Hastings, C. G. S. C. Kendall, H. W. Posamentier, C. A. Ross, and J. C. Van Wagoner (Eds.), *Sea level changes:an integrated approach* (Vol. 42, pp. 47–69). Society of Economic Paleontologists and Mineralogists Special Publication.
- Jha, S., Bhattacharya, B., and Nandwani, S. (2016). Significance of seismites in the Late Cretaceous transgressive Nimar Sandstone succession, Son-Narmada rift valley, Central India. *Geological Journal*, 52, 768–783.

- Joseph, J. K., and Patel, S. J. (2018). Facies characterisation of a shallow-water deltaic succession: the Upper Jurassic Wagad Sandstone Formation of Kachchh, western India. *Geologos*, 24, 137–150.
- Joseph, J. K., Patel, S. J., and Bhatt, N. Y. (2012). Trace fossil assemblages in mixed siliciclastic-carbonate sediments of the Kaladongar Formation (Middle Jurassic), Patcham Island, Kachchh, Western India. *Journal of the Geological Society of India*, 80, 189–214.
- Joseph, J. K., Patel, S. J., Darngawn, J. L., and Shitole, A. D. (2020). Ichnological analysis of Jurassic shallow to marginal marine deposits: example from Wagad Highland, Western India. *Ichnos*, 27, 35–63.
- Kaila, K. L. (1986). Tectonic framework of Narmada-Son Lineament – A continental rift system in Central India from Deep Seismic soundings. Reflection Seismology: A global perspective. *Geodynamics Series*, 13, 133–150.
- Karaszewski, W. (1971). Some fossil traces from the lower Liassic of the Holy Cross Mts., Central Poland. *Bulletin de l'Academie Polonaise des Sciences, Serie des Sciences de la Terre*, 19, 101–105.
- Keatinge, R. H. (1856). On Neocomian fossils from Bagh and its neighbourhood. *Journal of the Bombay Branch of the Royal Asiatic Society of India*, 5, 621–625.
- Keighley, D. G. and pickerill, R. K. (1994). The ichnogenus *Beaconites* and its distinction from *Ancorichnus* and *Taenidium*. *Palaeontology*, 37, 305–337.
- Keller, G., Nagori, M. L., Chaudhary, M., Reddy, A. N., Jaiprakash, B. C., Spangenberg, J. E., Mateo, P. and Adatte, T. (2021). Cenomanian-Turonian sea-level transgression and OAE2 deposition in the Western Narmada Basin, India. *Gondwana Research*, 94, 73–86.
- Kelly, S. R. A., and Bromley, R. G. (1984). Ichnological nomenclature of clavate borings. *Palaeontology*, 27, 793–807.
- Kennedy, W. J. (1967). Burrows and surface traces from the Lower Chalk of Southern England. *Bulletin of the British Museum (Natural History), Geology*, 5, 127–167.

- Kennedy, W. J., Phansalkar, V. G., and Walaszczyk, I. (2003). *Prionocyclus germari* (Reuss, 1845), a Late Turonian marker fossil from the Bagh Beds of central India. *Cretaceous Research*, 24, 433–438.
- Kent, R. (1991). Lithospheric uplift in eastern Gondwana: Evidence for a long-lived mantle plume system?. *Geology*, 19, 19–23.
- Kent, R. W., Pringle, M. S., Müller, R. D., Saunders, A. D., and Ghose, N. C. (2002). $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of the Rajmahal basalts, India, and their relationship to the Kerguelen Plateau. *Journal of Petrology*, 43, 1141–1153.
- Khosla, A., and Sahni, A. (2003). Biodiversity during the Deccan volcanic eruptive episode. *Journal of Asian Earth Sciences*, 21, 895–908.
- Khosla, A., Kapur, V. V., Sereno, P. C., and Wilson, J. A. (2003). First dinosaur remains from the Cenomanian-Turonian Nimar Sandstone (Bagh Beds), District Dhar, Madhya Pradesh, India. *Journal of the Palaeontological Society of India*, 48, 115–127.
- Kim, J. Y. (1994). A unique occurrence of *Lockeia* from the Yeongheung Formation (Middle Ordovician), Yeongweol, Korea. *Ichnos: An International Journal of Plant and Animal*, 3, 219–225.
- Knaust, D. (2002). Ichnogenus pholeus Fiege, 1944, revisited. *Journal of Paleontology*, 76, 882–891.
- Knaust, D. (2012). Trace fossil systematics. In D. Knaust and R. G. Bromley (Eds.), *Trace fossils as indicators of sedimentary environments. Developments in sedimentology* (Vol. 64, pp. 79–101). Amsterdam: Elsevier.
- Knaust, D. (2013). The ichnogenus Rhizocorallium: classification, trace makers, palaeoenvironments and evolution. *Earth-Science Reviews*, 126, 1–47.
- Knaust, D. (2015). Siphonichnidae (new ichnofamily) attributed to the burrowing activity of bivalves: Ichnotaxonomy, behaviour and palaeoenvironmental implications. *Earth-Science Reviews*, 150, 497–519.
- Knaust, D. and Bromley, R. G. (Eds.). (2012). *Trace fossils as indicators of sedimentary environments. Developments in Sedimentology*, 64. Elsevier, Amsterdam.

- Knaust, D., and Hauschke, N. (2004). Trace fossils versus pseudofossils in Lower Triassic playa deposits, Germany. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 215, 87–97.
- Kominz, M. A., Browning, J. V., Miller, K. G., Sugarman, P. J., Mizintseva, S., and Scotese, C. R. (2008). Late Cretaceous to Miocene sea-level estimates from the New Jersey and Delaware coastal plain coreholes: An error analysis. *Basin Research*, 20, 211–226.
- Krishna, J. (1987). An overview of the Mesozoic stratigraphy of Kachchh and Jaisalmer basins. *Journal of the Palaeontological Society of India*, 32, 136–149.
- Krishna, J. (2017). The Indian Mesozoic chronicle: sequence stratigraphic approach. Singapore: Springer Geology.
- Krishna, K. S., Gopala Rao, D., and Sar, D. (2006). Nature of the crust in the Laxmi Basin (14–20°N), western continental margin of India. *Tectonics*, 25, C1006.
- Kříž, J., and Mikuláš, R., (2006). Bivalve wood borings of the ichnogenus *Teredolites* Leymerie from the Bohemian Cretaceous Basin (Upper Cretaceous, Czech Republic). *Ichnos*, 13, 159–174.
- Krumbein, W. C., and Sloss, L. L. (1963). *Stratigraphy and sedimentation*, 2nd ed.: Freeman, San Francisco, 660p.
- Ksiazkiewicz, M. (1977). Trace fossils in the flysch of the Polish Carpathians. *Palaeontologia Polonica* 36, 1–208.
- Kuhnt, W., Thurow, J., Wiedmann, J., and Herbin, J. P. (1986). Oceanic anoxic conditions around the Cenomanian/Turonian boundary and the response of the biota. In E. T. Degens, P. A. Meyers and S. C. Brassell (Eds.), *Biogeochemistry of black shales* (Vol. 60, pp. 205–246). Mitteilungen der Geologisch-Palaontologisches Institut der Universität Hamburg, Germany.
- Kulkarni, K. G., Borkar, V. D., and Petare, T. J. (2008). Ichnofossils from the Fort Member (Middle Jurassic), Jaisalmer Formation, Rajasthan. *Journal of the Geological Society of India*, 71, 731–738.

- Kumar, P. (1994). Palynology of Carbonaceous Clays of Nimar Formation, Jhabua District, Madhya Pradesh, India. *Journal of the Geological Society of India*, 44, 671–674.
- Kumar, S., Gautam, J. P., Pandey, B., Pathak, D. B., and Jaitly, A. K. (2016). Some Unknown Arcid Bivalves from the Turonian (Late Cretaceous) of Central India. *International Journal of Science and Research*, 5, 1405–1410.
- Kumar, S., Pathak, D. B., Pandey, B., Jaitly, A. K., and Gautam, J. P. (2018). The age of the Nodular Limestone Formation (Late Cretaceous), Narmada Basin, central India. *Journal of Earth System Science*, 127, 1–7.
- Kumar, S., Singh, M. P., and Mohabey, D. M. (1999). *Lameta and Bagh Beds, Central India* (pp. 1–49). Palaeontological Society of India, Department of Geology, University of Lucknow: Field Guide.
- Kundal, P., and Sanganwar, B. N. (1998a). Stratigraphical, palaeogeographical and palaeoenvironmental significance of fossil calcareous algae from Nimar Sandstone Formation, Bagh Group (Cenomanian–Turonian) of Pipaldehla, Jhabua Dt, MP. *Current Science*, 75, 702–708.
- Kundal, P., and Sanganwar, B. N. (1998b). Stratigraphy and Palichnology of Nimar Sandstone, Bagh Beds of Jabot Area, Jhabua District, Madhya Pradesh. *Journal of the Geological Society of India*, 51, 619–634.
- Kundal, P., and Sanganwar, B. N. (2000). Ichnofossils from Nimar Sandstone Formation, Bagh Group of Manawar area, Dhar district, Madhya Pradesh. *Memoirs of the Geological Society of India*, 46, 229–244.
- Kuypers, M. M., Pancost, R. D., and Damste, J. S. S. (1999). A large and abrupt fall in atmospheric CO₂ concentration during Cretaceous times. *Nature*, 399, 342–345.
- Larson, R. L. (1991). Latest pulse of Earth: Evidence for a mid-Cretaceous superplume. *Geology*, 19, 547–550.
- Lawver, L. A., Gahagan, L. M., and Coffin, M. F. (1992). The development of paleoseaways around Antarctica. *Antarctic Research Series*, 56, 7–30.

- Lesquereux, L. (1876). Species of fossil marine plants from the Carboniferous Measures. *Indiana Geological Survey, Annual Report*, 7, 134–145.
- Lesquereux, L. (1880). Description of the coal flora of the Carboniferous formation in Pennsylvania and throughout the United States. *Geological Survey of Pennsylvania, Report of Progress*, 2, 694pp. (85 Pls. publ. in 1879).
- Lewis, C. J., & Sircombe, K. N. (2013). Use of U-Pb geochronology to delineate provenance of North West Shelf sediments, Australia. In M. Keep and S. J. Moss (Eds.), *The Sedimentary Basins of Western Australia IV*. Proceedings of the Petroleum Exploration Society of Australia Symposium. Perth, WA.
- Leymerie, M. A. (1842). Suite du memoire sur le Terrain Crétacé du Département de l'Aube. *Mémoires de la Société Géologique de la France*, 5, 1–34.
- Lindholm, R. C. (1987). Analysis of sedimentary structures. In *A Practical Approach to Sedimentology* (pp. 43–58). Dordrecht: Springer.
- Longley, I. M., Buessenschuett, C., Clydsdale, L., Cubitt, C. J., Davis, R. C., Johnson, M. K., Marshall, N. M., Murray, A. P., Somerville, R., Spry, T. B. and Thompson, N. B. (2002). The North West Shelf of Australia – a Woodside perspective. In M. Keep and S. J. Moss (Eds.), *The Sedimentary Basins of Western Australia 3: Proceedings of the Petroleum Exploration Society of Australia Symposium*, Perth: PESA.
- Loyal, R. S., Mohabey, D. M., Khosla, A., Sahani, A. (1999). Status and paleobiology of the Late Cretaceous Indian theropods with description of new theropod eggshells oogenus *Ellipsoolithus khedaensis* from the Lameta Formation, district Kheda, western India. In, B. P. Perez-Moreno, A. Holtz, J. L. Sanz, J. Mortella (Eds.), *Aspects of Theropod Biology* (Vol. 15, 379–387). GAIA.
- MacEachern, J. A., and Burton, J. A. (2000). Firmground *Zoophycos* in the Lower Cretaceous Viking Formation, Alberta: a distal expression of the *Glossifungites* Ichnofacies. *Palaios*, 15, 387–398.
- MacEachern, J. A., Bann, K. L., Pemberton, S. G. and Gingras, M. K. (2007). The Ichnofacies paradigm: High-resolution paleoenvironmental interpretation of the rock record. In J. A.

MacEachern, K. L. Bann, M. K. Gingras and S. G. Pemberton (Eds.), *Applied Ichnology* (Vol. 52, pp. 27–64). *Society for Sedimentary Geology Short Course Notes*.

MacEachern, J. A., Pemberton, S. G., Gingras, M. K., and Bann, K. L. (2007). The Ichnofacies Paradigm-Chapter 4: A Fifty-Year Retrospective. Martinsson, A. (1970). Toponomy of trace fossils. In T. P. Crimes and J. C. Harper (Ed.), *Trace Fossils* (Vol. 3, pp. 323–330). *Geological Journal Special Issue*.

MacEachern, J. A. and Pemberton, S. G. (1992). Ichnological aspects of Cretaceous shoreface successions and shoreface variability in the Western Interior Seaway of North America. In S. G. Pemberton (Ed.), *Applications of Ichnology to Petroleum Exploration*, *Society for Sedimentary Geology (SEPM)*, pp.57–84.

MacEachern, J. A., Zaitlin, B. A. and Pemberton, S. G. (1999). A sharpbased sandstone of the Viking Formation, Joffre Field, Alberta, Canada: criteria for recognition of transgressively incised shoreface complexes. *Journal of Sedimentary Research*, 69, 876–892.

Madhavaraju, J., Ramasamy, S., Mohan, S. P., Hussain, S. M., Gladwin Gnana Asir, N., and Stephen Pitchaimani, V. (2004). Petrography and surface textures on quartz grains of Nimar Sandstone, Bagh Beds, Madhya Pradesh-Implications for provenance and depositional environment. *Journal of the Geological Society of India*, 64, 747–762.

Mamtani, M. A. (1998). *Deformational mechanisms of the Lunavada Pre-Cambrian rocks, Panchmahal District, Gujarat*. (Unpublished PhD thesis). The Maharaja Sayajirao University of Baroda, Vadodara.

Mángano, M. G. and Buatois, L. A. (2004). Reconstructing early Phanerozoic intertidal ecosystems: Ichnology of the Cambrian Campanario Formation in northwest Argentina. *Fossils and Strata*, 51, 1–22.

Mángano, M. G., and Rindsberg, A. (2003). Carboniferous Psammichnites: systematic re-evaluation, taphonomy and autecology. *Ichnos*, 9, 1–22.

Mángano, M. G., Buatois, L. A., and West, R. R. and Maples, C. G. (2002). Ichnology of a Pennsylvanian Equatorial Tidal Flat-The Stull Shale Member at Waverly, Eastern Kansas. *Kansas Geological Survey, Bulletin*, 245, 1–133.

- Mankar, R. S., and Srivastava, A. K. (2019). Revised paleogeography of the dinosaur bearing Maastrichtian Lameta Formation, Central and Western India: in the perspective of newly identified Salbardi-Belker inland basin. *Journal of Sedimentary Environments*, 4, 53–71.
- Marck, W. [von der] (1894). *Dreginozoum nereitiforme*, ein vergessenes Fossil der oberen Kreide Westfalens von Dolberg bei Hamm. *Verhandlungen des Naturhistorischen Vereins der preußischen Rheinlande und Westphalens*, 51, 1–9.
- Martinsson, A. (1970). Toponomy of trace fossils. In T. P. Crimes, and J. C. Harper (Eds.), *Trace fossils* (Vol. 3, pp. 323–330). Geological Journal, Special Issue.
- Mayoral, E., Santos, A., Vintaned, J. G., Wissak, M., Neumann, C., Uchman, A., and Nel, A. (2020). Bivalve bioerosion in Cretaceous-Neogene amber around the globe, with implications for the ichnogenera *Teredolites* and *Apectoichnus*. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 538, 109410.
- McCall, P. L., and Tevesz, M. J. (1983). Soft-bottom succession and the fossil record. In M. J. S. Tevesz and P. L. McCall (Eds.), *Biotic Interactions in Recent and Fossil Benthic Communities* (pp. 157–194). New York: Plenum Press.
- McCann, T. (1993). A Nereites ichnofacies from the Ordovician-Silurian Welsh Basin. *Ichnos: An International Journal of Plant and Animal*, 3, 39–56.
- McCann, T., and Pickerill, R. K. (1988). Flysch trace fossils from the Cretaceous Kodiak Formation of Alaska. *Journal of Paleontology*, 62, 330–348.
- McIlroy, D. (1998). Additional Ichnotaxa from the Flysch-Type Deposits of the Arenig Ribband Group, Co. Wexford, Ireland. *Irish Journal of Earth Sciences*, 17, 103–113.
- McIlroy, D. (2004). Some Ichnological concepts, methodologies, applications and frontiers. In D. McIlroy (Ed.), *The Application of Ichnology to Palaeoenvironmental and Stratigraphic Analysis* (Vol. 228, pp. 3–27). Geological Society Special Publication.
- McLean, D. M. (1985). Deccan Traps mantle degassing in the terminal Cretaceous marine extinctions. *Cretaceous Research*, 6, 235–259.
- McLoughlin, S., Haig, D. W., Backhouse, J., Holmes, M. A., Ellis, G., Long, J. A., and McNamara, K. J. (1995). Oldest Cretaceous sequence, Giralia Anticline, Carnarvon

- Basin, Western Australia: late Hauerivian-Barremian. *Journal of Australian Geology and Geophysics*, 15, 445e468.
- Medlicott, H. B. (1875). The Shahpur coalfield with notice of coal exploration in Narbada region. *Records of the Geological Survey of India*, 8, 65–86.
- Melchor, R. N., Bromley, R. G., and Bedatou, E. (2009). Spongeliomorpha in nonmarine settings: an ichnotaxonomic approach. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 100, 429–436.
- Merh, S. S. (1995). Geology of Gujarat (p. 222). Bangalore, India: Geological Society of India.
- Merh, S. S. and Chamyal, L. S. (1997). The Quaternary geology of Gujarat alluvial plains. *Indian National Science Academy*, 63, 1–98.
- Miall A. D. (2000). Principles of sedimentary basin analysis, 3rd edition. Berlin: Springer.
- Miall, A. D. (1977). A review of the braided-river depositional environment. *Earth-Science Reviews*, 13, 1–62.
- Miall, A. D. (1984). Principles of sedimentary basin analysis, Springer-Verlag, 490p.
- Middleton, G. V. (1973). Johannes Walther's Law of the Correlation of Facies. *Geological Society of America Bulletin*, 84, 979–988.
- Miall, A. D. (1996). The geology of fluvial deposits: sedimentary facies, basin analysis, and petroleum geology. Berlin: Springer.
- Miall, A. D. (1997). The Geology of Stratigraphic Sequences. Springer-Verlag, Berlin, 433pp.
- Miall, A. D. (2015). Updating uniformitarianism: stratigraphy as just a set of “frozen accidents”. In D. G. Smith, R. J. Bailey, P. Burgess and A. Fraser, (Eds.), *Strata and Time* (Vol. 404, pp. 11–36). Geological Society, London, Special Publications.
- Middleton, G. V. (1973). Johannes Walther's Law of the Correlation of Facies. *Geological Society of America Bulletin*, 84, 979–988.
- Middleton, G. V. (1978). Facies, In R. W. Fairbridge, and J. Bourgeois (Eds.), *Encyclopedia of Sedimentology* Stroudsbury, Pennsylvania, Dowden, Hutchinson and Ross, pp. 323–325.

- Mikuláš, R., Rindsberg, A. K., Santos, A., and Pavela, M. (2017). A Carboniferous chiton (Mollusca, Polyplacophora) at the end of its trail: a unique find from the Czech Republic. *Bollettino della Società Paleontologica Italiana*, 56, 172.
- Miller, W., III (Ed.) (2007). Trace Fossils: Concepts, Problems, Prospects. Elsevier, Amsterdam, 632 pp.
- Miller, K. G., Kominz, M. A., Browning, J. V., Wright, J. D., Mountain, G. S., Katz, M. E., Sugarman, P. J., Cramer, B. S., Christie-Blick, N. and Pekar, S. F. (2005). The Phanerozoic record of global sea-level change. *science*, 310, 1293–1298.
- Miller. S. A. (1889). North American Geology' and Paleontology for the Use of Amateurs, Students and Scientists. Western Methodist Book Concern, Cincinnati, 664 pp
- Minter, N. J., and Braddy, S. J. (2009). Ichnology of an early permian intertidal flat: the Robledo Mountains Formation of southern New Mexico, USA. *Special Papers in Palaeontology*, 82, 5–107.
- Mitchum, R. M., JR. (1977). Seismic stratigraphy and global changes of sea level, part 11: glossary of terms used in seismic stratigraphy. In C. E. Payton (Ed.), *Seismic Stratigraphy—Applications to Hydrocarbon Exploration* (v. 26, pp. 205–212). American Association of Petroleum Geologists Memoir.
- Mohabey D. M., (1991). *Palaeontological studies of the Lameta Formation with special reference to dinosaurian eggs from Kheda and Panchmahal districts of Gujarat, India* (Unpublished Ph. D. thesis). Nagpur University, Nagpur.
- Mohabey, D. M. (1983). Note on the occurrence of dinosaurian fossil eggs from Infratrappean Limestone in Kheda district, Gujarat. *Current Science*, 52, 1194–1194.
- Moriya, K., Wilson, P. A., Friedrich, O., Erbacher, J., and Kawahata, H. (2007). Testing for ice sheets during the mid-Cretaceous greenhouse using glassy foraminiferal calcite from the mid-Cenomanian tropics on Demerara Rise. *Geology*, 35, 615–618.
- Mount, J. (1985). Mixed siliciclastic and carbonate sediments: a proposed first-order textural and compositional classification. *Sedimentology*, 32, 435–442.

- Mude, S. N. (2012a). Paleoenvironmental significance of ichnofossils from the Kand Formation of the Cambay Basin, Gujarat, India. *Proceedings of the International Academy of Ecology and Environmental Sciences*, 2, 12–20.
- Mude, S. N. (2012b). Paleo-Environmental Significance of Ichnofossils from the Babaguru Formation of the Cambay Basin, Gujarat, India. *Universal Journal of Environmental Research and Technology*, 2, 52–57.
- Mude, S. N., Jagtap, S. A., Kundal, P., Sarkar, P. K., and Kundal, M. P. (2012). Paleoenvironmental significance of ichnofossils from the Mesozoic Jaisalmer Basin, Rajasthan, north western India. *Proceedings of the International Academy of Ecology and Environmental Sciences*, 2, 150.
- Müller, R. D., Mihut, D., and Baldwin, S. (1998). A new kinematic model for the formation and evolution of the West and Northwest Australian Margin. In P. G. Purcell and R. R. Purcell (Eds.), *The Sedimentary Basins of Western Australia 2* (pp. 56–72): Proceedings of the Petroleum Exploration Society of Australia Symposium, Perth: PESA.
- Müller, R. D., Sdrolias, M., Gaina, C., Steinberger, B., and Heine, C. (2008). Long-term sea-level fluctuations driven by ocean basin dynamics. *Science*, 319, 1357–1362.
- Murphy, M. A., and Salvador, A. (1999). International subcommission on stratigraphic classification of IUGS international commission on stratigraphy-international stratigraphic guide—an abridged version. *Episodes*, 22, 255–271.
- Murty, A. S. N., Rao, P. K., Dixit, M. M., Rao, G. K., Reddy, M. S., Prasad, B. R., and Sarkar, D. (2011). Basement configuration of the Jhagadia–Rajpipla profile in the western part of Deccan syncline, India from travel-time inversion of seismic refraction and wide-angle reflection data. *Journal of Asian Earth Sciences*, 40, 40–51.
- Murty, A. S. N., Sarkar, D., Sen, M. K., Sridher, V., and Prasad, A. S. S. S. R. S. (2014). Mapping the sub-trappean Mesozoic sediments in the western part of Narmada–Tapti region of Deccan Volcanic Province, India. *Journal of Asian Earth Sciences*, 93, 15–24.
- Murty, K. N., Rao, R. P., Dhokarikar, B. G., and Varma, C. P. (1963). On the occurrence of plant fossils in the Nimar sandstone near Umrali, District Jhabua, Madhya Pradesh. *Current Science*, 32, 21–22.

- Murty, K. N., Rao, R. P., Dhokarikar, B. G., and Varma, C. P. (1963). On the occurrence of plant fossils in the Nimar sandstone near Umrali, District Jhabua, Madhya Pradesh. *Current Science*, 32, 21–22.
- Myannil, R. M. (1966). A small vertically excavated cavity in Baltic Ordo-vician limestone. Akademiya Nauk SSSR, Paleontologicheskiy Institut, pp.200-207.
- Myrow, P. M. (1995). *Thalassinoides* and the enigma of Early Paleozoic open-framework burrow systems. *Palaios*, 10, 58–74.
- Nagendra, R., Patel, S. J., Deepankar, R., and Reddy, A. N. (2010). Bathymetric significance of the ichnofossil assemblages of the Kulakkalnattam sandstone, Ariyalur area, Cauvery Basin. *Journal of the Geological Society of India*, 76, 525–532.
- Nagm, E. (2009). *Integrated Stratigraphy, Palaeontology and Facies Analysis of the Cenomanian-Turonian (Upper Cretaceous) Galala and Maghra El Hadida Formations of the Western Wadi Araha, Eastern Desert, Egypt* (Unpublished Ph.D. thesis). Würzburg University, Germany.
- Nagm, E., and Wilmsen, M. (2012). Late Cenomanian–Turonian (Cretaceous) ammonites from Wadi Qena, central Eastern Desert, Egypt: taxonomy, biostratigraphy and palaeobiogeographic implications. *Acta Geologica Polonica*, 62, 63–89.
- Nagm, E., El-Qot, G., and Wilmsen, M. (2014). Stable-isotope stratigraphy of the Cenomanian–Turonian (upper Cretaceous) boundary event (CTBE) in Wadi Qena, eastern Desert, Egypt. *Journal of African Earth Sciences*, 100, 524–531.
- Nair, K. K., Jain, S. C., and Yedekar, D. B. (1985). Geology, structure and tectonics of the Son-Narmada-Tapti lineament zone. *Records of the Geological Survey of India*, 117, 138–147.
- Nayak, K. K. (1987). Foraminifera from the Nimar Sandstone of Bagh beds, Jhabua district, Madhya Pradesh. *Biovigyanam*, 13, 30–39.
- Nayak, K. K. (2000). Additional trace fossils from the Nimar Sandstone, Bagh Group, Jhabua District, Madhya Pradesh. *Bulletin Oil and Natural Gas commission India*, 37, 189–197.

- Nayak, K. K. (2000a). Additional Bivalves from the Nimar Sandstone, Bagh Beds, Jhabua District, Madhya Pradesh. *Memoirs of the Geological Society of India*, 46, 139–158.
- Nayak, K. K. (2000b). Additional trace fossils from the Nimar Sandstone, Bagh Group, Jhabua District, Madhya Pradesh. *Bulletin Oil and Natural Gas commission India*, 37, 189–197.
- Nayak, K. K. (2004). Ostracods from Nimar Sandstone Formation, Bagh Group, Pipaldehla, Jhabua District, Madhya Pradesh, India. *Gondwana Geological Magazine*, 19, 85–91.
- Nemec, W., and Steel, R. J. (1984). Alluvial and coastal conglomerates: their significant features and some comments on gravelly mass-flow deposits. In E. H. Koster and R. J. Steel (Eds.). *Sedimentology of Gravels and Conglomerates* (Vol. 10, pp. 1–31). Canadian Society of Petroleum Geologists, Memoir.
- Nguyen, L. C., Hall, S. A., Bird, D. E., and Ball, P. J. (2016). Reconstruction of the East Africa and Antarctica continental margins. *Journal of Geophysical Research: Solid Earth*, 121, 4156–4179.
- Nicholls, K. H. (2019). A Geoconservation perspective on the trace fossil record associated with the end-Ordovician mass extinction and glaciation in the Welsh Basin. (Unpublished Ph.D. thesis). University of Chester, UK.
- Nichols, G. (2009). Sedimentology and Stratigraphy. Wiley-Blackwell publications, 410p.
- Nicholson and G. J. Hinde. (1875). Notes on the fossils of the Clinton, Niagara, and Guelph Formations of Ontario, with descriptions of new species. *The Canadian Journal of Science, Literature and History*, 14, 137–160.
- Nicholson, H. A. (1873). Contributions to the study of the errant annelides of the older Palaeozoic rock. – *Proceedings of the Royal society of London*, 21, 288–290; London [also: Geological Magazine 1: 309–310].
- Nielsen, J. K., Milà, J., and Mesfun, D. (2015). Trace fossils from the Eocene Lillebælt Clay Formation, Røsnæs, Denmark. In *Annales Societatis Geologorum Poloniae*, 85, 493–505).

- Obrist-Farner, J., Ball, P. J., McGilvery, T. A. M., and Rogers, R. R. (2017). A prograding margin during global sea-level maxima: an example from Mahajanga Basin, northwest Madagascar. *Basin Research*, 30, 671–687.
- Ogg, J. G., Ogg, G. M. and Gradstein, F. M. (2008). The Concise Geologic Time Scale (177pp.). Cambridge, U. K: Cambridge University Press.
- Olivero, D. (2003). Early Jurassic to Late Cretaceous evolution of Zoophycos in the French subalpine basin (southeastern France). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 192, 59–78.
- Osgood, R. G. (1970). Trace fossils of the Cincinnati Area. *Palaeontographica Americana*, 6, 281–444.
- Pal, A. K. (1970). On Reappraisal of the fossil fauna from the Bagh Beds of the Man River section, M.P. *The Quarterly Journal of the Geological, Mining, and Metallurgical Society of India*, 42, 101–105.
- Pal, A. K. (1971). *A note on the biozonation of the Bagh Group of Madhya Pradesh* (p. 333). Bangalore, India: Proceedings of the Indian Science Congress.
- Congress.Pal, T. K., Ray, S. K., Talukder, B., and Naik, M. K. (2007). On the megainvertebrate fauns (mollusca and brachiopoda) of Cenozoic and Mesozoic of Jaisalmer, Rajasthan and their stratigraphic implications. *Records of the Zoological Survey of India, Occasional paper No. 280*, pp. 40.
- Paranjape, A. R., Kulkarni, K. G., and Gurav, S. S. (2013). Significance of Lockeia and associated trace fossils from the Bada Bagh Member, Jaisalmer Formation, Rajasthan. *Journal of Earth System Science*, 122, 1359–1371.
- Paranjape, A. R., Kulkarni, K. G., and Kale, A. S. (2011). Report of trace fossils from the syn-rift Terani Plant beds, Early Cretaceous, Cauvery Basin, Tamil Nadu: Implications on basin evolution and hydrocarbon exploration. Abstract Volume of the 23rd Indian Colloquium on Micropalaeontology and Stratigraphy and International Bioevents in Earth's History, Bangalore, 4–5.

- Paranjape, A. R., Kulkarni, K. G., and Kale, A. S. (2015). Sea level changes in the upper Aptian–lower/middle (?) Turonian sequence of Cauvery Basin, India—An ichnological perspective. *Cretaceous Research*, 56, 702–715.
- Paranjape, A. R., Kulkarni, K. G., and Kale, A. S. (2016). Sedimentology and Ichnology of the syn-rift Terani Member, Cauvery Basin, India: Implications for early evolution of a passive margin. In A. Khosla, and S. G. Lucas (Eds.), *Cretaceous Period: Biotic Diversity and Biogeography* (Vol. 71, 263–273). Bulletin.
- Parihar, V. S., Nama, S. L., and Mathur, S. C. (2017). Discovery of Trace Fossils from Lower Odania Member of Lathi Formation of Jaisalmer Basin, Akal area, District-Jaisalmer, Western Rajasthan, India. *Advances in Ecological and Environmental Research*, pp. 195–210.
- Parihar, V. S., Nama, S. L., Khichi, C. P., Shekhawat, N. S., Snehlata, M., and Mathur, S. C. (2016). Near Shore-Shallow Marine (*Ophiomorpha* and *Margaritichnus*) Trace Fossils from Fatehgarh Formation of Barmer Basin, Western Rajasthan, India. *Journal of Ecosystem and Ecography*, 6, 180.
- Pascoe, E. H. (1959). *A manual of the geology of India and Burma* (pp. 485–1343). Calcutta, India: Government of India Press.
- Patel, S. J., Joseph, J. K., and Bhatt, N. Y. (2014). Ichnology of the Goradongar Formation, Goradongar Hill Range, Patcham Island, Kachchh, Western India. *Journal of the Geological Society of India*, 84, 129–154.
- Patel, S. J., and Patel, N. J. (2015). Sedimentological and palaeoecological significance of the trace fossils of the Jurassic rocks of the Jhura Dome, Mainland Kachchh, western India. *Volumina Jurassica*, 13, 101–140.
- Patel, S. J., Shitole, A. D., and Joseph, J. K. (2018). Plug shaped burrows *Conichnus-Conostichus* from the Late Cretaceous of Bagh Group, Gujarat, western India. *Journal of the Geological Society of India*, 91, 41–46.
- Patro, B. P. K., Harinarayana, T., Sastry, R. S., Rao, M., Manoj, C., Naganjaneyulu, K., and Sarma, S. V. S. (2005). Electrical imaging of Narmada–Son lineament zone, central India from magnetotellurics. *Physics of the Earth and Planetary Interiors*, 148, 215–232.

- Paumard, V., Bourget, J., Payenberg, T., Ainsworth, R. B., George, A. D., Lang, S., Posamentier, H. W. and Peyrot, D. (2018). Controls on shelf-margin architecture and sediment partitioning during a syn-rift to post-rift transition: Insights from the Barrow Group (Northern Carnarvon Basin, North West Shelf, Australia). *Earth-Science Reviews*, 177, 643–677.
- Payton, C. E. (1977). Seismic stratigraphy-applications to hydrocarbon exploration. *American Association of Petroleum Geologists Memoir*, 26, 516pp.
- Pearson, P. N., and Gooday, R. J. (2019). A large *Taenidium* burrow from the Upper Carboniferous of Corrie, Isle of Arran, and remarks on the association of *Taenidium* burrows and *Diplichnites* trails. *Scottish Journal of Geology*, 55, 135–140.
- Pemberton, R. K. and Frey, R. W. (1982). Trace fossils nomenclature and *Planolites Palaeophycus* dilemma. *Journal of Palaeontology*, 56, 416–439.
- Pemberton, S. G. and Frey, R. W. (1984). Ichnology of storm-influenced shallow marine sequence: Cardium Formation (Upper Cretaceous) at Seebe, Alberta, In D. F. Stott and D. J. Glass (Eds.), *The Mesozoic of middle North America* (Vol. 9, pp. 281–304). Canadian Society of Petroleum Geologists, Memoir.
- Pemberton, S. G., MacEachern, J. A., Dashtgard, S. E., Bann, K. L., Gingras, M. K., and Zonneveld, J. P. (2012). Shorefaces. In D. Knaust and R. G. Bromley (Eds.), *Trace fossils as indicators of sedimentary environments. Developments in sedimentology* (Vol. 64, pp. 563–603). Amsterdam: Elsevier.
- Pemberton, S. G., MacEachern, J. A., Gingras, M. K., and Zhang, J. (2000). Significance of ichnofossils to genetic stratigraphy. *Science in China Series D: Earth Sciences*, 43, 541–560.
- Pemberton, S. G., R. W. Frey, and R. G. Bromley. (1988). The ichnotaxonomy of *Conostichus* and other plug-shaped ichnofossils. *Canadian Journal of Earth Sciences*, 25, 866–892.
- Pemberton, S. G., Spila, M., Pulham, A. J., Saunders, T., Maceachern, J. A., Robbins, D. and Sinclair, I. K. (2001). Ichnology and sedimentology of shallow to marginal marine systems: Ben Nevis and Avalon Reservoirs, Jeanne d'Arc Basin. Geological Association of Canada, Short Course Notes 15, 353.

- Pemberton, S. G., MacEarchern, J. A., Saunders, T. (2004). Stratigraphic applications of substrate-specific ichnofacies: delineating discontinuities in the rock record. In D. McIlroy, (Ed.), *The Application of Ichnology to Palaeoenvironmental and Stratigraphic Analysis* (Vol. 228, pp. 29–62). Special Publication Geological Society. London: UK
- Perch-Nielsen, K. J. McKenzie, and He, Q. (1982). "Biostratigraphy and isotope stratigraphy and the 'catastrophic' extinction of calcareous nannoplankton at the Cretaceous/Tertiary boundary." *Geological implications of impacts of large asteroids and comets on the Earth. Geological Society of America, Special Paper*, 190, 353–371.
- Pervinquier, L. (1910). Sur quelques ammonites du Crétacé algérien. Mémoires de la Société Géologique de France. *Paléontologie*, 17, 1–86.
- Pfefferkorn, H. W. (1971). Note on *Conostichus broadheadi* Lesquereux (Trace Fossil: Pennsylvanian). *Journal of Paleontology*, 45, 888–892
- Phelps, R. M., Kerans, C., Da-Gama, R. O., Jeremiah, J., Hull, D., and Loucks, R. G. (2015). Response and recovery of the Comanche carbonate platform surrounding multiple Cretaceous oceanic anoxic events, northern Gulf of Mexico. *Cretaceous Research*, 54, 117–144.
- Pickerill, R. K., Romano, M., Melendez, B. (1984). Arenig trace fossils from Salamanca area, western Spain. *Geological Journal*, 19, 249–269.
- Plint, A. G., Eyles, N., Eyles, C. H., and Walker, R. G. (1992). Control of sea level change. In R. G. Walker and N. P. James (Eds.), *Facies Models: Response to Sea Level Change*. Geological Association of Canada, Alberta, pp. 15–25.
- Plint, A. G., (1988). Sharp-based shoreface sequences and “offshore bars” in the Cardium Formation of Alberta; their relationship to relative changes in sea level. In C. K. Wilgus, B. S. Hastings, C. G. S. C. Kendall, H. W. Posamentier, C. A. Ross, and J. C. Van Wagoner (Eds.), *Sea level changes:an integrated approach* (Vol. 42, pp. 357–370). Society of Economic Paleontologists and Mineralogists Special Publication.
- Plummer, C. C., and McGeary, D. (1996). *Physical Geology*. Wm. C. Brown Publishers, p. 539.

- Poddar, M. C. (1964). Mesozoics of Western India—their geology and oil possibilities. Report 22nd Session International Geological Congress (pp. 126–143). New Delhi, India, part-1, Geology of Petroleum.
- Posamentier, H. and Allen, G. (1999). Siliciclastic Sequence Stratigraphy—Concepts and Applications. SEPM Concepts in Sedimentology and Paleontology, 7, 210p.
- Posamentier, H. W., Allen, G. P., James, D. P., and Tesson, M. (1992). Forced regressions in a sequence stratigraphic framework: concepts, examples, and exploration significance. *American Association of Petroleum Geologists Bulletin*, 76, 1687–1709.
- Posamentier, H. W., Jervey, M. T. and Vail, P. R. (1988). Eustatic controls on clastic deposition II-conceptual framework. In C. K. Wilgus, B. S. Hastings, C. G. S. C. Kendall, H. W. Posamentier, C. A. Ross, and J. C. Van Wagoner (Eds.), *Sea level changes:an integrated approach* (Vol. 42, pp. 125–154). Society of Economic Paleontologists and Mineralogists Special Publication.
- Powell, C. M., Roots, S. R., and Veevers, J. J. (1988). Pre-breakup continental extension in East Gondwanaland and the early opening of the eastern Indian Ocean. *Tectonophysics*, 155, 261–283.
- Prantl, F. (1945). Dvě záhadné zkameněliny (stopy) z vrstev chrustenických—dδ3. *Rozpravy II. Třídy České Akademie*, 55, 3–8.
- Prasad, G. V., Verma, V., Sahni, A., Lourembam, R. S., and Rajkumari, P. (2017). Elasmobranch fauna from the upper most part of the Cretaceous Bagh Group, Narmada Valley, India. *Island Arc*, 26, e12200.
- Pratt, L. M., and Threlkeld, C. N. (1984). Stratigraphic significance of $^{13}\text{C}/\text{ }^{12}\text{C}$ ratios in mid-Cretaceous rocks of the western interior, U.S.A. In D. F. Stott and D. J. Glass (Eds.), *The Mesozoic of middle North America* (Vol. 9, pp. 305–312). Canadian Society of Petroleum Geologists Memoir.
- Pugh, A. T., Schröder-Adams, C. J., Carter, E. S., Herrle, J. O., Galloway, J., Haggart, J. W., Andrews, J. L. and Hatsukano, K. (2014). Cenomanian to Santonian radiolarian biostratigraphy, carbon isotope stratigraphy and paleoenvironments of the Sverdrup

- Basin, Ellef Ringnes Island, Nunavut, Canada. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 413, 101–122.
- Quenstedt, F. A. (1879). *Petrefactenkunde Deutschlands. Korallen (Sternkorallen). Erste Abtheilung, Sechster Band, Drittes Heft, Atlas*. Fues's Verlag (R. Reisland), Leipzig, pp. 337–496.
- Racey, A., Fisher, J., Bailey, H., and Roy, S. K. (2016). The value of fieldwork in making connections between onshore outcrops and offshore models: an example from India. *Geological Society, London, Special Publications*, 436, 21–53.
- Radhakrishna, T., Maluski, H., Mitchell, J. G., and Joseph, M. (1999). $^{40}\text{Ar}/^{39}\text{Ar}$ and K/Ar geochronology of the dykes from the south Indian granulite terrain. *Tectonophysics*, 304, 109–129.
- Rafferty, J. P. (Ed.). (2010). The Mesozoic Era: Age of dinosaurs: The geologic history of the earth. Britannica Educational Publishing, Newcastle upon Tyne, UK.
- Rajanikanth, A., and Chinnappa, C. H. (2016). Early Cretaceous flora of India – A review. *The Palaeobotanist*, 65, 209–245.
- Rajshekhar, C. (1991). Foraminifera from the Nodular Limestone, Bagh Beds, Madhya Pradesh, India. *Journal of the Geological Society of India*, 38, 151–168.
- Rajshekhar, C. (1995). Foraminifera from the Bagh Group, Narmada Basin, India. *Journal of the Geological Society of India*, 46, 413–428.
- Rajshekhar, C. (1997). Significance of fossils from the Rajpipla limestone, Cretaceous of Narmada basin, Gujarat. *Journal of the Geological Society of India*, 50, 339–342.
- Rajshekhar, C., and Atpalkar, S. (1995). Foraminifera from the Nodular Limestone, Bilthana, Gujarat-stratigraphic significance. *Journal of the Geological Society of India*, 45, 585–593.
- Raju, A. T. R., Chaube, A. N., and Chowdhary, L. R. (1971). Deccan Trap and the geologic framework of the Cambay Basin. *Bulletin Volcanologique*, 35, 521–538.

- Ramasamy, S., and Madhavaraju, J. (1993). Petrographic studies of the Bagh Beds along the Narmada Valley in Madhya Pradesh, India. *Gondwana Geological Magazine*, 4 and 5, 65–79.
- Ramkumar, M. (2015). *Cretaceous Sea Level Rise: Down Memory Lane and the Road Ahead*. Amsterdam: Elsevier.
- Ramkumar, M., Stüben, D., and Berner, Z. (2004). Lithostratigraphy, depositional history and sea level changes of the Cauvery basin, South India. *Ann Geol Penins Balk*, 65, 1–27.
- Raup, D. M., and Sepkoski, J. J. (1984). Periodicity of extinctions in the geologic past. *Proceedings of the National Academy of Sciences*, 81, 801–805.
- Raup, D. M., and Sepkoski, J. J. (1986). Periodic extinction of families and genera. *Science*, 231, 833–836.
- Ray, D. (1981). Stratigraphy and sedimentation of the Bagh Beds of Lower Narmada valley, Gujarat. (Unpublished Ph.D. thesis). The Maharaja Sayajirao University of Baroda, Vadodara.
- Reading, H. G. and Levell, B. K. (1996) Controls on the sedimentary rocks record. In H. G. Reading, (Ed.), *Sedimentary environments: Processes, facies and stratigraphy* (3rd edition). Blackwell Science Ltd., London, pp. 5–36.
- Reeves, C. (2009). Re-examining the evidence from plate-tectonics for the initiation of Africa's passive margins. London: Geological Society of Houston/Petroleum Exploration Society of Great, 9–10, 1–4.
- Reeves, C. V., Teasdale, J. P., and Mahanjane, E. S. (2016). Insight into the Eastern Margin of Africa from a new tectonic model of the Indian Ocean. In M. Nemcok, S. Rubar, S. T. Sinha, S. A. Hermeston, and L. Lendenyiova (Eds.), *Transform margins: Development, controls, and petroleum systems* (Vol. 431, pp. 299–322). London: Geological Society Special Publications.
- Reineck, H. E., and Wunderlich, F. (1968). Classification and origin of flaser and lenticular bedding. *Sedimentology*, 11, 99–104.

- Richards, M. A., Duncan, R. A., and Courtillot, V. E. (1989). Flood basalts and hot-spot tracks: plume heads and tails. *Science*, 246, 103–107.
- Richter, R. (1937). Marken und Spuren aus allen Zeiten. I–II. *Senckenbergiana*, 19: 150–169.
- Rieth, A. (1932). Neue Funde spongeliomorpher Fucoiden aus dem Jura Schwabens. *Geologische und Paläontologische Abhandlungen, N.F.* 19, 257–294.
- Rindsberg, A. K., and Kopaska-Merkel, D. C. (2003). *Treptichnus* and related trace fossils from the Union Chapel Mine (Langsettian, Alabama, USA). In A. J. Martin (Ed.), *Workshop on Permo-Carboniferous ichnology, program and abstracts, University of Alabama, Tuscaloosa*, pp.27–28.
- Rindsberg, A. K., and Kopaska-Merkel, D. C. (2005). *Treptichnus* and *Arenicolites* from the Steven C. Minkin paleozoic footprint site (Langsettian, Alabama, USA). *Pennsylvanian Footprints in the Black Warrior Basin of Alabama*, 1, 121–141.
- Rode, K. P., and Chiplonkar, G. W. (1935). A contribution to the stratigraphy of Bagh Beds. *Current Science*, 4, 322–323.
- Rodríguez-Tovar, F. J., Miguez-Salas, O., and Duarte, L. V. (2017). Toarcian Oceanic Anoxic Event induced unusual behaviour and palaeobiological changes in *Thalassinoides* tracemakers. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 485, 46–56.
- Rogers, R. R., Hartman, J. H., and Krause, D. W. (2000). Stratigraphic analysis of Upper Cretaceous rocks in the Mahajanga Basin, northwestern Madagascar: implications for ancient and modern faunas. *The Journal of Geology*, 108, 275–301.
- Rolfe, W. D. I. (1980). Early invertebrate terrestrial faunas: In A. L. Panchen (Ed.), *The terrestrial Environment and the Origin of Land Vertebrates* (pp. 117–157). London: Academic Press.
- Rouault, M. (1850). Note préliminaire sur une nouvelle formation découverte dans le terrain silurien inférieur de la Bretagne. *Société Géologique de France Bulletin, série*, 2, 724–744.
- Roy-chowdhary, M. K., and Sastri, V. V. (1962). On the revised classification of the Cretaceous and associated rocks of the Man River section of lower Narbada Valley. *Record Geological Survey of India*, 91, 283–4.

- Royer, J. Y., Sclater, J. G., Sandwell, D. T., Cande, S. C., Schlich, R., Munsch, M., Dyment, J., Fisher, R.L., Müller, R.D., Coffin, M.F. Patriat, P. and Bergh, H. W. (1992). Indian Ocean plate reconstructions since the Late Jurassic. In R. A. Duncan, D. K. Rea, R. B. Kidd, U. Von Rad and J. K. Weissel (Eds.). *Synthesis of results from scientific drilling in the Indian Ocean* (Vol. 70, pp. 471–475). Washington DC: American Geophysical Union.
- Ruban, D. A. (2015). Mesozoic long-term eustatic cycles and their uncertain hierarchy. *Geoscience Frontiers*, 6, 503–511.
- Ruban, D. A. (2016). A “chaos” of Phanerozoic eustatic curves. *Journal of African Earth Sciences*, 116, 225–232.
- Ruidas, D. K., Paul, S., and Gangopadhyay, T. K. (2018). A reappraisal of stratigraphy of Bagh Group of Rocks in Dhar District, Madhya Pradesh with an outline of origin of nodularity of Nodular Limestone Formation. *Journal of the Geological Society of India*, 92, 19–26.
- Sageman, B. B., Meyers, S. R., and Arthur, M. A. (2006). Orbital time scale and new C-isotope record for Cenomanian-Turonian boundary stratotype. *Geology*, 34, 125–128.
- Saha, O., Shukla, U. K., and Rani, R. (2010). Trace fossils from the late cretaceous Lameta Formation, Jabalpur area, Madhya Pradesh: Paleoenvironmental implications. *Journal of the Geological Society of India*, 76, 607–620.
- Sahni, A. (1995). Dinosaur eggshells from India. *Journal of the Palaeontological Society of India*, 40, 87–102.
- Sahni, M. R., and Jain, S. P. (1966). Note on a revised classification of the Bagh Beds, Madhya Pradesh. *Journal of the Palaeontological Society of India*, 11, 24–25.
- Salter, J. W. (1857). On annelide-burrows and surface markings from the Cambrian rocks of the Longmynd. *Quarterly Journal of the Geological Society*, 13, 199–206.
- Salvador, A. (Ed.). (1994). *International Stratigraphic Guide: A guide to stratigraphic classification, terminology, and procedure* (2nd edition). International Commission on Stratigraphy, Geological Society of America: Boulder, CO.
- Sanders, D., Lukesch, M., Rasser, M., and Skelton, P. (2007). Shell beds of diceratid rudists ahead of a low-energy gravelly beach (Tithonian, northern calcareous Alps, Austria): palaeoecology and taphonomy. *Austrian Journal of Earth Sciences*, 100, 186–199.

- Sanganwar, B. N., and Kundal, P. (1997). Ichnofossils from Nimar Sandstone Formation. Bagh Group of Barwah area, Khargaon district, MP. *Gondwana Geological Magazine*, 12, 12–33.
- Santos, H. P., Mángano, M. G., Soares, J. L., Nogueira, A. C., Bandeira, J., and Rudnitzki, I. D. (2017). Ichnologic evidence of a Cambrian age in the Southern Amazon Craton: Implications for the onset of the Western Gondwana history. *Journal of South American Earth Sciences*, 76, 482–488.
- Santos, M. N. D., Chemale Jr, F., Dussin, I. A., Martins, M. D. S., Queiroga, G., Pinto, R. T. R., Santos, A. N. Armstrong, R. (2015). Provenance and paleogeographic reconstruction of a mesoproterozoic intracratonic sag basin (Upper Espinhaço Basin, Brazil). *Sedimentary Geology*, 318, 40–57.
- Sarkar, S. K. (1973). *Sedimentology of the Nilmara-Bagh-Lameta Complex around Awaldaman and Bagh areas, Dhar district, Madhya Pradesh*. (Unpublished Ph.D. thesis). Jadavpur University, India.
- Sastray, M. V. A., and Mamgain, V. D. (1971). The marine Mesozoic Formations of India: a review. *Records of the Geological Survey of India*, 101, 162–177.
- Savi, P. and Meneghini, G. G. (1850). Osservazioni stratigrafiche e paleontologische concernenti la geologia della Toscana e dei paesi limitrofi. [Appendix to Murchison, R.I. Memoir sulla struttura geologica delle Alpi, degli Apennini et dei Carpazi]: 246-528. Stamparia granuale, Firenze
- Savrda, C. E. (1991). Ichnology in sequence stratigraphic studies: An example from the Lower Paleocene of Alabama. *Palaios*, 6, 39–53.
- Savrda, C. E., and King Jr, D. T. (1993). Log-ground and Teredolites lagerstatte in a Transgressive Sequence, Upper Cretaceous (Lower Campanian) Mooreville Chalk, central Alabama. *Ichnos: An International Journal of Plant and Animal*, 3, 69–77.
- Schatz, E. R., Mángano, M. G., Aitken, A. E., and Buatois, L. A. (2013). Response of benthos to stress factors in Holocene Arctic fjord settings: Maktak, Coronation, and North Pangnirtung Fjords, Baffin Island, Canada. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 386, 652–668.

- Schlanger, S. O., and Jenkyns, H. C. (1976). Cretaceous oceanic anoxic events: Causes and consequences. *Geologie en mijnbouw*, 55, 179–184.
- Schlirf, M. and Uchman, A. (2001). [see Schlirf, M., Uchman, A. and Kümmel, M. 2001.]
- Schlirf, M. (2000). Upper Jurassic trace fossils from the Boulonnais (northern France). *Geologica et Palaeontologica*, 34, 145–213.
- Schlirf, M., Uchman, A., and Kümmel, M. (2001). Upper Triassic (Keuper) non-marine trace fossils from the Haßberge area (Franconia, south-eastern Germany). *Paläontologische Zeitschrift*, 75, 71–96.
- Schmid, E. E. (1876). *Der Muschelkalk des östlichen Thüringen*.
- Scholle, P. A., and Arthur, M. A. (1980). Carbon isotope fluctuations in cretaceous pelagic limestones: potential stratigraphic and petroleum exploration tool. *The American Association of Petroleum Geologists Bulletin*, 64, 67–87.
- Schumm, S. A. (1993). River response to baselevel change: implications for sequence stratigraphy. *Journal of Geology*, 101, 279–294.
- Segev, A. (2000). Synchronous magmatic cycles during the fragmentation of Gondwana: radiometric ages from the Levant and other provinces. *Tectonophysics*, 325, 257–277.
- Segev, A. (2002). Flood basalts, continental breakup and the dispersal of Gondwana: evidence for periodic migration of upwelling mantle flows (plumes). *EGU Stephan Mueller Special Publication Series*, 2, 171–191.
- Seilacher, A. (1953). Studien zur Palichnologie II. Die fossilen Ruhespuren (Cubichnia). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 98, 87–124.
- Seilacher, A. (1964). Sedimentological classification and nomenclature of trace fossils. *Sedimentology*, 3, 256–253.
- Seilacher, A. (1967). Bathymetry of trace fossils. *Marine geology*, 5, 413–428.
- Seilacher, A. (1990). Paleozoic trace fossils. In R Said (Ed.), *The Geology of Egypt* (pp. 649–670) Balkema: Rotterdam.
- Seilacher, A. and Seilacher, E. (1994). Bivalvian trace fossils: A lesson from actuopaleontology. *Courier Forschungsinstitut Senckenberg*, 169, 5–15.

- Sepkoski, J. J. (1990). Mass extinction processes: Periodicity. In D. E. G. Briggs and P. R. Crowther (Eds.), *Palaeobiology* (pp. 171-179). Oxford: Blackwell.
- Shanley, K. W., McCabe, P. J., Hettinger, R. D. (1992). Significance of tidal influence in fluvial deposits for interpreting sequence stratigraphy. *Sedimentology* 39, 905–930.
- Sharma, K. K. (2007). KT magmatism and basin tectonism in western Rajasthan, India, results from extensional tectonics and not from Réunion plume activity. *Geological Society of America Special Papers*, 430, 775–784.
- Shibata, M., and Varricchio, D. J. (2020). Horseshoe crab trace fossils from the Upper Cretaceous Two Medicine Formation of Montana, USA, and a brief review of the xiphosurid ichnological record. *Journal of Paleontology*, 94, 887–905.
- Shipway, J. R., Rosenberg, G., Concepcion, G. P., Haygood, M. G., Savrda, C., and Distel, D. L. (2019). Shipworm bioerosion of lithic substrates in a freshwater setting, Abatan River, Philippines: Ichnologic, paleoenvironmental and biogeomorphological implications. *PLoS one*, 14, e0224551.
- Shitole, A. D., Patel, S. J., Darngawn, J. L., and Joseph, J. K. (2021). Amended lithostratigraphy of the Cretaceous Bagh Group, Western Lower Narmada Valley, India: A comparison with pervasive Tethyan basins. *Geological Journal*, 56, 5058–5093.
- Shitole, A. D., Patel, S. J., Joseph, J. K., and Darngawn, J. L. (2019). Ethological and environmental significance of *Bergaueria hemispherica* from the Late Cretaceous of the Bagh Group, Western India. *Comptes Rendus Palevol*, 18, 287–297.
- Singh, D., Alat, C. A., Singh, R. N., and Gupta, V. P. (1997). Source rock characteristics and hydrocarbon generating potential of Mesozoic sediments in Lodhika area, Saurashtra basin, Gujarat, India (pp. 205–220). Proceedings of Second International Petroleum Conference and Exhibition, PETROTECH., New Delhi, India.
- Singh, N. P. (2006). Mesozoic lithostratigraphy of the Jaisalmer Basin, Rajasthan. *Journal of the Palaeontological Society of India*, 51, 1–25.
- Singh, P. (1998). An atlas of the Cretaceous Foraminifera from the subsurface of Jaisalmer Basin, Rajasthan, India. Paleontology Laboratory, K D Malaviya Institute of Petroleum

Exploration, Oil and Natural Gas Corporation Limited, DehraDun, Project NO. GR1. 03. R06, 1–72.

Singh, S. (1982). *Bergaueria* Prantl from upper Cretaceous rocks of Madhya Pradesh, India. *Journal of the Palaeontological Society of India*, 27, 62–63.

Singh, S. K., and Dayal, R. M. (1979). Trace fossils and environment of deposition of Nimir Sandstone, Bagh Beds. *Journal of the Geological Society of India*, 20, 34–239.

Singh, S. K., and Srivastava, H. K. (1981). Lithostratigraphy of the Bagh Beds and its correlation with Lameta Beds. *Journal of the Paleontological Society of India*, 26, 77–85.

Sinton, C. W., and Duncan, R. A. (1997). Potential links between ocean plateau volcanism and global ocean anoxia at the Cenomanian-Turonian boundary. *Economic Geology*, 92, 836–842.

Sisodia, M. S., and Singh, U. K. (2000). Depositional environment and hydrocarbon prospects of the Barmer Basin, Rajasthan, India. *Nafia*, Zagreb (Croatia), 51, 309–326.

Skelton, P. W. (2003). Introduction to the Cretaceous. In P. W. Skelton (Ed.), *The cretaceous world*. Cambridge, UK: Cambridge University Press.

Sloss, L. L. (1962). Stratigraphic models in exploration. *American Association of Petroleum Geologists Bulletin*, 46, 1050–1057.

Smith, A. B. (2010). The Cretaceous Bagh Formation, India: a Gondwanan window onto Turonian shallow-water echinoid faunas. *Cretaceous Research*, 31, 368–386.

Smith, J. (1893). Peculiar U-shaped tubes in sandstone near Crawfurdland Castle, and in Gowkha Quarry near Kilwinning. *Transactions of the Geological Society of Glasgow*, 9, 289–292.

Spasojevic, S., and Gurnis, M. (2012). Sea level and vertical motion of continents from dynamic earth models since the Late Cretaceous. *American Association of Petroleum Geologists Bulletin*, 96, 2037–2064.

Srivastava, D. K., Rajshekhar, C., and Awere, C. M. (2011). *Mecaster mutabilis* (Lambert, 1933) from the Middle Cretaceous Nodular Limestone (Bagh Beds), Gujarat, India. *Journal of the Palaeontological Society of India*, 56, 231–236.

- Stanley, T. M., and Feldmann, R. M. (1998). Significance of nearshore trace-fossil assemblages of the Cambro-Ordovician Deadwood Formation and Aladdin Sandstone, south Dakota. *Annals of Carnegie Museum*, 67, 1–51.
- Steel, R. J., and Thompson, D. B. (1983). Structures and textures in Triassic braided stream conglomerates ('Bunter' pebble beds) in the Sherwood Sandstone Group, North Staffordshire, England. *Sedimentology*, 30, 341–367.
- Stefani, C. D. (1885). Studi paleozoologici sulle creta superiore e media dell'Apenninisettentrionale. *Atti Accademia dei Lincei, Memorie seri*, 4, 73–121.
- Stewart, J. (1821). Geological notes on the strata between Malwa and Guzerat. *Transactions of Literature Society of Bombay*, 3, 538–541.
- Storey, M., Mahoney, J. J., Saunders, A. D., Duncan, R. A., Kelley, S. P., and Coffin, M. F. (1995). Timing of hot spot-related volcanism and the breakup of Madagascar and India. *Science*, 267, 852–855.
- Tandon, S. K. (2000). Spatio-temporal patterns of environmental changes in the Late Cretaceous sequences of Central India. *Developments in Palaeontology and Stratigraphy*, 17, 225–241.
- Taylor, P. D., and Badve, R. M. (1995). A new cheilostome bryozoan from the Cretaceous of India and Europe: a cyclostome homeomorph. *Palaeontology*, 38, 627–658.
- Teichert, C. (1958). Concepts of facies. *The American Association of Petroleum Geologists Bulletin*, 42, 2718–2744.
- Tewari, A., Hart, M. B., and Watkinson, M. P. (1998). *Teredolites* from the Garudamangalam sandstone formation (Late Turonian-Coniacian), Cauvery basin, southeast India. *Ichnos: An International Journal of Plant and Animal*, 6, 75–98.
- Tewari, A., Hart, M. B. and Watkinson, M. P. (1996). A revised lithostratigraphic classification of the Cretaceous rocks of the Trichinopoly district, Cauvery basin, Southeast India. In J. Pandey, R. J. Azmi, A. Bhandari and A. Dave (Eds.), *Contributions to the XV Indian Colloquium on Micropalaeontology and Stratigraphy* (pp. 789–800).
- Tewari, H. C. (2001). A tectonic model of the Narmada region. *Current Science*, 80, 873–878.

- Tiwari, R. P., Rajkonwar, C., Malsawma, P. L. J., Ralte, V. Z., and Patel, S. J. (2011). Trace fossils from Bhurban Formation, Surma Group (lower to middle miocene) of Mizoram India and their palaeoenvironmental significance. *Journal of Earth System Science*, 120, 1127–1143.
- Torell, O. (1869). *Petrificata suecana formationis cambricae*, 6, 1–17.
- Torsvik, T. H., Tucker, R. D., Ashwal, L. D., Carter, L. M., Jamtveit, B., Vidyadharan, K. T., Venkataramana, P. (2000). Late Cretaceous India–Madagascar fit and timing of break-up related magmatism. *Terra Nova*, 12, 220–224.
- Tripathi, S. C. (1995). Palaeontological and palaeoenvironmental studies of Bagh Group, Madhya Pradesh. *Records of the Geological Survey of India*, 128, 104–105.
- Tripathi, S. C. (2006). Geology and evolution of the Cretaceous infratrappean basins of lower Narmada valley, western India. *Journal of the Geological Society of India*, 67, 459–468.
- Tripathi, S. C., and Lahiri, T. C. (2000). Marine oscillation event stratification: An example from Late Cretaceous Bagh carbonate sequence of Narmada valley, India. *Memoirs of the Geological Society of India*, 46, 15–24.
- Tripathi, S. C., and Lahiri, T. C. (2000). Marine oscillation event stratification: An example from Late Cretaceous Bagh carbonate sequence of Narmada valley, India. *Memoirs of the Geological Society of India*, 46, 15–24.
- Tsikos, H., Jenkyns, H. C., Walsworth-Bell, B., Petrizzo, M. R., Forster, A., Kolonic, S., Erba, E., Silva, I. P., Baas, M., Wagner, T. and Damsté, J. S. (2004). Carbon-isotope stratigraphy recorded by the Cenomanian–Turonian Oceanic Anoxic Event: Correlation and implications based on three key localities. *Journal of the Geological Society*, 161, 711–719.
- Tsujita, C. J., (2003). Smothered scampi: taphonomy of lobsters in the Upper Cretaceous Bearpaw Formation, southern Alberta. *Journal of Taphonomy*, 1, 187–206.
- Tunis, G., and Uchman, A. (1996). Trace fossils and facies changes in Cretaceous-Eocene flysch deposits of the Julian Prealps (Italy and Slovenia): Consequences of regional and world-wide changes. *Ichnos: An International Journal of Plant and Animal*, 4, 169–190.

- Twenhofel, W. H. (1939). Principles of Sedimentation. McGraw-Hill, New York. 610 pp.
- Uchman A., Mikulas R., Rindsberg, A. K. (2011). Mollusc trace fossils *Ptychoplasma* Fenton and Fenton, 1937 and *Oravaichnium* Plička and Uhrová, 1990: Their type material and ichnospecies. *Geobios*, 44, 387–397.
- Uchman, A. (1995). Taxonomy and palaeoecology of flysch trace fossils: The Marnosoarenacea Formation and associated facies (Miocene, Northern Apennines, Italy). *Beringeria*, 15, 1–115.
- Uchman, A. (1998). Taxonomy and ethology of flysch trace fossils: A revision of the Marian Ksiazkiewicz collection and studies of complementary material. *Annales Societatis Geologorum Poloniae*, 68, 105–218.
- Uchman, A., Kazakauskas, V., and Gaigalas, A. (2009). Trace fossils from Late Pleistocene varved lacustrine sediments in eastern Lithuania. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 272, 199–211.
- Vaidyanadhan, R. and Ramakrishnan, M. (2010). *Geology of India* (pp. 1–994). Bangalore, India: Geological Society of India.
- Vail, P. R., F. Audemard, S. A. Bowman, P. N. Eisner, and C. Perez-Cruz. (1991). The stratigraphic signatures of tectonics, eustasy and sedimentology -An overview. In G. Einsele, W. Ricken, and A. Seilacher (Ed.), *Cycles and Events in Stratigraphy* (pp. 617–659). Berlin: Springer-Verlag.
- Van Wagoner, J. C. (1995). Overview of sequence stratigraphy of foreland basin deposits: terminology, summary of papers, and glossary of sequence stratigraphy. In J. C. Van Wagoner and G. T. Bertram (Eds.), *Sequence Stratigraphy of Foreland Basin Deposits* (Vol. 64, 9–21). The American Association of Petroleum Geologists Bulletin.
- Van Wagoner, J. C., Mitchum, R. M. JR., Campion, K. M. and Rahmanian, V. D. (1990). Siliciclastic sequence stratigraphy in well logs, core, and outcrops: concepts for high resolution correlation of time and facies. *The American Association of Petroleum Geologists, Methods in Exploration, Series. 7*. Tulsa: AAPG.
- Van Wagoner, J. C., Posamentier, H. W., Mitchum, R. M. JR., Vail, P. R., Sarg, J. F., Loutit, T. S. and Hardenbol, J. (1988). An overview of sequence stratigraphy and key definitions. In

- C. K. Wilgus, B. S. Hastings, C. G. S. C. Kendall, H. W. Posamentier, C. A. Ross, and J. C. Van Wagoner (Eds.), *Sea level changes:an integrated approach* (Vol. 42, pp. 39–45). Society of Economic Paleontologists and Mineralogists Special Publication.
- Van Wagoner, J. C., Mitchum, R. M., Posamentier, H. W., Vail, P. R., (1987). An overview of sequence stratigraphy and key definitions. In A. W. Bally (Ed.), *Atlas of Seismic Stratigraphy* (Vol. 27, pp. 11–14). *The American Association of Petroleum Geologists*.
- Vassoevich, N. B. (1932). O nekotorykh priznakakh pozvolyyayushchikh otlicht'oprokinutoe polozhenie flishevых образований от normalynogo. *Akademie Nauka SSSR, Geologiduskiy Institut Trudy*, 2, 47–64.
- Veevers, J. J. (1988). Morphotectonics of Australia's northwestern margin – A review. In P. G. Purcell and R. R. Purcell (Eds.), *The North West Shelf, Australia* (pp. 19–27). Perth, WA: Proceedings of Petroleum Exploration Society of Australia Symposium.
- Veevers, J. J., and McElhinny, M. W. (1976). The separation of Australia from other continents. *Earth-Science Reviews*, 12, 139–143.
- Veevers, J. J., Powell, C. M., and Roots, S. R. (1991). Review of seafloor spreading around Australia. I. Synthesis of the patterns of spreading. *Australian Journal of Earth Sciences*, 38, 373–389.
- Verma, K. K. (1965). On fossil shark teeth from the Bagh Beds of Amba Dongar area, Gujarat State. *Current Science*, 34, 289–290.
- Verma, K. K. (1968). Bagh Beds their fauna and affinities with South Indian Cretaceous formations. *Memoirs of the Geological Society of India*, 2, 239–247.
- Verma, K. K. (1969). Critical review of the Bagh Beds of India. *Journal of the Indian Geoscience Association*, 10, 45–54.
- Verma, K. K. (1970). Occurrence of trace fossils in the Bagh Beds of Amba Dongar area, Gujarat State. *Journal of the Indian Geoscience Association*, 12, 37–40.
- Vianey-Liaud, M., Khosla, A., Garcia, G. (2003). Relationships between European and Indian dinosaur eggs and eggshells of the oofamily *Megaloolithidae*. *Journal of Vertebrate Paleontology*, 23, 575–585.

- Viglietti, P. A., McPhee, B. W., Bordy, E. M., Sciscio, L., Barrett, P. M., Benson, R. B. J., Wills, S., Chapelle, K. E. J., Dollman, K. N., Mdekazi, C. and Choiniere, J. N. (2020). Biostratigraphy of the *Massospondylus* assemblage zone (Stormberg Group, Karoo Supergroup), South Africa. *South African Journal of Geology*, 123, 249–262.
- Villegas-Martín, J., and Netto, R. G. (2019). Permian macroburrows as microhabitats for meiofauna organisms: an ancient behaviour common in extant organisms. *Lethaia*, 52, 31–43.
- Voigt, S., Erbacher, J., Mutterlose, J., Weiss, W., Westerhold, T., Wiese, F., Wilmsen, M. and Wonik, T. (2008). The Cenomanian-Turonian of the Wunstorf section-(North Germany): global stratigraphic reference section and new orbital time scale for Oceanic Anoxic Event 2. *Newsletters in Stratigraphy*, 43, 65-89.
- Vossler, S. M., and Pemberton, S. G. (1988). *Skolithos* in the Upper Cretaceous Cardium Formation: an ichnofossil example of opportunistic ecology. *Lethaia*, 21, 351–362.
- Vossler, S. M., Magwood, J. P., and Pemberton, S. G. (1989). The youngest occurrence of the ichnogenus *Didymaulichnus* from the Upper Cretaceous (Turonian) Cardium Formation. *Journal of Paleontology*, 63, 384–386.
- Vredenburg, E. W. (1907). The ammonites of the Bagh Beds. *Records of the Geological Survey of India*, 36, 109–125.
- Wadia, D. N. (1919). *Geology of India for students* (3rd ed., p. 553). London, England: MacMillan.
- Walaszczyk, I., Marcinowski, R., Praszki, T., Dembicz, K., and Bieńkowska, M. (2004). Biogeographical and stratigraphical significance of the latest Turonian and Early Coniacian inoceramid/ammonite succession of the Manasoa section on the Onilahy River, south-west Madagascar. *Cretaceous Research*, 25, 543–576.
- Walcott, C. D. (1890). Descriptive notes of new genera and species from the Lower Cambrian or Olenellus Zone of North America. *US National Museum Proceedings*, 12, 33–46.
- Walker, R. G. (1992) Facies models. In R. G. Walker and N. P. James (Eds.), *Facies Models: Response to Sea level Change* (pp.1–14). Geological Association of Canada, St. Johns.

- Walker, R. G. and Plint, A. G. (1992). Wave- and Storm-Dominated Shallow Marine systems. In R. G. Walker and N. P. James (Eds.), *Facies Models: Response to Sea Level Change* (pp. 219–238). Geological Association of Canada, St Johns.
- Wang, Y., Lin, J. P., Zhao, Y. L., and Orr, P. J. (2009). Palaeoecology of the trace fossil Gordia and its interaction with nonmineralizing taxa from the early Middle Cambrian Kaili Biota, Guizhou province, South China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 277, 141–148.
- Webby, B. D. (1970). Late Precambrian trace fossils from New South Wales. *Lethaia*, 3, 79–109.
- Weller, S. (1899). Kinderhook faunal studies 1: The fauna of the Vermicular sandstone at Northview, Webster County, Missouri. *St. Louis Academy of Science, Transactions*, 9, 9–51.
- West, W. D. (1962). The line of the Narmada and Son valleys. *Current Science*, 31, 143–144.
- Williamson, T., Henderson, R. A., Price, G. D., and Collerson, K. D. (2012). Strontium-isotope stratigraphy of the Lower Cretaceous of Australia. *Cretaceous Research*, 36, 24–36.
- Wilmsen, M., and Nagm, E. (2012). Depositional environments and faciesdevelopment of the Cenomanian–Turonian Galala and Maghra elHadida formations of the Southern Galala Plateau (Upper Cretaceous, Eastern Desert, Egypt). *Facies*, 58, 229–247.
- Wilmsen, M., and Nagm, E. (2013). Sequence stratigraphy of the lower Upper Cretaceous (Upper Cenomanian–Turonian) of the Eastern Desert, Egypt. *Newsletters on Stratigraphy*, 46, 23–46.
- Wilson, J. A., and Upchurch, P. (2003). A revision of Titanosaurus Lydekker (Dinosauria-Sauropoda), the first dinosaur genus with a ‘Gondwanan’ distribution. *Journal of Systematic Palaeontology*, 1, 125–160.
- Wilson, J. A., Sereno, P. C., Srivastava, S., Bhatt, D. K., Khosla, A., and Shani, A. (2003). A new abelisaurid (Dinosauria, Theropoda) from the Lameta formation (Cretaceous, Maastrichtian) of India. *Contributions from the Museum of Palaeontology, University of Michigan*, 31, 1–42.

- Yochelson, E. L., and Fedonkin, M. A. (1997). The type specimens (Middle Cambrian) of the trace fossil *Archaeonossa* Fenton and Fenton. *Canadian Journal of Earth Sciences*, 34, 1210–1219.
- Young, F. G. (1972). Early Cambrian and older trace fossils from the southern Cordillera of Canada. *Canadian Journal of Earth Sciences*, 9, 1–17.
- Zecchin, M. (2007). The architectural variability of small-scale cycles in shelf and ramp clastic systems: the controlling factors. *Earth-Science Reviews*, 84, 21–55.
- Zecchin, M. (2010). Towards the standardization of sequence stratigraphy: is the parasequence concept to be redefined or abandoned?. *Earth Science Reviews*, 102, 117–119.
- Zenker, J. C. (1836). Historisch-topographisches Taschenbuch von Jena und seiner Umgebung: besonders. In J. C. Zenker (Ed.), *naturwissenschaftlicher u. medicinischer Beziehung*. 338pp. Wackenhoder, Jena
- Zonneveld, J. P., Zaim, Y., Rizal, Y., Ciochon, R. L., Bettis III, E. A., and Gunnell, G. F. (2012). Ichnological constraints on the depositional environment of the Sawahlunto Formation, Kandi, northwest Ombilin Basin, west Sumatra, Indonesia. *Journal of Asian Earth Sciences*, 45, 106–113.
- Zorina, S. O., Dzyuba, O. S., Shurygin, B. N., Ruban, D. A. (2008). How global are the Jurassic - Cretaceous unconformities? *Terra Nova* 20, 341e346.