

LIST OF FIGURES

<i>Figure No.</i>	<i>Title</i>	<i>Page No.</i>
Figure 1.1	Phases of Gondwana breakup (after Reeves, 2009).	2
Figure 1.2	Reconstruction of Early Cretaceous Gondwana (eastern and central) after Segev (2002)	4
Figure 1.3	Extinction percentages of marine genera since last 270 Ma. The total number of genera is 10,383, of which 6350 are extinct (Sepkoski, 1990).	10
Figure 1.4	Conceptual contrast between lithostratigraphy and sequence stratigraphy (Emery and Myers, 1996 and Catuneanu, 2006).	13
Figure 1.5	a. Flow chart explaining the progression of data assemblage for depositional environment interpretation and construction of sequence stratigraphic model followed in the present study (Dalrymple, 2010). b. Applications, controls, integrated disciplines, and data of sequence stratigraphy (Catuneanu et al., 2009).	22
Figure 1.6	Map showing the studied sections of the Bagh Group of WLVN (location no. 1-23) and ELNV (location no. 24-36).	31
Figure 2.1	Geological and Tectonic map of the Lower Narmada Valley (Shitole et al., 2021; Abdul Azeez et al., 2013 and Jain et al., 1995) with studied localities of the Bagh Group.	36
Figure 2.2	An almost complete sequence of the Bagh Group (ELNV) exposing the basal Nimar Formation and Nodular Limestone capped by the younger Lameta Group. Note: Buddhist caves in the Nimar Sandstone at Bagh village, Dhar district of Madhya Pradesh.	38
Figure 2.3	Cretaceous paleogeography highlighting the position of India (blue) and Madagascar (yellow) during a. Berriasian-Aptian and b. Albian-Turonian (Boucot et al., 2013).	43
Figure 2.4	Tectonic Map of west-central India showing major faults, lineaments, and exposures of the Bagh Group, Narmada Basin (modified after Abdul Azeez et al., 2013; Jha et al., 2016).	44
Figure 4.1	Stratigraphic correlation of the Cretaceous Bagh Group rocks sections along the profile A-A' (Figure 2.1) in the Lower Narmada Valley; Men River Valley (8-12) and Mohanfort-Vajepur (17, 19, 23) are the	95

	composite profiles (Shitole et al., 2021).	
Figure 5.1	Two-dimensional strip log correlation of the WLNV sections.	99
Figure 6.1	Diagrammatic representation of a). <i>Archaeonassa</i> and b). <i>Bergaueria</i> assemblages.	190
Figure 6.2	Diagrammatic representation of a). <i>Conichmus- Conostichus</i> and b). <i>Helminthoidichnites-Gordia</i> assemblage.	192
Figure 6.3	Diagrammatic representation of a). <i>Lockeia-Planolites</i> and b). <i>Skolithos</i> assemblage.	193
Figure 6.4	Diagrammatic representation of a). <i>Taenidium</i> and b). <i>Thalassinoides</i> assemblages.	194
Figure 6.5	Schematic 3-dimensional diagram representing ichnoassemblages in the shallow marine depositional environment of the Western Lower Narmada Valley.	195
Figure 7.1	Transgression, regression, and forced regression defined by the interplay of base-level changes and sedimentation at the shoreline (Catuneanu, 2002, 2006; Catuneanu et al., 2009).	204
Figure 7.2	Position of sequence boundaries and the subdivision into systems tracts (Catuneanu et al., 2009).	208
Figure 7.3	Bagh Group succession of the Mohanfort-Vajepur section showing distribution of the ichnoassemblages and genetic sequence of 1 st , 2 nd and 3 rd order.	216
Figure 7.4	Bagh Group succession of the Men River Valley showing the distribution of ichnoassemblages and genetic sequence of 1 st , 2 nd and 3 rd order.	218
Figure 7.5	Comparison of Cretaceous shoreline trajectory of the WLNV with the eustatic curve (Haq, 2014) based on systems tracts, sequence boundary, sequence stratigraphy surfaces, and systems tract boundary (Ruban, 2014). The Pre-Albian succession is characterized by unconventional High Amalgamation Systems Tract, while conventional systems tracts characterize Post-Aptian succession.	225
Figure 7.6	Conceptual sequence stratigraphic model of the Cretaceous Bagh Group rocks, WLNV showing stratal stacking pattern.	226

- Figure 8.1 Paleogeographical map showing the distribution of the continents during Albian-Cenomanian (Bardhan et al., 2002) and position of the 1. Eastern Desert (Egypt), 2. Mahajanga Basin (Madagascar), 3. Carnarvon Basin (Australia), 4. Saurashtra-Kachchh basins (India), 5. Jaisalmer-Barmer basins (India), 6. Cauvery Basin (India) and 7. Narmada Basin (India) (modified after Shitole et al., 2021). 230
- Figure 8.2 Generalized stratigraphic succession and the depositional environments of the Cretaceous Tethyan basins. A. Eastern Desert, Mahajanga, Carnarvon, Saurashtra, Kachchh and Narmada, and b. Jaisalmer, Barmer, and Cauvery (modified after Shitole et al., 2021). 234-235