
References

- Almrud, J. J., Oliveira, M. A., Kern, A. D., Grishin, N. V., Phillips, M. A., and Hackert, M. L. (2000). Crystal structure of human ornithine decarboxylase at 2.1 Å resolution: structural insights to antizyme binding. *J Mol Biol* 295, 7-16.
- Apweiler, R., Bairoch, A., and Wu, C. H. (2004). Protein sequence databases. *Curr Opin Chem Biol* 8, 76-80.
- Bachmair, A., Finley, D., and Varshavsky, A. (1986). In vivo half-life of a protein is a function of its amino-terminal residue. *Science* 234, 179-186.
- Baranov, P. V., Gurvich, O. L., Fayet, O., Prere, M. F., Miller, W. A., Gesteland, R. F., Atkins, J. F., and Giddings, M. C. (2001). RECODE: a database of frameshifting, bypassing and codon redefinition utilized for gene expression. *Nucleic Acids Res* 29, 264-267.
- Bercovich, Z., Rosenberg-Hasson, Y., Ciechanover, A., and Kahana, C. (1989). Degradation of ornithine decarboxylase in reticulocyte lysate is ATP-dependent but ubiquitin-independent. *J Biol Chem* 264, 15949-15952.
- Caughey, B., and Lansbury, P. T. (2003). Protofibrils, pores, fibrils, and neurodegeneration: separating the responsible protein aggregates from the innocent bystanders. *Annu Rev Neurosci* 26, 267-298.
- Chen, B., Retzlaff, M., Roos, T., and Frydman, J. (2011). Cellular strategies of protein quality control. *Cold Spring Harb Perspect Biol* 3, a004374.
- Chen, D., and Dou, Q. P. (2010). The ubiquitin-proteasome system as a prospective molecular target for cancer treatment and prevention. *Curr Protein Pept Sci* 11, 459-470.
- Ciechanover, A., Heller, H., Elias, S., Haas, A. L., and Hershko, A. (1980). ATP-dependent conjugation of reticulocyte proteins with the polypeptide required for protein degradation. *Proc Natl Acad Sci U S A* 77, 1365-1368.
- Ciechanover, A., Hod, Y., and Hershko, A. (1978). A heat-stable polypeptide component of an ATP-dependent proteolytic system from reticulocytes. *Biochem Biophys Res Commun* 81, 1100-1105.
- Cohavi, O., Tobi, D., and Schreiber, G. (2009). Docking of antizyme to ornithine decarboxylase and antizyme inhibitor using experimental mutant and double-mutant cycle data. *J Mol Biol* 390, 503-515.
- Cohen, F. E., and Kelly, J. W. (2003). Therapeutic approaches to protein-misfolding diseases. *Nature* 426, 905-909.
- da Fonseca, P. C., and Morris, E. P. (2008). Structure of the human 26S proteasome: subunit radial displacements open the gate into the proteolytic core. *J Biol Chem* 283, 23305-23314.

-
- Dai, C., Whitesell, L., Rogers, A. B., and Lindquist, S. (2007). Heat shock factor 1 is a powerful multifaceted modifier of carcinogenesis. *Cell* 130, 1005-1018.
- De Duve, C., and Wattiaux, R. (1966). Functions of lysosomes. *Annu Rev Physiol* 28, 435-492.
- DeLano, W. L. (2010). The PyMOL Molecular Graphics System (New York, Schroödinger, LLC).
- Doshi, A., Mishra, P., Sharma, M., and Prabha, C. R. (2014). Functional characterization of dosage-dependent lethal mutation of ubiquitin in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 14, 1080-1089.
- Etlinger, J. D., and Goldberg, A. L. (1977). A soluble ATP-dependent proteolytic system responsible for the degradation of abnormal proteins in reticulocytes. *Proc Natl Acad Sci U S A* 74, 54-58.
- Finn, R. D., Bateman, A., Clements, J., Coghill, P., Eberhardt, R. Y., Eddy, S. R., Heger, A., Hetherington, K., Holm, L., Mistry, J., *et al.* (2013). Pfam: the protein families database. *Nucleic Acids Res* 42, D222-230.
- Fonzi, W. A., and Sypherd, P. S. (1987). The gene and the primary structure of ornithine decarboxylase from *Saccharomyces cerevisiae*. *J Biol Chem* 262, 10127-10133.
- Forsthoefel, A. M., Pena, M. M., Xing, Y. Y., Rafique, Z., and Berger, F. G. (2004). Structural determinants for the intracellular degradation of human thymidylate synthase. *Biochemistry* 43, 1972-1979.
- Gandre, S., Bercovich, Z., and Kahana, C. (2002). Ornithine decarboxylase-antizyme is rapidly degraded through a mechanism that requires functional ubiquitin-dependent proteolytic activity. *Eur J Biochem* 269, 1316-1322.
- Gerner, E. W., and Meyskens, F. L., Jr. (2004). Polyamines and cancer: old molecules, new understanding. *Nat Rev Cancer* 4, 781-792.
- Ghoda, L., Phillips, M. A., Bass, K. E., Wang, C. C., and Coffino, P. (1990). Trypanosome ornithine decarboxylase is stable because it lacks sequences found in the carboxyl terminus of the mouse enzyme which target the latter for intracellular degradation. *J Biol Chem* 265, 11823-11826.
- Ghoda, L., van Daalen Wetters, T., Macrae, M., Ascherman, D., and Coffino, P. (1989). Prevention of rapid intracellular degradation of ODC by a carboxyl-terminal truncation. *Science* 243, 1493-1495.
- Glickman, M. H., and Ciechanover, A. (2002). The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction. *Physiol Rev* 82, 373-428.
- Godderz, D., Schafer, E., Palanimurugan, R., and Dohmen, R. J. (2011). The N-terminal unstructured domain of yeast ODC functions as a transplantable and replaceable ubiquitin-independent degron. *J Mol Biol* 407, 354-367.

-
- Gosink, M. M., and Vierstra, R. D. (1995). Redirecting the specificity of ubiquitination by modifying ubiquitin-conjugating enzymes. *Proc Natl Acad Sci U S A* 92, 9117-9121.
- Gupta, R., Hamasaki-Katagiri, N., White Tabor, C., and Tabor, H. (2001). Effect of spermidine on the in vivo degradation of ornithine decarboxylase in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A* 98, 10620-10623.
- Hannah, J., and Zhou, P. (2015). Methods: A small-molecule SMASh hit. *Nat Chem Biol* 11, 637-638.
- Hayashi, S., Murakami, Y., and Matsufuji, S. (1996). Ornithine decarboxylase antizyme: a novel type of regulatory protein. *Trends Biochem Sci* 21, 27-30.
- Heller, J. S., Fong, W. F., and Canellakis, E. S. (1976). Induction of a protein inhibitor to ornithine decarboxylase by the end products of its reaction. *Proc Natl Acad Sci U S A* 73, 1858-1862.
- Hershko, A. (1996). Lessons from the discovery of the ubiquitin system. *Trends Biochem Sci* 21, 445-449.
- Hoffman, D. W., Carroll, D., Martinez, N., and Hackert, M. L. (2005). Solution structure of a conserved domain of antizyme: a protein regulator of polyamines. *Biochemistry* 44, 11777-11785.
- Ichiba, T., Matsufuji, S., Miyazaki, Y., Murakami, Y., Tanaka, K., Ichihara, A., and Hayashi, S. (1994). Functional regions of ornithine decarboxylase antizyme. *Biochem Biophys Res Commun* 200, 1721-1727.
- Ivanov, I. P., Anderson, C. B., Gesteland, R. F., and Atkins, J. F. (2004). Identification of a new antizyme mRNA +1 frameshifting stimulatory pseudoknot in a subset of diverse invertebrates and its apparent absence in intermediate species. *J Mol Biol* 339, 495-504.
- Ivanov, I. P., Gesteland, R. F., and Atkins, J. F. (1998). A second mammalian antizyme: conservation of programmed ribosomal frameshifting. *Genomics* 52, 119-129.
- Ivanov, I. P., Gesteland, R. F., and Atkins, J. F. (2000a). Antizyme expression: a subversion of triplet decoding, which is remarkably conserved by evolution, is a sensor for an autoregulatory circuit. *Nucleic Acids Res* 28, 3185-3196.
- Ivanov, I. P., Rohrwasser, A., Terreros, D. A., Gesteland, R. F., and Atkins, J. F. (2000b). Discovery of a spermatogenesis stage-specific ornithine decarboxylase antizyme: antizyme 3. *Proc Natl Acad Sci U S A* 97, 4808-4813.
- Jariel-Encontre, I., Bossis, G., and Piechaczyk, M. (2008). Ubiquitin-independent degradation of proteins by the proteasome. *Biochim Biophys Acta* 1786, 153-177.
- Jariel-Encontre, I., Pariat, M., Martin, F., Carillo, S., Salvat, C., and Piechaczyk, M. (1995). Ubiquitylation is not an absolute requirement for degradation of c-Jun protein by the 26 S proteasome. *J Biol Chem* 270, 11623-11627.

-
- Jin, L., Williamson, A., Banerjee, S., Philipp, I., and Rape, M. (2008). Mechanism of ubiquitin-chain formation by the human anaphase-promoting complex. *Cell* 133, 653-665.
- Joshi, R. G., Kulkarni, S., and Ratna Prabha, C. (2015). Engineering degrons of yeast ornithine decarboxylase as vehicles for efficient targeted protein degradation. *Biochim Biophys Acta* 1850, 2452-2463.
- Ju, D., and Xie, Y. (2004). Proteasomal degradation of RPN4 via two distinct mechanisms, ubiquitin-dependent and -independent. *J Biol Chem* 279, 23851-23854.
- Kern, A. D., Oliveira, M. A., Coffino, P., and Hackert, M. L. (1999). Structure of mammalian ornithine decarboxylase at 1.6 Å resolution: stereochemical implications of PLP-dependent amino acid decarboxylases. *Structure* 7, 567-581.
- Koulov, A. V., LaPointe, P., Lu, B., Razvi, A., Coppinger, J., Dong, M. Q., Matteson, J., Laister, R., Arrowsmith, C., Yates, J. R., 3rd, and Balch, W. E. (2010). Biological and structural basis for Aha1 regulation of Hsp90 ATPase activity in maintaining proteostasis in the human disease cystic fibrosis. *Mol Biol Cell* 21, 871-884.
- Krappmann, D., Wulczyn, F. G., and Scheidereit, C. (1996). Different mechanisms control signal-induced degradation and basal turnover of the NF-kappaB inhibitor IkkappaB alpha in vivo. *Embo J* 15, 6716-6726.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-685.
- Lansbury, P. T., and Lashuel, H. A. (2006). A century-old debate on protein aggregation and neurodegeneration enters the clinic. *Nature* 443, 774-779.
- Li, X., and Coffino, P. (1992). Regulated degradation of ornithine decarboxylase requires interaction with the polyamine-inducible protein antizyme. *Mol Cell Biol* 12, 3556-3562.
- Li, X., and Coffino, P. (1993). Degradation of ornithine decarboxylase: exposure of the C-terminal target by a polyamine-inducible inhibitory protein. *Mol Cell Biol* 13, 2377-2383.
- Li, X., and Coffino, P. (1994). Distinct domains of antizyme required for binding and proteolysis of ornithine decarboxylase. *Mol Cell Biol* 14, 87-92.
- Li, X., Lonard, D. M., Jung, S. Y., Malovannaya, A., Feng, Q., Qin, J., Tsai, S. Y., Tsai, M. J., and O'Malley, B. W. (2006). The SRC-3/AIB1 coactivator is degraded in a ubiquitin- and ATP-independent manner by the REGgamma proteasome. *Cell* 124, 381-392.
- Manning, M. C., and Woody, R. W. (1991). Theoretical CD studies of polypeptide helices: examination of important electronic and geometric factors. *Biopolymers* 31, 569-586.

-
- Matsufuji, S., Matsufuji, T., Miyazaki, Y., Murakami, Y., Atkins, J. F., Gesteland, R. F., and Hayashi, S. (1995). Autoregulatory frameshifting in decoding mammalian ornithine decarboxylase antizyme. *Cell* 80, 51-60.
- Matsuzawa, S., Cuddy, M., Fukushima, T., and Reed, J. C. (2005). Method for targeting protein destruction by using a ubiquitin-independent, proteasome-mediated degradation pathway. *Proc Natl Acad Sci U S A* 102, 14982-14987.
- Melo, S. P., Yoshida, A., and Berger, F. G. (2010). Functional dissection of the N-terminal degron of human thymidylate synthase. *Biochem J* 432, 217-226.
- Mishra, P., Prabha, C. R., Rao Ch, M., and Volety, S. (2011). Q2N and S65D substitutions of ubiquitin unravel functional significance of the invariant residues Gln2 and Ser65. *Cell Biochem Biophys* 61, 619-628.
- Mishra, P., Volety, S., Rao Ch, M., and Prabha, C. R. (2009). Glutamate64 to glycine substitution in G1 beta-bulge of ubiquitin impairs function and stabilizes structure of the protein. *J Biochem* 146, 563-569.
- Moriishi, K., Okabayashi, T., Nakai, K., Moriya, K., Koike, K., Murata, S., Chiba, T., Tanaka, K., Suzuki, R., Suzuki, T., *et al.* (2003). Proteasome activator PA28gamma-dependent nuclear retention and degradation of hepatitis C virus core protein. *J Virol* 77, 10237-10249.
- Morimoto, R. I. (2008). Proteotoxic stress and inducible chaperone networks in neurodegenerative disease and aging. *Genes Dev* 22, 1427-1438.
- Munoz, V., and Serrano, L. (1994). Elucidating the folding problem of helical peptides using empirical parameters. *Nat Struct Biol* 1, 399-409.
- Murakami, Y., Matsufuji, S., Kameji, T., Hayashi, S., Igarashi, K., Tamura, T., Tanaka, K., and Ichihara, A. (1992). Ornithine decarboxylase is degraded by the 26S proteasome without ubiquitination. *Nature* 360, 597-599.
- Palanimurugan, R., Scheel, H., Hofmann, K., and Dohmen, R. J. (2004). Polyamines regulate their synthesis by inducing expression and blocking degradation of ODC antizyme. *Embo J* 23, 4857-4867.
- Pegg, A. E. (1988). Polyamine metabolism and its importance in neoplastic growth and a target for chemotherapy. *Cancer Res* 48, 759-774.
- Persson, L., Jeppsson, A., and Nasizadeh, S. (2003). Turnover of trypanosomal ornithine decarboxylases. *Biochem Soc Trans* 31, 411-414.
- Pickart, C. M., and Fushman, D. (2004). Polyubiquitin chains: polymeric protein signals. *Curr Opin Chem Biol* 8, 610-616.
- Porat, Z., Landau, G., Bercovich, Z., Krutauz, D., Glickman, M., and Kahana, C. (2008). Yeast antizyme mediates degradation of yeast ornithine decarboxylase by yeast but not by mammalian proteasome: new insights on yeast antizyme. *J Biol Chem* 283, 4528-4534.

Prabha, C. R., Mukherjee, S., Raman, R., and Kulkarni, S. (2012). The ends and means of artificially induced targeted protein degradation. *Appl Microbiol Biotechnol* 96, 1111-1123.

Raina, K., and Crews, C. M. (2010). Chemical inducers of targeted protein degradation. *J Biol Chem* 285, 11057-11060.

Ratnaprabha, C., and U. Sasidhar, Y. (1998). Conformational features of disulfide intact and reduced forms of hen egg white lysozyme in aqueous solution in the presence of trifluoroethanol (TFE): Implications for protein folding intermediates. *Journal of the Chemical Society, Faraday Transactions* 94, 3631-3637.

Rock, K. L., Gramm, C., Rothstein, L., Clark, K., Stein, R., Dick, L., Hwang, D., and Goldberg, A. L. (1994). Inhibitors of the proteasome block the degradation of most cell proteins and the generation of peptides presented on MHC class I molecules. *Cell* 78, 761-771.

Rosenberg-Hasson, Y., Bercovich, Z., Ciechanover, A., and Kahana, C. (1989). Degradation of ornithine decarboxylase in mammalian cells is ATP dependent but ubiquitin independent. *Eur J Biochem* 185, 469-474.

Roy, A., Kucukural, A., and Zhang, Y. (2010). I-TASSER: a unified platform for automated protein structure and function prediction. *Nat Protoc* 5, 725-738.

Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4, 406-425.

Sambrook, J., Fritsch, E.F., Maniatis, T., (1989). *Molecular cloning: a laboratory manual*, 2 edn (Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press).

Sasidhar, Y. U., and Prabha, C. R. (2000). Conformational features of reduced and disulfide intact forms of hen egg white lysozyme in aqueous solution in presence of 3-chloro-1, 2-propanediol and dioxane: implications for protein folding intermediates. *Indian J Biochem Biophys* 37, 97-106.

Sawkar, A. R., D'Haese, W., and Kelly, J. W. (2006). Therapeutic strategies to ameliorate lysosomal storage disorders--a focus on Gaucher disease. *Cell Mol Life Sci* 63, 1179-1192.

Schrader, E. K., Harstad, K. G., and Matouschek, A. (2009). Targeting proteins for degradation. *Nat Chem Biol* 5, 815-822.

Sheaff, R. J., Singer, J. D., Swanger, J., Smitherman, M., Roberts, J. M., and Clurman, B. E. (2000). Proteasomal turnover of p21Cip1 does not require p21Cip1 ubiquitination. *Mol Cell* 5, 403-410.

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Soding, J., *et al.* (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 7, 539.

Snapir, Z., Keren-Paz, A., Bercovich, Z., and Kahana, C. (2009). Antizyme 3 inhibits polyamine uptake and ornithine decarboxylase (ODC) activity, but does not stimulate ODC degradation. *Biochem J* 419, 99-103, 101 p following 103.

Sreerama, N., and Woody, R. W. (2000). Estimation of protein secondary structure from circular dichroism spectra: comparison of CONTIN, SELCON, and CDSSTR methods with an expanded reference set. *Anal Biochem* 287, 252-260.

Su, V., Nakagawa, R., Koval, M., and Lau, A. F. (2010). Ubiquitin-independent proteasomal degradation of endoplasmic reticulum-localized connexin43 mediated by CIP75. *J Biol Chem* 285, 40979-40990.

Swapnali Kulkarni: Studies on degradation determinant signals of ornithone decarboxylase. Ph.D. Thesis 2012

Tabor, C. W., and Tabor, H. (1984). Polyamines. *Annu Rev Biochem* 53, 749-790.

Tachihara, K., Uemura, T., Kashiwagi, K., and Igarashi, K. (2005). Excretion of putrescine and spermidine by the protein encoded by YKL174c (TPO5) in *Saccharomyces cerevisiae*. *J Biol Chem* 280, 12637-12642.

Tamura, K., Dudley, J., Nei, M., and Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* 24, 1596-1599.

Tarcsa, E., Szymanska, G., Lecker, S., O'Connor, C. M., and Goldberg, A. L. (2000). Ca²⁺-free calmodulin and calmodulin damaged by in vitro aging are selectively degraded by 26 S proteasomes without ubiquitination. *J Biol Chem* 275, 20295-20301.

Teixeira, M. C., Cabrito, T. R., Hanif, Z. M., Vargas, R. C., Tenreiro, S., and Sa-Correia, I. Yeast response and tolerance to polyamine toxicity involving the drug : H⁺ antiporter Qdr3 and the transcription factors Yap1 and Gcn4. *Microbiology* 157, 945-956.

Thrower, J. S., Hoffman, L., Rechsteiner, M., and Pickart, C. M. (2000). Recognition of the polyubiquitin proteolytic signal. *Embo J* 19, 94-102.

UniProtConsortium, T. (2014). UniProt: a hub for protein information. *Nucleic Acids Res* 43, D204-212.

Varshavsky, A. (1996). The N-end rule: functions, mysteries, uses. *Proc Natl Acad Sci U S A* 93, 12142-12149.

Weissmann, A. M. (1997). Regulating protein degradation by ubiquitination. *Immunology Today* 8, 189-196.

Wiggins, C. M., Tsvetkov, P., Johnson, M., Joyce, C. L., Lamb, C. A., Bryant, N. J., Komander, D., Shaul, Y., and Cook, S. J. (2011). BIM(EL), an intrinsically disordered protein, is degraded by 20S proteasomes in the absence of poly-ubiquitylation. *J Cell Sci* 124, 969-977.

-
- Yang, J., Yan, R., Roy, A., Xu, D., Poisson, J., and Zhang, Y. (2015). The I-TASSER Suite: protein structure and function prediction. *Nat Methods* 12, 7-8.
- Yu, H., Kaung, G., Kobayashi, S., and Kopito, R. R. (1997). Cytosolic degradation of T-cell receptor alpha chains by the proteasome. *J Biol Chem* 272, 20800-20804.
- Yukse, K., Chen, W. L., Chien, D., and Ou, J. H. (2009). Ubiquitin-independent degradation of hepatitis C virus F protein. *J Virol* 83, 612-621.
- Zhang, M., MacDonald, A. I., Hoyt, M. A., and Coffino, P. (2004). Proteasomes begin ornithine decarboxylase digestion at the C terminus. *J Biol Chem* 279, 20959-20965.
- Zhang, M., Pickart, C. M., and Coffino, P. (2003). Determinants of proteasome recognition of ornithine decarboxylase, a ubiquitin-independent substrate. *Embo J* 22, 1488-1496.
- Zhang, Y. (2008). I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics* 9, 40.