

---

## REFERENCES

- Adhikary L, et al. Abnormal p38mitogen-activated protein kinase signalling in human and experimental diabetic nephropathy. *Diabetologia* 2004; 47:1210–1222.
- Aguirre V, et al. Phosphorylation of Ser307 in insulin receptor substrate-1 blocks interactions with the insulin receptor and inhibits insulin action. *J Biol Chem*. 2002; 277:1531–1537.
- Aguirre V, et al. The c-Jun NH(2)- terminal kinase promotes insulin resistance during association with insulin receptor substrate-1 and phosphorylation of Ser(307). *J Biol Chem* 2000; 275: 9047-9054.
- Ahima RS, Flier JS Adipose tissue as an endocrine organ. *Trends Endocrinol Metab* 2000;11: 327–332.
- Alessi MC, et al.. Production of plasminogen activator inhibitor 1 by human adipose tissue: possible link between visceral fat accumulation and vascular disease. *Diabetes* 1997; 46: 860-867.
- Arai T, et al. Increased plasma cholesteryl ester transfer protein in obese subjects. A possible mechanism for the reduction of serum HDL cholesterol levels in obesity. *Arterioscler Thromb* 1994; 14:1129–1136.
- Araki T, et al. Effect of adiponectin on carotid arterial stiffness in type 2 diabetic patients treated with pioglitazone and metformin. *Metabolism*. 2006; 55: 996–1001.
- Arita Y, et al. Adipocyte-derived plasma protein, adiponectin, suppresses lipid accumulation and class A scavenger receptor expressionin human monocyte- derived macrophages. *Circulation* 2001;103: 1057–1063
- Arita Y, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophy Res Commun* 1999; 257: 79-83.
- Arkan MC, et al. IKK-beta links infl ammation to obesity-induced insulin resistance. *Nat Med*, 2005; 11:191–198.
- Aso Y, et al. Comparison of serum high-molecular weight (HMW) adiponectin with total adiponectin concentrations in type 2 diabetic patients with coronary artery disease using a novel enzyme-linked immunosorbent assay to detect HMW adiponectin. *Diabetes* 2006; 55: 1954-1960.
- Atzmon G, et al. Differential gene expression between visceral and subcutaneous fat depots. *Horm Metab Res* 2002;34:622–628.
- Balagopal P, et al. Reduction of Elevated Serum Retinol Binding Protein (RBP4) in Obese Children by Lifestyle Intervention: Association with Sub-clinical Inflammation *J Clin Endocrin Metab*.2007; 92:1971-1974.

- Baratta R, et al. Adiponectin relationship with lipid metabolism is independent of body fat mass: evidence from both cross-sectional and intervention studies. *J Clin Endocrinol Metab* 2004; 89: 2665-2671.
- Barrett-Connor E and Bush TL. Estrogen and coronary heart disease in women. *JAMA* 1991; 265: 1861–1867,
- Barter PJ, et al. Cholesteryl ester transfer protein: a novel target for raising HDL and inhibiting atherosclerosis. *Arterioscler Thromb Vasc Biol* 2003; 23:160–167.
- Barzilai N, et al. Surgical removal of visceral fat reverses hepatic insulin resistance. *Diabetes*. 1999; 48: 94 –98.
- Bastard JP, et al. Elevated levels of interleukin 6 are reduced in serum and subcutaneous adipose tissue of obese women after weight loss. *J Clin Endocrinol Metab* 2000; 85: 3338-3342.
- Basualdo CG, et al. Vitamin A (retinol) status of First Nation adults with non-insulin-dependent diabetes mellitus. *J Am Coll Nutr* 1997; 16: 39-45.
- Baynes JW, Thorpe SR. Role of oxidative stress in diabetic complications: a new perspective on an old paradigm. *Diabetes*. 1999; 48:1–9.
- Bennett CN, et al. Regulation of osteoblastogenesis and bone mass by Wnt10b. *Proc Natl Acad Sci U S A*. 2005; 102: 3324–3329.
- Bensaid M et al. The cannabinoid CB1 receptor antagonist SR141716 increases Acrp30 mRNA expression in adipose tissue of obese fa/fa rats and in cultured adipocyte cells. *Mol Pharmacol* 2003; 63: 908–914.
- Berg AH, et al. Adipocyte differentiation induces dynamic changes in NF-kappaB expression and activity. *Am J Physiol Endocrinol Metab*. 2004; 287: E1178–E1188.
- Berg AH, et al. The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med* 2001; 7: 947–953
- Berg AH, Scherer PE. Adipose Tissue, Inflammation, and Cardiovascular Disease. *Circ Res*. 2005; 96: 939-49.
- Berger J and Moller DE. The mechanisms of action of PPARs. *Annu. Rev. Med*. 2002; 53, 409–435
- Bergman RN, Mittelman SD. Central role of the adipocyte in insulin resistance. *J Basic Clin Physiol Pharmacol* 1998; 9:205–221.
- Bermúdez-Siva FJ et al. Activation of cannabinoid CB1 receptors induces glucose intolerance in rats. *Eur J Pharmacol* 2006; 531: 282–284.
- Bhagar K and Vallance P. Inflammatory cytokines impairs endothelium- dependent dilatation in human veins in vivo. *Circulation* 1997; 96: 3042– 3047
- Bierman EL. George Lyman Duff Memorial Lecture. Atherogenesis in diabetes. *Arterioscler Thromb* 1992;12: 647–656.

- Blake, GJ, Ridker PM. Novel Clinical Markers of Vascular Wall Inflammation. *Circ Res.* 2001; 89: 763-771
- Bluher M, et al. Dysregulation of the peripheral and adipose tissue endocannabinoid system in human abdominal obesity. *Diabetes* 2006; 55: 3053–3060.
- Bodzioch M, et al. The gene encoding ATP-binding cassette transporter 1 is mutated in Tangier disease. *Nat Genet* 1999; 22: 347–551
- Bokarewa M, et al. Resistin an adipokine with potent pro-inflammatory properties. *J Immunol* 2005;174: 5789–5795.
- Boring L, et al. Decreased lesion formation in CCR2 ko mice reveals a role for chemokines in the initiation of atherosclerosis. *Nature.* 1998; 394: 894–897.
- Bottner A, et al. Gender differences of adiponectin levels develop during the progression of puberty and are related to serum androgen levels. *J Clin Endocrinol Metab* 2004; 89: 4053-4061.
- Bouhrel MA, et al. PPARgamma activation primes human monocytes into alternative M2 macrophages with antiinflammatory properties. *Cell Metab*, 2007; 6: 137–143.
- Branen L, et al. Inhibition of tumor necrosis factor-alpha reduces atherosclerosis in apolipoprotein E knockout mice. *Arterioscler Thromb Vasc Biol.* 2004; 24:2137–2142.
- Brendle DC, et al. Effects of exercise rehabilitation on endothelial reactivity in older patients with peripheral arterial disease. *Am J Cardiol* 2001; 87: 324–329.
- Brentano F, et al. Pre B cell colony enhancing factor/ visfatin a new marker of inflammation in rheumatoid arthritis with proinflammatory and matrix degrading activities. *Arthritis. Rheum.* 2007; 56: 2829-2839.
- Brochu M, et al. Is there a threshold of visceral fat loss that improves the metabolic profile in obese postmenopausal women? *Metabolism.* 2003; 52: 599–604.
- Bruun JM, et al. Regulation of adiponectin by adipose tissue-derived cytokines: in vivo and in vitro investigations in humans. *Am J Physiol Endocrinol Metab* 2003; 285: E527-33.
- Burnett MS, et al. The potential role of resistin in atherogenesis. *Atherosclerosis* 2005;182: 241–8.
- Busso N, et al. Leptin signaling deficiency impairs humoral and cellular immune responses and attenuates experimental arthritis. *J Immunol* 2002;168:875–882.
- Butte NF, et al. Genetic and environmental factors influencing fasting serum adiponectin in Hispanic children. *J Clin Endocrinol Metab* 2005; 90: 4170-4176.
- Cai D, et al. Local and systemic insulin resistance resulting from hepatic activation of IKK-beta and NF-kappaB. *Nat Med*, 2005 ; 11:183–190.
- Cai Q, et al. Growth factors induce monocyte binding to vascular smooth muscle cells: implications for monocyte retention in atherosclerosis *Am J Physiol* 2004; 287:C707–C714,

- Calabro P, et al. Release of C-reactive protein in response to inflammatory cytokines by human adipocytes: linking obesity to vascular inflammation. *J Am Coll Cardiol* 2005; 46: 1112-1113.
- Caldefie-Chezet F, et al. Leptin regulates functional capacities of polymorphonuclear neutrophils. *Free Radic Res* 2003; 37: 809-814.
- Caldefie-Chezet F, et al. Leptin: a potential regulator of polymorphonuclear neutrophil bactericidal action? *J Leukoc Biol* 2001; 69: 414-418.
- Calles-Escandon J, et al. Induction of hyperinsulinemia combined with hyperglycemia and hypertriglyceridemia increases plasminogen activator inhibitor 1 in blood in normal human subjects. *Diabetes* 1998, 47:290-293.
- Campbell JH, Campbell GR: Cell biology of atherosclerosis. *J Hypertension* 1994, 12:S129-S132.
- Campbell LA, Kuo CC. Chlamydia pneumoniae: an infectious risk factor for atherosclerosis? *Nat Rev Microbiol*. 2004; 2: 23-32.
- Canault M, et al. Exclusive expression of transmembrane TNF-alpha in mice reduces the inflammatory response in early lipid lesions of aortic sinus. *Atherosclerosis*. 2004;172: 211-218.
- Cao Z, et al. Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes Dev*. 1991; 5:1538-1352
- Cawthorn WP, et al. Tumour necrosis factor-alpha inhibits adipogenesis via a beta-catenin/TCF4(TCF7L2)-dependent pathway. *Cell Death Differ*. 2007;14:1361-1373.
- Charriere G, et al. Preadipocyte conversion to macrophage. Evidence of plasticity. *J Biol Chem*. 2003;278:9850 -9855.
- Chawla A, et al. PPAR-gamma dependent and independent effects on macrophage-gene expression in lipid metabolism and inflammation. *Nat Med* 2001; 7: 48-52.
- Chen H, et al. Adiponectin stimulates production of nitric oxide in vascular endothelial cells. *J Biol Chem*. 2003; 278: 45021- 6.
- Chen Z, et al. Troglitazone inhibits atherosclerosis in apoli- *Arterioscler Thromb Vasc Biol* 2001; 21:372-377.
- Cho YM, et al. Plasma retinol-binding protein-4 concentrations are elevated in human subjects with impaired glucose tolerance and type 2 diabetes. *Diabetes Care* 2006; 29: 2457-2461.
- Choi KC, et al. Effect of PPAR-alpha and -gamma agonist on the expression of visfatin, adiponectin, and TNF-alpha in visceral fat of OLETF rats. *Biochem Biophys Res Commun* 2005; 28: 747-753
- Chowdhury B, et al. A multicompartiment body composition technique based on computerized tomography. *Int J Obes Relat Metab Disord*. 1994;18:219-234.

- Christiansen T, et al. Monocyte chemoattractant protein-1 is produced in isolated adipocytes, associated with adiposity and reduced after weight loss in morbid obese subjects. *Int J Obes* 2005; 29: 146-150.
- Chu X, et al. Suppression of adipogenesis program in cultured preadipocytes transfected stably with cyclooxygenase isoforms. *Biochim Biophys Acta*. 2009; 1791: 273-280.
- Chu X, et al. Up-regulation of adipogenesis in adipocytes expressing stably cyclooxygenase-2 in the antisense direction. *Prostaglandins Other Lipid Mediat*. 2010; 91:1-9.
- Chung S, et al Preadipocytes Mediate Lipopolysaccharide-Induced Inflammation and Insulin Resistance in Primary Cultures of Newly Differentiated Human Adipocytes. *Endocrinology*. 2006; 147: 5340–5351.
- Cinti S, et al. Adipocyte death defines macrophage localization and function in adipose tissue of obese mice and humans. *J Lipid Res*, 2005; 46: 2347–2355.
- Clarkson P, et al. Exercise training enhances endothelial function in young men. *J Am Coll Cardiol* 1999; 33:1379–1385.
- Cnop M, et al. Relationship of adiponectin to body fat distribution, insulin sensitivity and plasma lipoproteins: evidence for independent roles of age and sex. *Diabetologia* 2003; 46: 459-469.
- Collins AR, et al. Troglitazone inhibits formation of early atherosclerotic lesions in diabetic and nondiabetic low density lipoprotein receptor-deficient mice. *Arterioscler Thromb Vasc Biol* 2001; 21:365–371
- Combs, T.P. et al. A transgenic mouse with a deletion in the collagenous domain of adiponectin displays elevated circulating adiponectin and improved insulin sensitivity. *Endocrinology* 2004; 145: 367–383
- Considine RV, et al. Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med* 1996; 334: 292–295.
- Cooke JP and Oka RK. Does leptin cause vascular disease? *Circulation* 2002;106: 1904–1905.
- Cota D et al. The endogenous cannabinoid system affects energy balance via central orexigenic drive and peripheral lipogenesis. *J Clin Invest* 2003; 112: 423–431.
- Cousin B, et al. A role for preadipocytes as macrophage-like cells. *FASEB J*, 1999 ; 13: 305–312.
- Coutts AA, Izzo AA. The gastrointestinal pharmacology of cannabinoids: an update. *Curr Opin Pharmacol* 2004; 4: 572–579.
- Cowie CC, et al. Prevalence of diabetes and impaired fasting glucose in adults—United States, 1999–2000. *MMWR* 2003; 52: 833–837.
- Croci T, et al. Role of cannabinoid CB1 receptors and tumor necrosis factor- $\alpha$  in the gut and systemic anti-inflammatory activity of SR 141716 (Rimonabant) in rodents. *Br. Journal. Pharmacol*. 2003; 140:115–122.

- Dahl TB, et al. Increased expression of visfatin in macrophage of human unstable carotid and coronary atherosclerosis. *Circulation* 2007; 115: 972-980.
- Dandona P, Aljada A. A rational approach to pathogenesis and treatment of type 2 diabetes mellitus, insulin resistance, inflammation, and atherosclerosis. *Am. J. Cardiol.* 2002; 90: 27-33.
- Dandona P, et al. Tumor necrosis factor-alpha in sera of obese patients: fall with weight loss. *J Clin Endocrinol Metab* 1998; 83:2907-2910
- Danforth E Jr. Failure of adipocyte differentiation causes type II diabetes mellitus? *Nat Genet.* 2000; 26: 13.
- de Lemos JA, et al. Association between plasma levels of monocyte chemoattractant protein-1 and long-term clinical outcomes in patients with acute coronary syndromes. *Circulation* 2003; 107: 690-695.
- de Souza CJ, et al. Effects of pioglitazone on adipose tissue remodeling within the setting of obesity and insulin resistance. *Diabetes* 2001; 50: 1863-1871.
- Dekker JM, et al. Prognostic value of adiponectin for cardiovascular disease and mortality. *J Clin Endocrinol Metab* 2008; 93: 1489-1496.
- Delaigle AM, et al. Induction of adiponectin in skeletal muscle by inflammatory cytokines: in vivo and in vitro studies. *Endocrinology* 2004;145, 5589-5597.
- Despres JP, et al. Effects of rimonabant on metabolic risk factors in overweight patients with dyslipidemia. *N Engl J Med* 2005; 353:2121-2134.
- Devaraj S, Jialal I. Alpha tocopherol supplementation decreases serum C-reactive protein and monocyte interleukin-6 levels in normal volunteers and type 2 diabetic patients. *Free Radic Biol Med* 2000; 29:790-2.
- Di Marzo V, Matias I. Endocannabinoid control of food intake and energy balance. *Nat Neurosci* 2005; 8: 585-589.
- Diep QN, et al. Long-term effects of the PPAR gamma activator pioglitazone on cardiac inflammation in stroke-prone spontaneously hypertensive rats. *Can J Physiol Pharmacol.* 2004; 82: 976-985.
- Dietze-Schroeder D, et al. Autocrine action of adiponectin on human fat cells prevents the release of insulin resistance-inducing factors. *Diabetes* 2005; 54: 2003-2011.
- Dol-Gleizes F, et al. Rimnabant, a Selective Cannabinoid CB1 Receptor Antagonist, Inhibits Atherosclerosis in LDL Receptor-Deficient Mice. *Arteriosclerosis Thrombosis and Vascular Biology* 2009; 29: 12-18.
- Doll R and Hill AB. Lung cancer and other causes of death in relation to smoking: a second report on the mortality of British doctors. *Br Med* 1956; J 2: 1071-1081

- Dusserre E, et al. Differences in mRNA expression of the proteins secreted by the adipocytes in human subcutaneous and visceral adipose tissues. *Biochim Biophys Acta* 2000; 1500: 88–96
- Dutta K, et al. Repeated systemic *Escherichia coli* infection enhances anti-oxidant response in hypercholesterolemic mice inducing cardiovascular inflammation. *Inflammation* 2009; 32: 89–98.
- Eckel RH. Obesity and heart disease: a statement for healthcare professionals from the Nutrition Committee, American Heart Association. *Circulation* 1997; 96: 3248–50.
- Ehtisham NZ: Do resistin resist insulin? *Curr Sci* 2002, 83:1190–1191.
- Elgazar-Carmon V, Rudich A, Hadad N, et al. 2008. Neutrophils transiently infiltrate intra-abdominal fat early in the course of high fat feeding. *J Lipid Res*, 49:1894–903.
- Engelberts I, et al. Evidence for different effects of soluble TNF-receptors on various TNF measurements in human biological fluids. *Lancet* 1991; 338: 515–516.
- Engell SJ, et al. Activation of the peripheral endocannabinoid system in human obesity. *Diabetes* 2005; 54: 2838–2843
- English JP, et al. Tobacco and coronary disease. *JAMA* 1940;115: 1327–1329
- Eriksson P, et al. Adipose tissue secretion of plasminogen activator inhibitor-1 in non-obese and obese individuals. *Diabetologia* 1998; 41: 65–71.
- Eriksson P, et al. Regional variation in plasmin-ogen activator inhibitor-1 expression in adipose tissue from obese individuals. *Thromb Haemost* 2000; 83: 545–548
- Erikstrup C, et al. Retinol-Binding Protein 4 and Insulin Resistance. *N Engl J Med* 2006; 355:1393–1394
- Esposito K, et al. Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women: a randomized trial. *JAMA* 2003; 289: 1799–1804.
- Faggioni R, et al. Leptin deficiency enhances sensitivity to endotoxin induced lethality. *Am J Physiol* 1999; 276:136–142.
- Faggioni R, et al. Leptin regulation of the immune response and the immunodeficiency of malnutrition. *FASEB J* 2001; 14: 2565–2571.
- Fain JN, et al. Comparison of the release of adipokines by adipose tissue, adipose tissue matrix, and adipocytes from visceral and subcutaneous abdominal adipose tissues of obese humans. *Endocrinology*. 2004;145:2273–2282.
- Farooqi IS, et al. Beneficial effects of leptin on obesity, T cell hyporesponsiveness, and neuroendocrine/metabolic dysfunction of human congenital leptin deficiency. *J Clin Invest* 2002;110:1093–1103.
- Fasshauer M, et al. Adiponectin gene expression and secretion is inhibited by interleukin-6 in 3T3-L1 adipocytes. *Biochem. Biophys. Res. Commun.* 2003; 301, 1045–1050

- Fehmann H, Heyn J. Plasma resistin levels in patients with type 1 and type 2 diabetes mellitus and in healthy controls. *Horm Metab Res* 2002; 34: 671-673.
- Filippatos TD, et al. Increased plasma levels of visfatin/pre-B cell colony-enhancing factor in obese and overweight patients with metabolic syndrome. *J Endocrinol Invest* 2007; 30:323–326.
- Fleming RM. The Pathogenesis of Vascular Disease. *Textbook of Angiology*. New York: Springer-Verlag; 1999:787-798.
- Flier JS, et al. Severely impaired adiponin expression in genetic and acquired obesity. *Science*, 1987; 237: 405-408
- Flier JS. The adipocyte: storage depot or node on the energy information superhighway? *Cell*. 1995; 80:15– 18.
- Flier JS: Obesity wars: Molecular progress confronts an expanding epidemic. *Cell* 2004; 116: 337–350.
- Folsom AR, et al. Impact of weight loss on plasminogen activator inhibitor (PAI- 1), factor VII, and other hemostatic factors in moderately overweight adults. *Arterioscler Thromb* 1993; 13: 162–169
- Forman BM, et al. 15-deoxyprostaglandin J2 is a ligand for the adipocyte determination factor PPAR $\gamma$ . *Cell* 1995; 83, 803–812
- Forst T, et al. Pleiotropic and anti-inflammatory effects of pioglitazone precede the metabolic activity in type 2 diabetic patients with coronary artery disease. *Atherosclerosis* 2008; 197: 311-317.
- Freeman DJ, et al. C-reactive protein is an independent predictor of risk for the development of diabetes in the West of Scotland Coronary Prevention Study. *Diabetes* 2002; 51: 1596-600.
- Fried SK, et al. Omental and subcutaneous adipose tissues of obese subjects release interleukin-6: depot difference and regulation by glucocorticoid. *J Clin Endocrinol Metab* 1998; 83: 847–850
- Friedman J. Fat in all the wrong places. *Nature* 2002, 415:268–269.
- Fruzzetti F, et al. Effect of long-term naltrexone treatment on endocrine profile, clinical features, and insulin sensitivity in obese women with polycystic ovary syndrome. *Fertil Steril* 2002; 77:936–944.
- Fukuhara A, et al Visfatin: a protein secreted by visceral fat that mimics the effects of insulin. *Science* 2005; 307: 426–430
- Garbriely I, et al. Removal of visceral fat prevents insulin resistance and glucose intolerance of aging: an adipokine-mediated process? *Diabetes*. 2002; 51: 2951–2958.
- Garg A. Lipodystrophies. *Am J Med*. 2000;108:143–52.



- Garlichs CD, et al. Upregulation of CD40 and CD40 ligand (CD154) in patients with moderate hypercholesterolemia. *Circulation* 2001; 104: 2395-2400.
- Gary-Boboo M et al. The cannabinoid CB1 receptor antagonist rimonabant (SR141716) inhibits cell proliferation and increases markers of adipocyte maturation in cultured mouse 3T3 F442A preadipocytes. *Mol Pharmacol* 2006; 69: 471-478.
- Gasperi V, et al. Endocannabinoids in adipocytes during differentiation and their role in glucose uptake. *Cell. Mol. Life. Sci.* 2007; 64: 219-229.
- Gaziano JM. Epidemiology of risk factor reduction. In: *Vascular Medicine*, edited by Loscalzo J, Creagher M, and Dzau V. Boston, MA: Little Brown, 1996; p. 569-586.
- Gearing AJH, et al. Soluble forms of vascular adhesion molecules, E-selectin, ICAM-1, and VCAM-1. Pathological significance. *Ann N Y Acad Sci* 1992, 667:324-331.
- Gerhardt CC, et al. Chemokines control fat accumulation and leptin secretion by cultured human adipocytes. *Mol Cell Endocrinol* 2001; 175: 81-92.
- Gerstmayr B, et al. Identification of RELM gamma, a novel resistin-like molecule with a distinct expression pattern. *Genomics* 2003; 81: 588-595.
- Ghanim H, et al. Circulating mononuclear cells in the obese are in a proinflammatory state. *Circulation*, 2004; 110:1564-1571.
- Ghanim H, et al. Low-Dose Rosiglitazone Exerts an Antiinflammatory Effect with an Increase in Adiponectin Independently of Free Fatty Acid Fall and Insulin Sensitization in Obese Type 2 Diabetics. *J Clin Endocrinol Metab* 2006; 91: 3553-3558.
- Ghanim H, et al. Suppression of nuclear factor-kappaB and stimulation of inhibitor kappaB by troglitazone: Evidence for an anti-inflammatory effect and a potential antiatherosclerotic effect in the obese. *J Clin Endocrinol Metab* 2001; 86: 1306-1312.
- Giacchetti G, et al. Overexpression of the renin-angiotensin system in human visceral adipose tissue in normal and overweight subjects. *Am J Hypertens* 2002; 15:381-388.
- Giltay EJ, et al. Visceral fat accumulation is an important determinant of PAI-1 levels in young, nonobese men and women: modulation by cross-sex hormone administration. *Arterioscler Thromb Vasc Biol* 1998; 18: 1716-1722.
- Giugliano G, et al. Effect of liposuction on insulin resistance and vascular inflammatory markers in obese women. *Br J Plast Surg* 2004; 57: 190-194.
- Gordon T, et al. High density lipoprotein as a protective factor against coronary heart disease. The Framingham Study. *Am J Med* 1977; 62: 707-714
- Gotto AM Jr and Farmer JA. Risk factors for coronary artery disease. In: *Heart Disease: A Textbook of Cardiovascular Medicine*, edited by Braunwald E. Philadelphia, PA: Saunders, 1988, p.1153-1190.
- Gotto AM Jr and Grundy SM. Lowering LDL cholesterol: questions from recent meta-analyses and subset analyses of clinical trial data. Issues from the Interdisciplinary Council on

- 
- Reducing the Risk for Coronary Heart Disease, ninth Council meeting. *Circulation* 1999; 99: E1–E7
- Graham TE, et al. Retinol-binding protein 4 and insulin resistance in lean, obese, and diabetic subjects. *N Engl J Med* 2006; 354: 2552–2563.
- Greenberg AS, et al. Interleukin 6 reduces lipoprotein lipase activity in adipose tissue of mice in vivo and in 3T3-L1 adipocytes: a possible role for interleukin-6 in cancer cachexia. *Cancer Res* 1992; 52: 4113–4116.
- Gregoire FM, et al. Understanding adipocyte differentiation. *Physiol Rev.* 1998; 78: 783–809.
- Griendling KK, et al. Angiotensin II signaling in vascular smooth muscle: new concepts. *Hypertension.* 1997; 29: 366–373.
- Grundy SM. Obesity, metabolic syndrome, and cardiovascular disease. *J Clin Endocrinol Metab* 2004; 89: 2595–2600.
- Gu L, et al. Absence of monocyte chemoattractant protein-1 reduces atherosclerosis in low-density lipoprotein-deficient mice. *Mol Cell.* 1998; 2: 275–281.
- Gustafson B, et al. Inflamed adipose tissue: a culprit underlying the metabolic syndrome and atherosclerosis. *Arterioscler Thromb Vasc Biol* 2007; 27: 2276–2283,
- Gustafson B, Smith U. Cytokines promote Wnt signaling and inflammation and impair the normal differentiation and lipid accumulation in 3T3-L1 preadipocytes. *J Biol Chem.* 2006; 281: 9507–9516.
- Gustafson B, Smith U. Cytokines promote Wnt signaling and inflammation and impair the normal differentiation and lipid accumulation in 3T3-L1 preadipocytes. *J Biol Chem* 2006; 281: 9507–9516
- Haffner SM, et al. Cardiovascular risk factors in confirmed prediabetic individuals. Does the clock for coronary heart disease start ticking before the onset of clinical disease? *JAMA* 1990; 263: 2893–2898
- Haffner SM, et al. Effect of rosiglitazone treatment on nontraditional markers of cardiovascular disease in patients with type 2 diabetes mellitus. *Circulation* 2002;106: 679–684.
- Halaas JL, et al. Weight-reducing effects of the plasma protein encoded by the obese gene. *Science* 1995; 269: 543–546.
- Hammarstedt A, et al. High circulating levels of RBP4 and mRNA levels of aP2, PGC-1 $\alpha$  and UCP-2 predict improvement in insulin sensitivity following pioglitazone treatment of drug-naïve type 2 diabetic subjects. *J. Intern Med.* 2008; 263: 440–449.
- Hammarstedt A, et al. Reduced expression of PGC-1 and insulin-signaling molecules in adipose tissue is associated with insulin resistance. *Biochem Biophys Res Commun.* 2003;301:578–582.
- Hammond EC and Horn D. Smoking and death rates: report on forty-four months of follow-up of 187,783 men. 2. Death rates by cause. *J Am Med Assoc* 1958;166: 1294–1308

- Hamsten A, et al. Increased plasma levels of a rapid inhibitor of tissue plasminogen activator in young survivors of myocardial infarction. *N Engl J Med* 1985; 313: 1557-1563.
- Han SH, et al. Antiatherosclerotic and Anti-Insulin Resistance Effects of Adiponectin: Basic and Clinical Studies. *Prog Cardiovasc Dis* 2009; 52:126-140.
- Hansson G, Libby P. The role of the lymphocyte. In: Fuster V, Ross R, Topol E, eds. *Atherosclerosis and Coronary Artery Disease*. New York, NY: Lippincott-Raven; 1996:557-568.
- Hardardo I, et al. Effects of endotoxin and cytokines on lipid metabolism. *Curr Opin Lipidol* 1994; 5: 207-215.
- Hauer AD, et al. Delivery of *Chlamydia pneumoniae* to the vessel wall aggravates atherosclerosis in LDLr KO mice. *Cardiovasc Res*. 2006; 69: 280-288.
- Havel PJ, et al. Relationship of plasma leptin to plasma insulin and adiposity in normal weight and overweight women: effects of dietary fat content and sustained weight loss. *J Clin Endocrinol Metab* 1996, 81:406-413.
- Heilbronn LK, et al. Energy restriction and weight loss on very-low-fat diets reduce C-reactive protein concentrations in obese, healthy women. *Arterioscler Thromb Vasc Biol* 2001; 21: 968-970.
- Heinonen MV, et al. Apelin, orexin-A and leptin plasma levels in morbid obesity and effect of gastric banding. *Regul. Pept.* 2005;130, 7-13.
- Herder C, et al. Low-grade inflammation, obesity, and insulin resistance in adolescents. *J Clin Endocrinol Metab* 2007; 92: 4569-4574.
- Hernandez-Presa M, et al. Angiotensin-converting enzyme inhibition prevents arterial nuclear factor- $\kappa$ B activation, monocyte chemoattractant protein-1 expression, and macrophage infiltration in a rabbit model of early accelerated atherosclerosis. *Circulation*. 1997; 95:1532-1541.
- Hevener AL, et al. Macrophage PPAR gamma is required for normal skeletal muscle and hepatic insulin sensitivity and full antidiabetic effects of thiazolidinediones. *J Clin Invest*, 2007; 117:1658-1669.
- Hirosumi J, et al. A central role for JNK in obesity and insulin resistance. *Nature*, 2002 ; 420:333-336.
- Holcomb IN, et al. FIZZ1, a novel cysteine-rich secreted protein associated with pulmonary inflammation, defines a new gene family. *EMBO J* 2000;19: 4046-4055.
- Hotamisligil GS, et al. Adipose expression of tumor necrosis factor-alpha: direct role in obesity-linked insulin resistance. *Science*. 1993; 259: 87-91.
- Hotamisligil GS, et al. Increased adipose tissue expression of tumor necrosis factor-alpha in human obesity and insulin resistance. *J Clin Invest*. 1995; 95: 2409-2415.

- Hotamisligil GS, et al. IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF alpha- and obesity-induced insulin resistance. *Science* 1996; 271:665–668.
- Hotamisligil GS. Inflammatory pathways and insulin action. *Int J Obes Relat Metab Disord* 2003; 27: S53–S55.
- Hotamisligil GS. Mechanisms of TNF-alpha-induced insulin resistance. *Exp. Clin. Endocrinol. Diabetes*. 1999; 107: 119-25.
- Hotta K, et al. Circulating concentrations of the adipocyte protein adiponectin are decreased in parallel with reduced insulin sensitivity during the progression to type 2 diabetes in rhesus monkeys. *Diabetes* 2001; 50: 1126-1133.
- Hotta K, et al. Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 2000; 20: 1595-9.
- Howard JK, et al. Leptin protects mice from starvation-induced lymphoid atrophy and increases thymic cellularity in ob/ob mice. *J Clin Invest* 1999; 104: 1051–9.
- Hsueh, W.A. and Law, R.E. PPARgamma and atherosclerosis: effects on cell growth and movement. *Arterioscler. Thromb. Vasc. Biol.* 2001; 21: 1891–1895
- Hu E, et al. AdipoQ is a novel adipose-specific gene dysregulated in obesity. *J. Biol. Chem.* 1996; 271, 10697–10703.
- Hube F, Hauner H. The role of TNF-alpha in human adipose tissue: prevention of weight gain at the expense of insulin resistance? *Horm Metab Res* 1999, 31:626–631.
- Hulley S, et al Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. Heart and Estrogen/progestin Replacement Study (HERS) Research Group. *JAMA* 1998; 280: 605–613
- Inadera H, et al. Increase in circulating levels of monocyte chemoattractant protein-1 with aging. *J Interferon Cytokine Res* 1999; 19: 1179-1182.
- Ingelsson E, et al. Retinol-binding protein-4 is associated with endothelial dysfunction in adults with newly diagnosed type 2 diabetes mellitus. *Atherosclerosis* 2009; 204: 23–25
- Inouye KE, et al. Absence of CC chemokine ligand 2 does not limit obesity-associated infiltration of macrophages into adipose tissue. *Diabetes*, 2007; 56: 2242–2250.
- Isobe T, et al. Influence of gender, age and renal function on plasma adiponectin level: the Tanno and Sobetsu study. *Eur J Endocrinol* 2005; 153: 91-98.
- Itoh K, et al. Relationship between changes in serum leptin levels and blood pressure after weight loss. *Hypertens Res* 2002; 25: 881–886.
- Iwaki M, et al. Induction of adiponectin, a fat-derived antidiabetic and antiatherogenic factor, by nuclear receptors. *Diabetes* 2003; 52, 1655–1663.
- Iwashima Y, et al. Hypoadiponectinemia is an independent risk factor for hypertension. *Hypertension* 2004; 43: 1318-1323.

- Janke J, et al. Resistin gene expression in human adipocytes is not related to insulin resistance. *Obes Res* 2002; 10: 1-5.
- Jia SH, et al. Pre-B cell colony-enhancing factor inhibits neutrophil apoptosis in experimental inflammation and clinical sepsis *J. Clin. Invest.* 2004; 113, 1318-1327
- Jia W, et al. Association of serum retinol-binding protein 4 and visceral adiposity in Chinese subjects with and without type 2 diabetes. *J Clin Endocrinol Metab* 2007; 92: 3224-3229.
- Jiao P, and Xu H. Adipose inflammation: cause or consequence of obesity-related insulin resistance. *Diabetes, Metabolic Syndrome and Obesity* 2008;1 25–31.
- Juhan-Vague I, et al. Increased plasminogen activator inhibitor activity in non insulin dependent diabetic patients-relationship with plasma insulin. *Thromb Haemost* 1989; 61: 370-373.
- Jung HS, et al. Resistin is secreted from macrophages in atheromas and promotes atherosclerosis. *Cardiovascular Research* 2006; 69: 76 – 85.
- Kahn R, et al. The metabolic syndrome: time for a critical appraisal. Joint statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care.* 2005; 28: 2289–2304.
- Kahn SE, et al. Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature* 2006; 444: 840–846.
- Kamei N, et al. Overexpression of monocyte chemoattractant protein-1 in adipose tissues causes macrophage recruitment and insulin resistance. *J Biol Chem*, 2006; 281:26602–26614.
- Kanaya AM, et al. Serum adiponectin and coronary heart disease risk in older Black and White Americans. *J Clin Endocrinol Metab* 2006; 91: 5044-5050.
- Kanazawa A, et al. Wnt5b partially inhibits canonical Wnt/beta-catenin signaling pathway and promotes adipogenesis in 3T3–L1 preadipocytes. *Biochem Biophys Res Commun.* 2005; 330: 505–510.
- Kanda H, et al. MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J Clin Invest*, 2006; 116:1494–505.
- Kaser S, et al. Resistin messenger-RNA expression is increased by proinflammatory cytokines in vitro. *FEBS Lett* 2003; 309: 286-290.
- Kato K, et al. Improvement by the insulin-sensitizing agent, troglitazone, of abnormal fibrinolysis in type 2 diabetes mellitus. *Metabolism* 2000; 49: 662–665.
- Katsuki A, et al. Serum levels of tumor necrosis factor-alpha are increased in obese patients with noninsulin-dependent diabetes mellitus. *J Clin Endocrinol Metab* 1998, 83: 859–862.
- Katsuki A, et al. Troglitazone reduces plasma levels of tumor necrosis factor-alpha in obese patients with type 2 diabetes. *Diabetes Obes Metab* 2000, 2:189–191.
- Kawanami D, et al. Direct reciprocal effects of resistin and adiponectin on vascular endothelial cells: a new insight into adipocytokine-endothelial cell interactions. *Biochem Biophys Res Commun* 2004; 314: 415–419.

## References

- Kellerer M, et al. Circulating TNF-alpha and leptin levels in offspring of NIDDM patients do not correlate to individual insulin sensitivity. *Horm Metab Res* 1996; 28: 737-743.
- Kern PA, et al. Adipose tissue tumor necrosis factor and interleukin-6 expression in human obesity and insulin resistance. *Am J Physiol Endocrinol Metab* 2001; 280:E745-E751.
- Kiechl S, et al. Chronic infections and the risk of carotid atherosclerosis: prospective results from a large population study. *Circulation* 2001; 103:1064.
- Kim CS, et al. Circulating levels of MCP-1 and IL-8 are elevated in human obese subjects and associated with obesity-related parameters. *Int J Obes* 2006; 30: 1347-1355.
- Kim JK, et al. Mechanism of insulin resistance in A-ZIP/F-1 fatless mice. *J Biol Chem*. 2000; 275: 8456-8460.
- Kim JK, et al. Prevention of fat-induced insulin resistance by salicylate. *J Clin Invest* 2001; 108: 437-446.
- Kim KH, et al. A cysteine-rich adipose tissue-specific secretory factor inhibits adipocyte differentiation. *J Biol Chem* 2001; 276, 11252-11256.
- Kimura M, et al. T lymphopenia in obese diabetic (db/db) mice is non-selective and thymus independent. *Life Sci* 1998; 62:1243-50.
- King AB, Armstrong DU. Lipid response to pioglitazone in diabetic patients: clinical observations from a retrospective chart review. *Diabetes Technol. Ther.* 2002; 4: 145-151.
- Kintscher U, et al. T-lymphocyte Infiltration in Visceral Adipose Tissue. A Primary Event in Adipose Tissue Inflammation and the Development of Obesity-Mediated Insulin Resistance. *Arterioscler Thromb Vasc Biol*. 2008; 28:1304-1310
- Kirkham TC, et al. Endocannabinoid levels in rat limbic forebrain and hypothalamus in relation to fasting, feeding and satiation: stimulation of eating by 2-arachidonoyl glycerol. *Br J Pharmacol* 2002; 136: 550-557.
- Kistorp C, et al. Plasma adiponectin, body mass index, and mortality in patients with chronic heart failure. *Circulation* 2005; 112: 1756-1762.
- Kleemann R, et al. Apolipoprotein C-I Is Crucially Involved in Lipopolysaccharide-Induced Atherosclerosis Development in Apolipoprotein E Knockout Mice. *Circulation* 2007;116: 2173-2181.
- Klotting N, et al. Serum Retinol-Binding Protein Is More Highly Expressed in Visceral than in Subcutaneous Adipose Tissue and Is a Marker of Intra-abdominal Fat Mass. *Cell Metabolism* 2007; 6: 79-87.
- Koerner A, et al. Adipocytokines: leptin – the classical. Resistin – the controversial, adiponectin – the promising, and more to come. *Best Pract Res Clin Endocrinol Metab* 2005;19: 525-546.
- Konstantinides S, et al. Leptin-independent platelet aggregation and arterial thrombosis suggests a mechanism for atherothrombotic disease in obesity. *J Clin Invest* 2001; 108: 1533- 1540.

- Kopelman PG. Obesity as a medical problem. *Nature* 2000; 404: 635–643.
- Kopp HP, et al. Impact of weight loss on inflammatory proteins and their association with the insulin resistance syndrome in morbidly obese patients. *Arterioscler Thromb Vasc Biol* 2003; 23: 1042–1047.
- Koshiyama H, et al. Rapid communication: inhibitory effect of pioglitazone on carotid arterial wall thickness in type 2 diabetes. *J Clin Endocrinol Metab*. 2001; 86: 3452–3456.
- Kralisch S, et al. Hormonal regulation of the novel adipocytokine visfatin in 3T3-L1 adipocytes. *J Endocrinol* 2005; 185: R1–8.
- Kranzhofer R, et al. Angiotensin induces inflammatory activation of human vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol*. 1999;19:1623–1629.
- Kumada M, et al. Association of hypoadiponectinemia with coronary artery disease in men. *Arterioscler Thromb Vasc Biol* 2003; 23: 85–89.
- Kusminski CM, et al. Role of resistin in obesity, insulin resistance and Type II diabetes. *Clin. Sci. (Lond)* 2005; 109, 243–256.
- Kwon K, et al. Reciprocal association between visceral obesity and adiponectin: in healthy premenopausal women. *Int J Cardiol* 2005; 101: 385–390.
- Kyriakis JM, Avruch J. Mammalian mitogen-activated protein kinase signal transduction pathways activated by stress and inflammation. *Physiol Rev* 2001; 81: 807–869.
- La Cava A. & Matarese, G. The weight of leptin in immunity. *Nature Rev. Immunol*. 2004; 4, 371–379.
- Lacasa D, et al. Macrophage-Secreted Factors Impair Human Adipogenesis: Involvement of Proinflammatory State in Preadipocytes. *Endocrinology* 2006; 148: 868–77.
- Lam QLK, et al. Involvement of leptin signaling in the survival and maturation of bone marrow-derived dendritic cells. *Eur J Immunol* 2006; 36: 3118–3130.
- Lau DCW, et al. Adipokines: molecular links between obesity and atherosclerosis *Am J Physiol Heart Circ Physiol* 2005; 288: H2031–H2041.
- Lavie CJ, et al. Obesity and Cardiovascular Disease. Risk Factor, Paradox, and Impact of Weight Loss. *J Am Coll Cardiol*, 2009; 53:1925–32.
- Lawlor DA, et al. Plasma adiponectin levels are associated with insulin resistance, but do not predict future risk of coronary heart disease in women. *J Clin Endocrinol Metab* 2005; 90: 5677–5683.
- Lawn RM, et al. The Tangier disease gene product ABC1 controls the cellular apolipoprotein-mediated lipid removal pathway. *J Clin Invest* 1999; 104: R25–R31
- Lee GH, et al. Abnormal splicing of the leptin receptor in diabetic mice. *Nature* 1996; 379: 632–635.
- Lee J, et al. Saturated fatty acids but not unsaturated fatty acids induce the expression of COX-2 mediated through Toll like receptor 4. *J. Biol. Chem* 2001; 276:16683–16689.

- Lee JW, et al. Visceral adiposity is associated with serum retinol binding protein-4 levels in healthy women. *Obesity* 2007; 15: 2225-2232.
- Lee JY, et al. Saturated fatty acids, but not unsaturated fatty acids, induce the expression of cyclooxygenase-2 mediated through Toll-like receptor 4. *J Biol Chem*, 2001; 276:16683–16689.
- Lehr HA, et al. Immunopathogenesis of atherosclerosis: endotoxin accelerates atherosclerosis in rabbits on hypercholesterolemic diet. *Circulation* 2001; 104: 914.
- Lehrke M, et al. An inflammatory cascade leading to hyperresistinemia in humans. *PLoS Med* 2004; 1: e45.
- Lemieux I, et al. Elevated C-reactive protein: another component of the atherothrombotic profile of abdominal obesity. *Arterioscler Thromb Vasc Biol* 2001; 21: 961-7.
- Li AC, et al. 2000. Peroxisome proliferator-activated receptor  $\gamma$  ligands inhibit development of atherosclerosis in LDL receptor-deficient mice. *J. Clin. Invest.* 106:523–531.
- Li H, et al. An atherogenic diet rapidly induces VCAM-1, a cytokine regulatable mononuclear leukocyte adhesion molecule, in rabbit endothelium. *Arterioscler Thromb.* 1993; 13: 197–204.
- Li H, Sun B. Toll-like receptor 4 in atherosclerosis *J Cell Mol Med.* 2007;11: 88-95.
- Li S, et al. Enhanced Proatherogenic Responses in Macrophages and Vascular Smooth Muscle Cells Derived From Diabetic db/db Mice. *Diabetes* 2006; 55: 2611–2619
- Libby P, et al. Inflammation and Atherosclerosis. *Circulation.* 2002;105:1135-1143.
- Libby P, et al. Macrophages and atherosclerotic plaque stability. *Curr Opin Lipidol.* 1996; 7: 330–335.
- Libby P. Current concepts of the pathogenesis of the acute coronary syndromes. *Circulation.* 2001;104: 365–372.
- Libby P. Inflammation in atherosclerosis. *Nature* 2002; 420: 868–874.
- Lim S, et al. Insulin-Sensitizing Effects of Exercise on Adiponectin and Retinol-Binding Protein-4 Concentrations in Young and Middle-Aged Women. *J. Clin. Endocrinol. Metab.* 2008; 93: 2263-2268.
- Lin Y, et al. The lipopolysaccharide-activated toll-like receptor (TLR)-4 induces synthesis of the closely related receptor TLR-2 in adipocytes. *J Biol Chem.* 2000; 275: 24255–24263.
- Lindsay RS, et al. Adiponectin and coronary heart disease: the Strong Heart Study. *Arterioscler Thromb Vasc Biol* 2005; 25: e15-6.
- Lindsay RS, et al. Adiponectin and development of type 2 diabetes in the Pima Indian population. *Lancet* 2002; 360: 57-58.
- Lion CJ, et al. Mini Review; Adiposity, inflammation, and atherogenesis. *Endocrinology* 2003; 144: 2195-2200.
- Loffreda S, et al. Leptin regulates proinflammatory immune responses. *FASEB J* 1998;12: 57–65,



- Longo KA, et al. Wnt10b inhibits development of white and brown adipose tissues. *J Biol Chem*. 2004; 279 :35503–35509.
- Lord GM, et al. Leptin modulates the T-cell immune response and reverses starvation- induced immunosuppression. *Nature* 1998; 394: 897–901.
- Loskutoff DJ, Samad F. The adipocyte and hemostatic balance in obesity: studies of PAI-1. *Arterioscler Thromb Vasc Biol* 1998; 18: 1-6.
- Lu SC, et al. Lipopolysaccharide increases resistin gene expression in vivo and in vitro. *FEBS Letters*. 2002; 530, 158-162
- Lumeng CN, et al. Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *J Clin Invest*, 2007;117: 175–184.
- Lundgren M, et al. Fat cell enlargement is an independent marker of insulin resistance and 'hyperleptinaemia'. *Diabetologia*. 2007; 50: 625–633.
- Luo et al. Macrophage Adiponectin Expression Improves Insulin Sensitivity and Protects Against Inflammation and Atherosclerosis. *Diabetes* 2010; 59:791-799.
- Mach F, et al. Differential expression of three T lymphocyte-activating CXC chemokines by human atheroma-associated cells. *J Clin Invest*. 1999; 104: 1041–1050.
- MacMahon S, et al. Blood pressure, stroke, and coronary heart disease. Part 1. Prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet* 1990; 335: 765–774
- Madonna R, et al. Insulin enhances vascular cell adhesion molecule-1 expression in human cultured endothelial cells through a pro-atherogenic pathway mediated by p38 mitogen-activated protein-kinase. *Diabetologia* 2004; 47:532–536.
- Maeda K., et al. cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1 (AdiPose Most abundant Gene transcript 1). *Biochem. Biophys. Res. Commun*. 1996; 221, 286–289.
- Maeda N, et al. Diet-induced insulin resistance in mice lacking adiponectin/ACRP30. *Nature Med*. 2002; 8: 731–737.
- Maeda N, et al. PPARgamma ligands increase expression and plasma concentrations of adiponectin, an adipose-derived protein. *Diabetes* 2001; 50: 2094–2099.
- Maeda N. et al. Diet induced insulin resistance in mice lacking adiponectin/ACRP30, *Nat Med*. 2002; 8: 731-737.
- Mancuso P, et al. Leptin augments alveolar macrophage leukotriene synthesis by increasing phospholipase activity and enhancing group IVC iPLA2 (cPLA2gamma) protein expression. *Am J Physiol Lung Cell Mol Physiol* 2004; 287: 497–502.
- Mannino, D. M. et al. Boys with high body masses have an increased risk of developing asthma: findings from the National Longitudinal Survey of Youth (NLSY). *Int. J. Obesity (Lond)*. 2006; 30, 6–13.

- Marcil M, et al. Mutations in the ABC1 gene in familial HDL deficiency with defective cholesterol efflux. *Lancet* 1999; 354: 1341–1346
- Marx N, et al. Effect of rosiglitazone treatment on soluble CD40L in patients with type 2 diabetes and coronary artery disease. *Circulation* 2003, 107: 1954–1957.
- Matarese G, et al. Leptin in immunology. *J Immunol* 2005;174:3137–3142.
- Matias I et al. Regulation, function, and dysregulation of endocannabinoids in models of adipose and beta-pancreatic cells and in obesity and hyperglycemia. *J Clin Endocrinol Metab* 2006; 91: 3171–3180.
- Matsushita K, et al. Comparison of circulating adiponectin and proinflammatory markers regarding their association with metabolic syndrome in Japanese men. *Arterioscler Thromb Vasc Biol* 2006; 26: 871-876.
- Matsuzawa Y, et al. Adiponectin and metabolic syndrome. *Arterioscler Thromb Vasc Biol.* 2004; 24: 29–33.
- Matsuzawa Y. Therapy insight: adipocytokines in metabolic syndrome and related cardiovascular disease. *Nat Clin Pract Cardiovasc Med* 2006;3:35–42.
- McGill HC Jr, et al. Obesity accelerates the progression of coronary atherosclerosis in young men. *Circulation* 2002;105:2712–2718.
- McGill JB, et al. Factors responsible for impaired fibrinolysis in obese subjects and NIDDM patients. *Diabetes* 1994, 43:104–109.
- McGill JB, et al. Factors responsible for impaired fibrinolysis in obese subjects and NIDDM patients. *Diabetes* 1994; 43: 104-109.
- Medzhitov R, et al. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature*, 1997; 388: 394–397.
- Medzhitov R, Janeway CJ. The Toll like receptor family and microbial recognition, *Trends Microbiol.* 2000; 8: 452-456
- Megawa H, et al. Short term low dosage pioglitazone treatment improves vascular dysfunction in patients with type 2 diabetes. *Endocr J* 2007; 54: 613-618.
- Mertens I, et al. Visceral fat is a determinant of PAI-1 activity in diabetic and non-diabetic overweight and obese women. *Horm Metab Res* 2001; 33: 602-607.
- Michelsen KS, et al. TLR Signaling: An Emerging Bridge from Innate Immunity to Atherogenesis. *J. Immunol.*2004; 173; 5901-5907
- Miner JL. The adipocyte as an endocrine cell. *J Anim Sci* 2004;82: 935–941.
- Mittra S, et al. Increase in weight induced by muraglitazar, a dual PPAR $\alpha$ /gamma agonist, in db/db mice: adipogenesis/or oedema? *Br J Pharmacol* 2007; 150: 480–487.
- Mohamed-Ali V, et al. Subcutaneous adipose tissue releases interleukin-6, but not tumor necrosis factor- $\alpha$ , in vivo. *J Clin Endocrinol Metab* 1997; 82: 4196-4200.

- Mohanty P, et al. Evidence for a potent antiinflammatory effect of rosiglitazone. *J. Clin. Endocrinol.Metab.* 2004; 89: 2728–2735.
- Mohapatra J, et al. Involvement of adipokines in rimonabant-mediated insulin sensitivity in ob/ob mice. *J Pharm pharmacol* 2009; 61: 1493-1498.
- Moldes M, et al. Peroxisome-proliferator-activated receptor gamma suppresses Wnt/betacatenin signalling during adipogenesis. *Biochem J.* 2003;376:607–613.
- Moller DE. Potential role of TNF- $\alpha$  in the pathogenesis of insulin resistance and type 2 diabetes. *Trends Endocrinol Metab* 2000, 11: 212–217
- Montague CT, et al. Depot-related gene expression in human subcutaneous and omental adipocytes. *Diabetes* 1998; 47:1384-1391
- Moore GBT, et al. Differential regulation of adipocytokine mRNA by rosiglitazone in db/db mice. *Biochem Biophys Res Commun* 2001; 286: 735–741.
- Morimoto Y, et al. KB-R7785, a novel matrix metalloproteinase inhibitor, exerts its antidiabetic effect by inhibiting tumor necrosis factor- $\alpha$  production. *Life Sci*, 1997; 61:795–803.
- Morise T, et al. Increased plasma levels of immunoreactive endothelin and von Willebrand factor in NIDDM patients. *Diabetes Care* 1995, 18:87–89.
- Mortensen RF. C-reactive protein, inflammation, and innate immunity. *Immunol Res* 2001; 24: 163-76.
- Moschen AR, et al. Visfatin, an adipocytokine with proinflammatory and immunomodulating properties. *J Immunol* 2007; 178: 1748–1758.
- Motoshima H, et al. Adiponectin suppresses proliferation and superoxide generation and enhances eNOS activity in endothelial cells treated with oxidized LDL. *Biochem Biophys Res Commun.* 2004; 315: 264 –71.
- Mussa FF, et al. Chlamydia pneumoniae and vascular disease: an update. *J Vasc Surg.* 2006; 43:1301–1307.
- Nabipour I, et al. The association of metabolic syndrome and Chlamydia pneumoniae, Helicobacter pylori, cytomegalovirus, and herpes simplex virus type 1: the Persian Gulf Healthy Heart Study. *Cardiovasc Diabetol.* 2006; 5: 25.
- Nagaev I, Smith U: Insulin resistance and type 2 diabetes are not related to resistin expression in human fat cells or skeletal muscle. *Biochem Biophys Res Commun* 2001, 285:561–564.
- Nagel T, et al. Shear stress selectively upregulates intercellular adhesion molecule-1 expression in cultured human vascular endothelial cells. *J Clin Invest.* 1994; 94: 885–891.
- Nesto RW. Beyond low-density lipoprotein: addressing the atherogenic lipid triad in type 2 diabetes mellitus and the metabolic syndrome. *Am J Cardiovasc Drugs* 2005; 5:379–387.
- Neumeier M, et al. Different effects of adiponectin isoforms in human monocytic cells. *J. Leukocyte Biol.* 2006; 79, 803–808.

- Ni M, et al. Atherosclerotic plaque disruption induced by stress and lipopolysaccharide in apolipoprotein E knockout mice. *Am J Physiol Heart Circ Physiol* 2009; 296: H1598-H1606
- Nishigaki I, et al. Lipid peroxide levels of serum lipoprotein fractions of diabetic patients. *Biochem Med* 1981; 25: 373–378.
- Nissen SE, et al. Effect of rimonabant on progression of atherosclerosis in patients with abdominal obesity and coronary artery disease: the STRADIVARIUS randomized controlled trial. *JAMA*. 2008; 299: 1547-1560.
- Niu J, Kolattukudy PE. Role of MCP-1 in cardiovascular disease: Molecular mechanism and clinical implications. *Clin Sci (Lond)* 2009; 117: 95-109.
- Nogueiras R, et al. Peripheral, but not central, CB1 antagonism provides food intake independent metabolic benefits in diet-induced obese rats. *Diabetes* 2008; 57: 2977-91.
- Nolan JJ, et al. Improvement in glucose tolerance and insulin resistance in obese subjects treated with troglitazone. *N Engl J Med* 1994; 331: 1188–1193.
- Nomura S, et al. Significance of chemokines and activated platelets in patients with diabetes. *Clin Exp Immunol* 2000; 121: 437-443.
- O'st A, et al. Retinol-binding protein-4 attenuates insulin-induced phosphorylation of IRS1 and ERK1/2 in primary human adipocytes. *FASEB J*. 2007; 21: 3696 –3704.
- O'Connor JC, et al. IL-1 -mediated innate immunity is amplified in the db/db mouse model of type 2 diabetes. *J Immunol* 2005; 174: 4991– 4997
- O'Rourke L, et al. Glucosedependent regulation of cholesterol ester metabolism in macrophages by insulin and leptin. *J Biol Chem* 2002; 277: 42557–42562.
- O'Rourke RW, et al. Alterations in peripheral blood lymphocyte cytokine expression in obesity, *Clinical and Experimental Immunology*, 2006; 146: 39–46
- Oberkofler H, et al. Uncoupling protein gene: quantification of expression levels in adipose tissues of obese and non-obese humans. *J Lipid Res*. 1997;38: 2125–2133.
- Odegaard JI, et al. Macrophage specific PPARgamma controls alternative activation and improves insulin resistance. *Nature*, 2007; 447:1116–1120.
- Okamoto Y, et al. Adiponectin reduces atherosclerosis in apolipoprotein E-deficient mice. *Circulation* 2002;106: 2767–2670.
- Ong KK, et al. Sex-discordant associations with adiponectin levels and lipid profiles in children. *Diabetes* 2006; 55: 1337-1341.
- Oram JF. Tangier disease and ABCA1. *Biochim Biophys Acta* 2000;1529: 321–330.
- Osei-Hyiaman D, et al. Endocannabinoid activation at hepatic CB1 receptors stimulates fatty acid synthesis and contributes to diet-induced obesity. *J Clin Invest* 2005; 115: 1298–1305.
- Otero M, et al. Leptin, from fat to inflammation: old questions and new insights. *FEBS Lett* 2005;579: 295–301.

- Otero M, et al. Towards a pro-inflammatory and immunomodulatory emerging role of leptin. *Rheumatology (Oxford)* 2006;45: 944–950.
- Otto TC, Lane MD. Adipose development: from stem cell to adipocyte. *Crit Rev Biochem Mol Biol*. 2005;40:229–242.
- Ouchi N, et al. Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF-kappa B signaling through a cAMP-dependent pathway. *Circulation* 2000; 102:1296–1301
- Ouchi N, et al. Association of hypoadiponectinemia with impaired vasoreactivity. *Hypertension* 2003; 42: 231-234.
- Ouchi N, et al. Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin. *Circulation* 1999;100, 2473–2476.
- Ouchi N, et al. Reciprocal association of C-reactive protein with adiponectin in blood stream and adipose tissue. *Circulation* 2003; 107: 671-674.
- Pacher P, and Ungvári Z. Pleiotropic effects of the CB2 cannabinoid receptor activation on human monocyte migration: implications for atherosclerosis and inflammatory diseases. *Am J Physiol Heart Circ Physiol*. 2008; 294: H1133–H1134.
- Packard RR and Libby P. Inflammation in atherosclerosis: from vascular biology to biomarker discovery and risk prediction. *Clin Chem* 2008; 54: 24–38.
- Pagano C, et al. Reduced Plasma Visfatin/Pre-B Cell Colony-Enhancing Factor in Obesity Is Not Related to Insulin Resistance in Humans. *J. Clin. Endocrinol. Metab*. 2006; 91: 3165–3170.
- Pagano C, et al. The endogenous cannabinoid system stimulates glucose uptake in human fat cells via phosphatidylinositol 3-kinase and calcium-dependent mechanisms. *J. Clin. Endocrinol. Metab*. 2007; 92: 4810-4819.
- Pajvani UB, et al. Structure-function studies of the adipocyte-secreted hormone Acrp30/adiponectin. Implications for metabolic regulation and bioactivity. *J Biol Chem* 2003; 278: 9073-9085.
- Pajvani, UB. et al. Complex distribution, not absolute amount of adiponectin, correlates with thiazolidinedione -mediated improvement in insulin sensitivity. *J. Biol. Chem*. 2004; 279: 12152–12162
- Pang S, Le Y. Role of Resistin in inflammation and inflammation related diseases. *Cell Mol Immunol* 2006; 3: 29–34.
- Paolisso G, et al. A high concentration of fasting plasma non-esterified fatty acids is a risk factor for the development of NIDDM. *Diabetologia*. 1995; 38:1213–1217.
- Papayianni A, et al. Circulating levels of ICAM-1, VCAM-1 and MCP-1 are increased in haemodialysis patients: association with inflammation, dyslipidaemia and vascular events. *Nephrol Dial Transplant* 2002; 17: 435-441.

- Parissis JT, et al. Serum profiles of granulocyte-macrophage colony-stimulating factor and C-C chemokines in hypertensive patients with or without significant hyperlipidemia. *Am J Cardiol* 2000; 85: 777-779.
- Patel L, et al. Resistin is expressed in human macrophages and directly regulated by PPAR gamma activators. *Biochem Biophys Res Commun* 2003; 300: 472-476.
- Perseghin G, et al. Serum retinol-binding protein-4, leptin, and adiponectin concentrations are related to ectopic fat accumulation. *J Clin Endocrinol Metab* 2007; 92: 4883-4888.
- Persson J, et al. Cytokine response to lipoprotein lipid loading in human monocyte-derived macrophages. *Lipids in Health and Disease* 2006; 5:17
- Pfeiffer A, et al. Circulating tumor necrosis factor alpha is elevated in male but not in female patients with type II diabetes mellitus. *Horm Metab Res* 1997; 29: 111-114.
- Pickup JC, Crook MA. Is type II diabetes mellitus a disease of the innate immune system? *Diabetologia* 1998; 41: 1241-1248.
- Pickup JC. Inflammation and activated innate immunity in the pathogenesis of type 2 diabetes. *Diabetes Care* 2004; 27:813-23.
- Piemonti L, et al. Fasting plasma leptin, tumor necrosis factor-alpha receptor 2, and monocyte chemoattracting protein 1 concentration in a population of glucose-tolerant and glucose-intolerant women: impact on cardiovascular mortality. *Diabetes Care* 2003; 26: 2883-2889.
- Pilz S, et al. Adiponectin and mortality in patients undergoing coronary angiography. *J Clin Endocrinol Metab* 2006; 91: 4277-4286.
- Pilz S, et al. Early atherosclerosis in obese juveniles is associated with low serum levels of adiponectin. *J Clin Endocrinol Metab* 2005; 90: 4792-4796.
- Pineiro, R. et al. Adiponectin is synthesized and secreted by human and murine cardiomyocytes. *FEBS Lett.* 2005; 579, 5163-5169
- Pischon T, et al. Association of plasma resistin levels with coronary heart disease in women. *Obes Res* 2005; 13: 1764-1771.
- Pischon T, et al. Plasma adiponectin levels and risk of myocardial infarction in men. *JAMA* 2004; 291: 1730-1737.
- Pi-Sunyer FX, et al. Effect of rimonabant, a cannabinoid-1 receptor blocker, on weight and cardiometabolic risk factors in overweight or obese patients: RIO-North America: a randomized controlled trial. *JAMA* 2006; 295:761-775.
- Poltorak A, et al. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science*, 1998; 282:2085-2088.
- Popa C, et al. The role of TNF-alpha in chronic inflammatory conditions, intermediary metabolism, and cardiovascular risk. *J Lipid Res.* 2007; 48:751-762.

- Poulain Godefroy O, Froguel P. Preadipocyte response and impairment of differentiation in an inflammatory environment. *Biochemical Biophysical Research Communication* 2007; 356: 662-667.
- Pradhan AD, et al. C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA* 2001, 286:327–334.
- Primrose JN, et al. Reduction in factor VII, fibrinogen and plasminogen activator inhibitor-1 activity after surgical treatment of morbid obesity. *Thromb Haemost* 1992; 68: 396-399.
- Pyyrala K, et al. Diabetes and atherosclerosis: an epidemiologic view. *Diabetes Metab* 1987; Rev 3: 463–524.
- Qi Q, et al. Elevated retinol-binding protein 4 levels are associated with metabolic syndrome in Chinese people. *J Clin Endocrinol Metab* 2007; 92: 4827-4834.
- Qiao JH, et al. Role of macrophage colony-stimulating factor in atherosclerosis: studies of osteopetrotic mice. *Am J Pathol.* 1997;150:1687–1699.
- Quadro L, et al. Understanding the physiological role of retinol-binding protein in vitamin A metabolism using transgenic and knockout mouse models. *Mol Aspects Med* 2003; 24: 421–430.
- Rabe K, et al. Adipokines and Insulin Resistance. *Mol Med*, 2008, 14: 741-751
- Rajala MW, et al. Adipose-derived resistin and gut-derived resistin-like molecule-beta selectively impair insulin action on glucose production. *J Clin Invest* 2003; 11: 225–230.
- Rajala MW, Scherer PE. Minireview: The adipocyte—at the crossroads of energy homeostasis, inflammation, and atherosclerosis. *Endocrinology* 2003; 144: 3765–3773.
- Raso GM, et al. Leptin potentiates IFN-gamma-induced expression of nitric oxide synthase and cyclo-oxygenase-2 in murine macrophage J774A. 1. *Br J Pharmacol* 2002;137: 799–804.
- Reddy MA, et al. Key role of Src kinase in S100B-induced activation of the receptor for advanced glycation end products in vascular smooth muscle cells. *J Biol Chem* 2006; 281:13685–13693.
- Reilly MP, et al. Resistin is an inflammatory marker of atherosclerosis in humans. *Circulation* 2005; 111: 932-939.
- Reitman ML, et al. Transgenic mice lacking white fat: models for understanding human lipotrophic diabetes. *Ann N Y Acad Sci.* 1999; 892:289 –296.
- Remesar X et al. Effect of oral oleoyl-estrone on adipose tissue composition in male rats. *Int J Obes* 2002; 26: 1092–1102.
- Reyes MR, Lazalde B. Aortic preadipocyte differentiation into adipocytes induced by rosiglitazone in an in vitro model. *In Vitro Cell Dev Biol Anim.* 2007; 43:159-161.
- Ridker PM, et al. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med.* 2000; 342:836–843.

- Ridker PM, et al. C-reactive protein, the metabolic syndrome, and risk of incident cardiovascular events: an 8-year follow-up of 14719 initially healthy American women. *Circulation* 2003; 107: 391-397.
- Ridker PM. Evaluating novel cardiovascular risk factors: can we better predict heart attacks? *Ann Intern Med.* 1999;130: 933–937.
- Rimm EB, et al. Body size and fat distribution as predictors of coronary heart disease among middle-aged and older US men. *Am J Epidemiol* 1995; 141: 1117–1127.
- Robinson LJ, et al. Agonist-modulated palmitoylation of endothelial nitric oxide synthase. *J Biol Chem* 1995; 270: 995–998.
- Romano M, et al. Role of IL-6 and its soluble receptor in induction of chemokines and leukocyte recruitment. *Immunity* 1997; 6:1–20.
- Rosen ED, et al. PPAR gamma is required for the differentiation of adipose tissue in vivo and in vitro. *Mol Cell* 1999; 4: 611–617.
- Rosen ED, et al. Transcriptional regulation of adipogenesis. *Genes Dev.* 2000;14:1293-1307
- Rosen ED, MacDougald OA. Adipocyte differentiation from the inside out. *Nat Rev Mol Cell Biol.* 2006; 7: 885–896.
- Rosen ED, Spiegelman BM. Molecular regulation of adipogenesis. *Annu Rev Cell Dev Biol.* 2000;16:145–171.
- Rosen ED. The transcriptional basis of adipocyte development. *Prostaglandins Leukot Essent Fatty Acids.* 2005; 73: 31–34.
- Rosenson RS: Assessing risk across the spectrum of patients with the metabolic syndrome. *Am J Cardiol* 2005; 96:8E-10E
- Ross R: Atherosclerosis—an inflammatory disease. *N Engl J Med* 1999, 340:115–126.
- Ross SE, et al. Inhibition of adipogenesis by Wnt signaling. *Science.* 2000;289:950–953.
- Rotter V, et al. Interleukin-6 (IL-6) induces insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor- $\alpha$ , overexpressed in human fat cells from insulin-resistant subjects. *J Biol Chem* 2003; 278: 45777–45784
- Roytblat L, et al. Raised interleukin-6 levels in obese patients. *Obes Res* 2000; 8: 673–675,
- Rucker D, et al. Long term pharmacotherapy for obesity and overweight: updated metaanalysis. *Br Med J* 2007; 335:1194–1199.
- Rust S, et al. Tangier disease is caused by mutations in the gene encoding ATP-binding cassette transporter 1. *Nat Genet* 1999; 22: 352– 355
- Ryden M, et al. Mapping of early signaling events in tumor necrosis factor-alpha-mediated lipolysis in human fat cells. *J Biol Chem.* 2002; 277:1085–1091.
- Saijo S, et al. Inhibition by adiponectin of IL-8 production by human macrophages upon coculturing with late apoptotic cells. *Biochem. Biophys. Res. Commun.* 2005 ; 334, 1180–1183.



- Salmenniemi U, et al. Multiple abnormalities in glucose and energy metabolism and coordinated changes in levels of adiponectin, cytokines, and adhesion molecules in subjects with metabolic syndrome. *Circulation* 2004; 110: 3842-3848.
- Samad F, et al. Elevated expression of transforming growth factor-alpha in adipose tissue from obese mice. *Mol Med* 1997; 3:37-48
- Samad F, Loskutoff DJ. Tissue distribution and regulation of plasminogen activator inhibitor-1 in obese mice. *Mol Med* 1996; 2: 568-582
- Sandouk T, et al. Antidiabetic agent pioglitazone enhances adipocyte differentiation of 3T3-F442A cells. *Am J Physiol Cell Physiol* 1993; 264: C1600-C1608.
- Sanna V, et al. Leptin surge precedes onset of autoimmune encephalomyelitis and correlates with development of pathogenic T cell responses. *J Clin Invest* 2003;111:241-250.
- Sartipy P, Loskutoff DJ. Monocyte chemoattractant protein 1 in obesity and insulin resistance. *Proc Natl Acad Sci USA* 2003; 100: 7265-70
- Savage DB, et al. Resistin/Fizz3 expression in relation to obesity and peroxisome proliferator-activated receptor-gamma action in humans. *Diabetes* 2001; 50: 2199-2202.
- Schafer A, et al. The cannabinoid receptor-1 antagonist rimonabant inhibits platelet activation and reduces pro-inflammatory chemokines and leukocytes in Zucker rats. *Br J Pharmacol* 2008; 154: 1047-1054.
- Schafer K, et al. Leptin promotes vascular remodeling and neointimal growth in mice. *Arterioscler Thromb Vasc Biol*, 2004; 24:112-117.
- Scheen AJ, et al. Efficacy and tolerability of rimonabant in overweight or obese patients with type 2 diabetes: a randomised controlled study. *Lancet* 2006; 368:1660-1672.
- Scherer PE, et al. A novel serum protein similar to C1q, produced exclusively in adipocytes. *J. Biol. Chem.* 1995; 270, 26746-26749.
- Schiffrin EL. Beyond blood pressure: the endothelium and atherosclerosis. *Am J Hypertens* 2002;15:115S-22S.
- Schmidt AM, et al. Activation of receptor for advanced glycation end products: a mechanism for chronic vascular dysfunction in diabetic vasculopathy and atherosclerosis. *Circ Res.* 1999; 84:489-497.
- Schreyer SA, et al. Obesity and diabetes in TNF-alpha receptor- deficient mice. *J Clin Invest*, 1998; 102:402-411.
- Schwartz MW, et al. Central nervous system control of food intake. *Nature* 2000; 404: 661-671.
- Semple RK, et al. Expression of the thermogenic nuclear hormone receptor coactivator PGC-1alpha is reduced in the adipose tissue of morbidly obese subjects. *Int J Obes Relat Metab Disord.* 2004; 28: 176-179.

- Senolt L, et al. Resistin is abundantly present in rheumatoid arthritis synovial tissue, synovial fluid, and elevated serum resistin reflects disease activity. *Ann Rheum Dis* 2007; 66: 458–63.
- Shadid S, et al. Diet/Exercise versus pioglitazone: effects of insulin sensitization with decreasing or increasing fat mass on adipokines and inflammatory markers. *J Clin Endocrinol Metab* 2006; 91: 3418–3425.
- Shao D, Lazar MA: Peroxisome proliferator activated receptor  $\gamma$ , CCAAT/enhancer-binding protein  $\alpha$ , and cell cycle status regulate the commitment to adipocyte differentiation. *J Biol Chem* 1997; 272: 21473–8.
- Shetty GK, et al. Circulating adiponectin and resistin levels in relation to metabolic factors, inflammatory markers, and vascular reactivity in diabetic patients and subjects at risk for diabetes. *Diabetes Care* 2004; 27: 2450–2457.
- Shi H, et al. TLR4 links innate immunity and fatty acid-induced insulin resistance. *J Clin Invest*, 2006; 116: 3015–3025.
- Shimomura I, et al. Leptin reverses insulin resistance and diabetes mellitus in mice with congenital lipodystrophy. *Nature*. 1999; 401:73–76.
- Sierra-Honigsmann MR, et al. Biological action of leptin as an angiogenic factor. *Science* 1998; 281: 1683–1386.
- Silha JV, et al. Plasma resistin, adiponectin and leptin levels in lean and obese subjects: correlations with insulin resistance. *Eur J Endocrinol* 2003; 149: 331–335.
- Skurk T, et al. Relationship between adipocytes size and adipokine expression and secretion. *J Clin Endocrinol Metab*. 2007;92:1023–1033.
- Skurk T, Hauner H. Obesity and impaired fibrinolysis: role of adipose production of plasminogen activator inhibitor-1. *Int J Obes Relat Metab Disord* 2004; 28: 1357–64.
- Smith JD, et al. Decreased atherosclerosis in mice deficient in both macrophage colony-stimulating factor (op) and apolipoprotein E. *Proc Natl Acad Sci U S A*. 1995;92: 8264–8268.
- Solini A, et al. Retinol-binding protein-4 in women with untreated essential hypertension. *Am J Hypertens* 2009; 22: 948–949.
- Soltis EE and Cassis LA. Influence of perivascular adipose tissue on rat aortic smooth muscle responsiveness. *Clin Exp Hypertens* 1991; 13: 277–96,
- Son MH, et al. Peripherally acting CB1-receptor antagonist: the relative importance of central and peripheral CB1 receptors in adiposity control. *International Journal of Obesity* 2010; 34: 547–556
- Sopasakis VR, et al. High local concentrations and effects on differentiation implicate interleukin-6 as a paracrine regulator. *Obes Res*. 2004;12:454–460.

- Sorrentino MJ: Implications of the metabolic syndrome the new epidemic. *Am J Cardiol* 2005; 96:3E-7E
- Sowers JR Obesity as a cardiovascular risk factor. *Am J Med* 2003; 115: 37S–41S.
- Srinivasan K, et al. Reversal of glucose intolerance by pioglitazone in high fat diet-fed rats. *Methods Find Exp Clin Pharmacol* 2004; 26: 327-33.
- Staiger H, et al. Relationship of serum adiponectin and leptin concentrations with body fat distribution in humans. *Obes Res* 2003; 11: 368-372.
- Statnick MA, et al. Decreased expression of apM-1 in omental and subcutaneous adipose tissue of humans with type 2 diabetes. *Int. J. Exp. Diabetes. Res.* 2000; 1: 81– 8.
- Stefan N, et al. High circulating retinol-binding protein 4 is associated with elevated liver fat but not with total, subcutaneous, visceral, or intramyocellular fat in humans. *Diabetes Care* 2007; 30: 1173-1178.
- Stefan N, et al. Plasma adiponectin concentration is associated with skeletal muscle insulin receptor tyrosine phosphorylation, and low plasma concentration precedes a decrease in whole-body insulin sensitivity in humans. *Diabetes* 2002; 51: 1884-1888.
- Steinberg BA, Cannon CP. Cannabinoid-1 receptor blockade in cardiometabolic risk reduction: safety, tolerability, and therapeutic potential. *Am J Cardiol* 2007; 100: 27–32.
- Steinberg HO, et al. Elevated circulating free fatty acid levels impair endothelium-dependent vasodilation. *J Clin Invest* 1997;100:1230–1239.
- Stejskal D, et al. Resistin- concentrations in persons with type 2 diabetes mellitus and in individuals with acute inflammatory disease. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2003; 147: 63-69.
- Stephens JM, Pekala PH: Transcriptional repression of the GLUT4 and C/EBP genes in 3T3-L1 adipocytes by tumor necrosis factor-alpha. *J Biol Chem* 1991, 266:21839–21845.
- Steppan CM, et al. The hormone resistin links obesity to diabetes. *Nature* 2001, 409:307–312.
- Stokes KY, et al. Dietary nitrite prevents hypercholesterolemic microvascular inflammation and reverses endothelial dysfunction. *Am J Physiol Heart Circ Physiol* 2009; 296:H1281-H1288.
- Stoll LL, et al. Endotoxin, TLR4 signaling and vascular inflammation: potential therapeutic targets in cardiovascular disease. *Curr Pharm Des* 2006; 12: 4229–4245
- Strackowski M, et al. Plasma interleukin-8 concentrations are increased in obese subjects and related to fat mass and tumor necrosis factor-alpha system. *J Clin Endocrinol Metab* 2002; 87: 4602-4606.
- Strassmann G, et al. The role of interleukin-6 in lipopolysaccharide-induced weight loss, hypoglycemia and fibrinogen production, in vivo. *Cytokine* 1993, 5: 285–290.
- Strissel KJ, et al. Adipocyte death, adipose tissue remodeling, and obesity complications. *Diabetes*, 2007; 56: 2910–2918.

- Stuart P, et al. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 2003; 112: 1796–1808.
- Suganami T, et al. A Paracrine Loop Between Adipocytes and Macrophages Aggravates Inflammatory Changes Role of Free Fatty Acids and Tumor Necrosis Factor alpha. *Arterioscler Thromb Vasc* 2005; 25:2062-2068.
- Takashima N, et al. Retinol-Binding Protein 4 and Insulin Resistance. *N Engl J Med* 2006; 355:1392.
- Takeda K, Akira S. Roles of Toll-like receptors in innate immune responses. *Genes Cells* 6: 2001; 733–42
- Tang QQ, et al. Mitotic clonal expansion: a synchronous process required for adipogenesis. *Proc Natl Acad Sci U S A.* 2003; 100: 44–49.
- Tchernof A, et al. Weight loss reduces C-reactive protein levels in obese postmenopausal women. *Circulation* 2002; 105: 564-569.
- Thorne A, et al. A pilot study of long-term effects of a novel obesity treatment: omentectomy in connection with adjustable gastric banding. *International J Obesity.* 2002; 26: 193–199.
- Tian Z, et al. Impaired natural killer (NK) cell activity in leptin receptor deficient mice: leptin as a critical regulator in NK cell development and activation. *Biochem Biophys Res Commun* 2002; 298: 297–302.
- Tilg H, Moschen AR. Adipocytokines: mediators linking adipose tissue, inflammation and immunity. *Nat Rev Immunol* 2006;6:772–783.
- Tominaga M, et al. Thiazolidinediones (AD-4833 and CS-045) improve hepatic insulin resistance in streptozotocin-induced diabetic rats. *Endocr J* 1993; 40: 343–349.
- Tonelli, J. et al. Mechanisms of early insulin-sensitizing effects of thiazolidinediones in type 2 diabetes. *Diabetes* 2004; 53: 1621–1629
- Tontonoz P, et al. mPPAR gamma 2: tissuespecific regulator of an adipocyte enhancer. *Genes Dev* 1994; 8: 1224–1234.
- Tontonoz P, et al. Regulation of adipocyte gene expression and differentiation by peroxisome proliferator activated receptor gamma. *Curr Opin Genet Dev* 1995; 5: 571–576.
- Tontonoz P, et al. Stimulation of adipogenesis in fibroblasts by PPAR  $\gamma$  2, a lipid-activated transcription factor. *Cell.* 1994; 79:1147-56
- Trillou CR, et al. Anti-obesity effect of SR141716, a CB1 receptor antagonist, in diet-induced obese mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 2003; 284: R345–R353.
- Trinchieri G, Sher A. Cooperation of toll-like receptor signals in innate immune defence. *Nat Rev Immunol* 2007; 7:179–190
- Tschritter O, et al. Plasma adiponectin concentrations predict insulin sensitivity of both glucose and lipid metabolism. *Diabetes* 2003; 52: 239-243.

- Tsukumo DM, et al. Loss-of function mutation in Toll-like receptor 4 prevents diet-induced obesity and insulin resistance. *Diabetes*, 2007; 56:1986–1998.
- Tsutsumi C, et al. Retinoids and retinoid-binding protein expression in rat adipocytes. *J Biol Chem* 1992; 267:1805-1810.
- Tuomainen AM, et al. *Aggregatibacter actinomycetemcomitans* induces MMP-9 expression and proatherogenic lipoprotein profile in apoE-deficient mice. *Microbial Pathogenesis* 2008; 44: 111–117.
- Ueki et al. Suppressor of Cytokine Signaling 1 (SOCS-1) and SOCS-3 Cause Insulin Resistance through Inhibition of Tyrosine Phosphorylation of Insulin Receptor Substrate Proteins by Discrete Mechanisms. *Molecular and Cellular Biology* 2004; 24: 5434-5446.
- United States Department of Health and Human Services. Reducing the health consequences of smoking: 25 years of progress. Report Surgeon General 1989; DHSS CDC 89–8411
- Usui S, et al. Association between serum retinol-binding protein 4 and small dense low-density lipoprotein cholesterol levels in young adult women. *Clin Chim Acta* 2009; 399: 45–48.
- Uysal KT, et al. Functional analysis of tumor necrosis factor (TNF) receptors in TNF-alpha-mediated insulin resistance in genetic obesity. *Endocrinology*, 1998; 139: 4832–4838.
- Uysal KT, et al. Protection from obesity-induced insulin resistance in mice lacking TNF-alpha function. *Nature*, 1997; 389: 610–614.
- Vaisse C, et al. Leptin activation of Stat3 in the hypothalamus of wild-type and ob/ob mice but not db/db mice. *Nat Genet* 1999;14: 95–99,
- Van Gaal et al. Mechanisms linking obesity with cardiovascular disease. *Nature* 2006; 444: 875-880.
- Van Gaal LF, et al. Effects of the cannabinoid-1 receptor blocker rimonabant on weight reduction and cardiovascular risk factors in overweight patients: 1-year experience from the RIO-Europe study. *Lancet* 2005; 365:1389–1397.
- Van Gall L et al. Efficacy and safety of rimonabant for improvement of multiple cardiometabolic risk factors in overweight/ obese patients. *Diabetes Care* 2008; 31: S229–S240.
- Van Snick J. Interleukin-6: an overview. *Annu Rev Immunol* 1990; 8: 253–78.
- Ventre J, et al. Targeted disruption of the tumor necrosis factor-alpha gene: metabolic consequences in obese and nonobese mice. *Diabetes* 1997, 46:1526–1531.
- Verma S, et al. Resistin promotes endothelial cell activation: further evidence of adipokine-endothelial interaction. *Circulation* 2003;108: 736–740.
- Viridis A, Schiffrin EL. Vascular inflammation: a role in vascular disease in hypertension. *Curr Opin Nephrol Hypertens* 2003;12:181–7.
- von Eynatten M, et al. Atherogenic dyslipidaemia but not total- and high-molecular weight adiponectin are associated with the prognostic outcome in patients with coronary heart disease. *Eur Heart J* 2008; 29: 1307-1315.

- von Eynatten M, et al. Retinol-binding protein 4 is associated with components of the metabolic syndrome, but not with insulin resistance, in men with type 2 diabetes or coronary artery disease. *Diabetologia* 2007; 50: 1930–1937
- Vozarova B, et al. Circulating interleukin-6 in relation to adiposity, insulin action, and insulin secretion. *Obes Res* 2001; 9: 414-417.
- Wang P, Ba ZF, and Chaudry IH. Administration of tumor necrosis factor- $\alpha$  in vivo depresses endothelium-dependent relaxation. *Am J Physiol Heart Circ Physiol* 1994; 266: H2535–H2541.
- Way et al. Adipose Tissue Resistin Expression Is Severely Suppressed in Obesity and Stimulated by Peroxisome Proliferator-activated Receptor  $\gamma$  Agonists. *The J Biol Chem* 2001; 276: 25651–25653.
- Weisberg SP, et al. CCR2 modulates inflammatory and metabolic effects of high-fat feeding. *J. Clin. Invest.* 2006; 116, 115–124.
- Weisberg SP, et al. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 2003; 112: 1796-1808.
- Wellen, KE. & Hotamisligil, GS. Inflammation, stress, and diabetes. *J. Clin. Invest.* 2005; 115, 1111–1119
- Wendt T, et al. RAGE modulates vascular inflammation and atherosclerosis in a murine model of type 2 diabetes. *Atherosclerosis* 2006;185: 70 –77.
- Weyer C, et al. Enlarged subcutaneous abdominal adipocyte size, but not obesity itself, predicts type ii diabetes independent of insulin resistance. *Diabetologia*. 2000; 43:1498 –1506.
- Wiedermann CJ, et al. Association of endotoxemia with carotid atherosclerosis and cardiovascular disease: prospective results from the Bruneck Study. *J. Am. Coll. Cardiol.* 1999; 34: 1975.
- Wilkison WO, et al. Control of the adipsin gene in adipocyte differentiation. Identification of distinct nuclear factors binding to single- and double-stranded DNA. *J Biol Chem.* 1990; 265: 477-482.
- Williams L, et al. Signal transducer and activator of transcription 3 is the dominant mediator of the anti-inflammatory effects of IL-10 in human macrophages. *J Immunol* 2004;172:567–576.
- Wilson PW, et al. Clustering of metabolic factors and coronary heart disease. *Arch Intern Med* 1999; 159: 1104–1109.
- Wilson PW, et al. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998; 97: 1837–1847,
- Wilson-Fritch L, et al. Mitochondrial biogenesis and remodeling during adipogenesis and in response to the insulin sensitizer rosiglitazone. *Mol Cell Biol.* 2003; 23:1085–1094.

- Winer JC, et al. Adiponectin in childhood and adolescent obesity and its association with inflammatory markers and components of the metabolic syndrome. *J Clin Endocrinol Metab* 2006; 91: 4415-4423.
- Wolf A.M. et al. Up-regulation of the anti-inflammatory adipokine adiponectin in acute liver failure in mice. *J. Hepatol.* 2006; 44, 537–543.
- Wolf AM, et al. Adiponectin induces the anti-inflammatory cytokines IL-10 and IL-1RA in human leukocytes. *Biochem. Biophys. Res. Commun.* 2004; 323, 630–635
- Wu Z, et al. Cross-regulation of C/EBP alpha and PPAR gamma controls the transcriptional pathway of adipogenesis and insulin sensitivity. *Mol Cell* 1999;3: 151–8.
- Xu H, et al. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 2003; 112:1821–1830.
- Yamaguchi N, et al. Adiponectin inhibits Toll-like receptor family-induced signaling. *FEBS Lett.* 2005; 579, 6821–6826.
- Yamauchi T, et al. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nat Med* 2001, 7:941–946.
- Yamauchi T, et al. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nat Med* 2001; 7: 946-9.
- Yamauchi T, et al. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nat. Med.* 2001; 7: 946-949.
- Yamuchi T. et al. Globular adiponectin protected ob/ob mice from diabetes and apoE deficient mice from atherosclerosis. *J. Biol. Chem.* 2003; 278: 2461-2468.
- Yang Q, et al. Serum retinol binding protein 4 contributes to insulin resistance in obesity and type 2 diabetes, *Nature* 2005; 436: 356-362.
- Yang WS, et al. Weight reduction increases plasma levels of an adipose-derived anti-inflammatory protein, adiponectin. *J Clin Endocrinol Metab* 2001; 86: 3815-3819.
- Yang X, Smith U. Adipose tissue distribution and risk of metabolic disease: does thiazolidinedione-induced adipose tissue redistribution provide a clue to the answer? *Diabetologia.* 2007;50:1127–1139.
- Yannakoulia M, et al. Body fat mass and macronutrient intake in relation to circulating soluble leptin receptor, free leptin index, adiponectin, and resistin concentrations in healthy humans. *J Clin Endocrinol Metab* 2003; 88: 1730-1736.
- Yao-Borengasser A, et al. Retinol Binding Protein 4 Expression in Humans: Relationship to Insulin Resistance, Inflammation, and Response to Pioglitazone. *J Clin Endocrinol Metab* 2007; 92: 2590–2597.
- Ye SQ, et al. Pre-B-cell colony-enhancing factor as a potential novel biomarker in acute lung injury *Am. J. Respir. Crit. Care Med.* 2005; 171,361-370.
- Yki-Jarvinen, H. Thiazolidinediones. *N. Engl. J. Med.* 2004; 351, 1106–1118

- Yokota T, et al. Adiponectin, a new member of the family of soluble defense collagens, negatively regulates the growth of myelomonocytic progenitors and the functions of macrophages. *Blood* 2000; 96: 1723–1732.
- Yu JG, et al. The effect of thiazolidinediones on plasma adiponectin levels in normal, obese, and type 2 diabetic subjects. *Diabetes* 2002; 51:2968–2974.
- Yuan M, et al. Reversal of obesity- and diet-induced insulin resistance with salicylates or targeted disruption of Ikk beta. *Science*, 2001; 293:1673–1677.
- Yudkin JS, et al. C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arterioscler Thromb Vasc Biol* 1999, 19: 972–978.
- Yudkin JS, et al. Inflammation, obesity, stress and coronary heart disease: is interleukin- 6 the link? *Atherosclerosis* 2000; 148: 209-214.
- Zarkesh-Esfahani H, et al. High-dose leptin activates human leukocytes via receptor expression on monocytes. *J Immunol* 2001;167: 4593–4599.
- Zhang B, et al. Down-regulation of the expression of the obese gene by an antidiabetic thiazolidinedione in Zucker diabetic fatty rats and db/db mice. *J Biol Chem* 1996; 271: 9455–9459
- Zhang JL, et al. Serum resistin level in essential hypertension patients with different glucose tolerance. *Diabet Med* 2003; 20: 828-831.
- Zhang Q, et al. Antiatherogenic property of triterpenoids-enriched extract from the aerial parts of *Salvia miltiorrhiza*. *Phytother Res* 2008; 22:1040-1045.
- Ziccardi P, et al. Reduction of inflammatory cytokine concentrations and improvement of endothelial functions in obese women after weight loss over one year. *Circulation* 2002;105: 804–809
- Zykova SN, et al. diabetic type II-like db/db mice. *Diabetes* 2000; 49:1451–1458.