

REFERENCES

REFERENCES

1. Arnone R A, Satellite--derived color-temperature relationship in the Alboran sea, *Rem. Sen. Env.*, 23 (1987) 417-437.
2. Banse K., 1968, Hydrography of the Arabian Sea Shelf of India and Pakistan, and effect on demersal fisheries, *Deep-Sea-Research*, 15, 45-79.
3. Banse K., 1973, Upsloping of isotherm on the continental shelf off Goa and Bombay in June, 1967, *Jr. Marine Biological Association of India*, 14, 344-356.
4. Barnard A H, Stegmann P M & Yoder J A, Seasonal surface ocean variability in the South Atlantic Bight derived from CZCS and AVHRR imagery, *Cont Shelf Res*, 17 (1997) 1181-1206.
5. Boje R., and Tomacz M, 1978, Ecosystem analysis and distribution of boundaries in upwellinng region, *Upwelling Ecosystem* (Editor: Boje and Tomacz), Springer – Verlag, New York. pp 3-11.
6. Carder, K. L., Steward, R. G., Paul, J. H., and Vargo, G. A., (1986), Relationships between chlorophyll and ocean color constituents as

- they affect remote-sensing reflectance models, *Limnology and Oceanography*, 34, 68-81, 1989
7. Carder, K. L., Hawes, S. K., Baker, K. A., Smith, R. C., Steward, R. G., and Mitchell, B. G., (1991), Reflectance model for quantifying chlorophyll *a* in the presence of productivity degradation products, *Journal of Geophysical Research*, 96, 20599-20611.
 8. Carr, M., E., and Broad, K., 2000, Satellite, society and the Peruvian fisheries during the year 1997-1998 El Nino, In *Satellite, Oceanography and Society*, Edited by David Halpern, Publisher. Elsevier Science B.V., pp.71-191
 9. Chaturvedi, N., Narain, A., and Pandey, P. C., (1998), Remote Sensing applications in oceanography with special reference to bio-physical coupling: A review, *Indian Journal of Marine Science*, 27, 1-9.
 10. Chaturvedi, N and Narain, A., (2003), Chlorophyll distribution pattern in the Arabian Sea: seasonal and regional variability, as observed from SeaWiFS data, *International Journal of Remote Sensing*, 24 (3), 511-518.
 11. Chauhan, P., (2002), Retrieval of water constituents using Ocean colour data: IRS P4 OCM data processing. Pre-Conference Training, PORSEC-2000, Goa, India, Dec.1-4, pp.6-10.

12. Chaturvedi, N , Narain A., and Pandey P C., (1998), Remote sensing applications in oceanography with special reference to bio-physical coupling: A review, *Indian Jr. of Marine Sciences*, 27, 1-9.
13. Clarke, G. L., Ewing, G. C., and Lorenzen, C. J., (1970a), Remote measurement of ocean color as an index of biological productivity. *In the Proceedings of Sixth International Symposium on Remote Sensing in the Environment*, Ann Arbor, Oct 13-16, 1969, 2: 991-1001.
14. Cushing D.H (1978), Upper tropic levels in upwelling areas. Pp. 101-110, Springer, Verlag, New York, 303 pp.
15. Düing W., 1970, The Monsoon Regime of the currents in the Indian ocean, East-West Centre, press, University of Hawaii, Honolulu, Hawaii, 68pp.
16. Freeerg, M.H., Kerma, D., Staples, D., Wrigley, R.C. and Sullivan, D.V., 1995, Application of space based radar systems to fisheries monitoring, control and surveillance, *Proceedings Third Thematic Conference, Remote Sensing for Marine and Coastal Environment*, 1, 278-289.

17. Fiedler, P.C., Smith, G.B. and Laurs, R.M., 1985, Fisheries application of satellite data in the eastern North Pacific. *Mar. Fish. Rev.* 46 (3):1-13.
18. Fiedler, P.C., 1984b, Satellite observation of El Nino along the Pacific Coast, *Science*, 224, 1251-1254.
19. Fiedler, P.C., and Bernard, H.J., 1987, Tunna aggregation and feeding near fronts observed in satellite imagery, *Cont. Shelf Res.* 7(8), 871-881.
20. Fiedler, P.C., 1984b, Satellite observations of El Nino along the U.S. Pacific coast, *Science*, 224, 1251-1254.
21. Fiedler, P.C., 1983, Satellite remote sensing of the habitat of spawning anchovy in the Southern California Bight, *CalCOFI Rep.* 24, 202-209.
22. Gordoa, A., Maso, M. and Voges, L. 2000, Satellite and fisheries: The Namibian hake, a case study, In *Satellite, Oceanography and Society*, Edited by David Halpern, Publisher: Elsevier Science B.V., pp. 192-205.
23. Gordon, H.R. and Clark, D.K., 1980, Atmospheric effect in the

- remote sensing of the phytoplankton pigments. *Boundary Layer Meteorology*, **18**, 229-313.
24. Gordon H R & Morel A., Water color measurements - an introduction. In *Oceanography from Space* In *Oceanography from space*, edited by J.F.R. Gower, (Plenum, New York), 1980. pp. 207-212.
25. Huntley M.E. and Lopez, M.D.G., 1992. Temperature dependent production of marine copepods: global synthesis, *American Naturalist*, **140**, 201-242.
26. Herron, T and Leming, T.D., and Li, J., 1989, Satellite detected fronts and butterfly aggregation in the Northern Gulf of Mexico, *Continental Shelf Research*, **9**, 569-588.
27. JAMES J MCCARTHY AND ALLAN R. ROBINSON, BRIAN J. ROTHSCHILD, 2002, Introduction biological-physical interactions in the sea: emergent findings and new directions
28. Lasker, R., Pelaez J. and Laurs, R.M., 1981, The use of satellite infrared imagery for describing ocean processes in relation to spawning spawning of the northern anchovy (*Engraulis mordax*), *Remote sensing of environment*, **11**, 439-453.

29. Laurs, R.M., 1971, Fishery-advisory information available to tropical Pacific tuna fleet via radio facsimiles broadcast, *Comer. Fish. Rev.*, 33, 40-42.
30. Laevastu, T. and Hayes, 1981, *Fisheries Oceanography*. Fishing News Books, Norwich, England, 199 pp.
31. Laurs, R.M. and Burcks, J.T., 1985, Living marine resources applications, In Satellite remote sensing, B. Saltzman (ed), Academic press, *Advances in Geophysics*, 27, 419-452.
32. Laurs, R.M., Fiedler, P.C., and Montgomery, 1994, Albacore catch distributions relative to environmental features observed from satellite. *Deep Sea Research*, 31, 1085-1099.
33. Leming, T.D. and Stuntz, W.E., 1984, Zones of coastal hypoxia revealed by satellite scanning have implications for strategic fishing, *Nature*, 389, 136-138.
34. Liu, A., C.Y. Peng and J.D. Schumacher, 1994, Wave current interaction study in the gulf of Alaska for detection of eddies by synthetic aperture radar, *J. Geophys. Res.*, (94 (C11):16189-16200.

35. Leming, T.D., May, N., and Jones, P. (1999), A geographic information System for near real time use of remote sensing of remote sensing in fisheries management in the gulf of Mexico, NOAA National Environmental Satellite, Data, and Information Service, Final report to US department of Commerce:
<http://www.sguit2.noaa.gov/COASTWATCH/reference.htm>
36. Lasker, R., Pelaez, J. and Laurs, R.M., 1981, The use of satellite infrared imagery for describing ocean processes in relation to spawning of the northern anchovy (*Engraulis mordax*), *Rem. Sen Environ.*, 11, 439-453.
37. Liu, A., C.Y. Peng and J.D. Schumacher, 1994, Wave current interaction study in the gulf of Alaska for detection of eddies by synthetic aperture radar, *J. Geophys. Res.*, (94 (C11):16189-16200.
38. Laurs R M, Fielder P C & Montgomery D R, Albacore tuna catch distribution relative to environmental features observed from satellite *Deep Sea Res.*, 31 (1984) 1085-1099.
39. Mayers, D.G., and Hike, P.P., 1990, An application of satellite derived sea surface temperature data to the Australian fishing industry in the near real time, *Int. J. Remote Sensing*, 11, 2103-2112

40. Maul, G., A., Roffer, M. and Sousa, F.M., 1984, Remotely sensed oceanographic patterns and variability of blue fin tuna catch in the Gulf of Mexico, *Oceano. Acta.*, &, 469-479.
41. McClain P.C., Pichel, W.G. and Walton, C.C., 1985, Comparative performance of AVHRR based multi channel sea surface temperature, Jr. *Geophysical Research*, 90, 11587-11601.
42. Mohan M, Chauhan P, Mathur A & Dwivedi R M, Atmospheric correction of MOS-B data using longwavelength and PCI based approaches, IRS-P3 MOS validation experiment: ocean applications *SAC Report No. SAC/RESA/MWRD/IRSP3/SN/02/98*, (Space Applications Centre, Ahmedabad) 1998, pp. 14-22.
43. Montgomery, D.R., 1981, Commercial applications of satellite oceanography, *Oceanus*, 24, 56-65.
44. Morel, A., (1988), Optical modeling of the upper ocean in relation to its biogenous matter content (Case I waters), *Journal of Geophysical Research*, 93, 10749 –10768.
45. Morel, A., (1991), Light and Marine photosynthesis: A spectral model with geochemical and climatological implications *Progress in Oceanography*, 263-306.

46. Morel, A , and Prieur, L , (1977), Analysis of variations in Ocean Colour, *Limnology and Oceanography*, 22, 709-722.
47. Morel, A. and Gentili, B., (1991), Diffuse reflectance of oceanic waters: its dependence on sun angle as influenced by the molecular scattering contribution, *Applied Optics*, 30, 4427-4438.
48. Mueller J L & La Violate P E, Colour and Temperature signatures of ocean fronts observed with Nimbus 7 CZCS *In Oceanography from space*, Edited by J.F.R. Gower, (Plenum, New York) 1980, pp. 295-302.
49. McCARTHY, J. J , ROBINSON A. R. and ROTHCHILD, B J., 2002, Introduction biological–physical interactions in the sea: emergent findings and new directions, *The Sea*, Volume 12, (editors: Allan R. Robinson, James J. McCarthy, and Brian J. Rothschild), ISBN 0-471-18901-4, John Wiley & Sons, Inc., New York, pp.1-18.
- Biophy Prestidge, M.C., R.P. Harris and A.H. Taylor (1995) A modelling investigation of copepod egg production in the Irish Sea. *ICES J. mar. Sci.*, 52, 693-70.
50. Narain, A., Dwivedi, R.M., Kumari, B , Chaturvedi, N., Solanki, H.U., Mankodi, P.C., 1991, Relationship between sea surface temperature (SST) and fish catch data: A feasibility Study, In *Proc.*

- Nat. Workshop on Fish. Res. Data and the Fishing Indu.*,
Visakhapatnam, 149-261.
51. Njoku, E.G., Barnett, T.P., Laurs, R.M. and Vasano, A.C., 1986,
Advances in satellite sea surface temperature measurement and
oceanographic applications, *Jr. Geophy. Res.*, 90, c6, 11571-
11586.
52. Narayana A, Solanki H U, Krishna B G, & Narain A, Geometric
correction and radiometric normalisation of NOAA AVHRR data for
fisheries application, *Int. J. Rem Sen* 16, (1995) 765-771.
53. Nath A., Rao M V, Rao T & Daniel N E, Retrieval of sea surface
temperature from NOAA AVHRR data – some results,
Mahasagar, Bull National Institute of Oceanography, 22, (1989) 83-
86
54. O'Reilly, J. E., Maritonena, S, Mitchell, B. G., Siegal, D. A., Carder,
K L, Graver, S. A., Kahru, M. and McClain, C. R., 1998, Ocean
color chlorophyll algorithms for SeaWiFs. *Journal of Geophysics*,
103, 24937-24963
55. Petit, M., Stretta, J.M., Farrugio, H. and Wadsworth, A., 1992,
Synthetic aperture radar imaging of sea surface life and fishing
activities. *IEEE, Transactions on Geoscience and Remote Sensing*,

30, 1085-1089.

56. Podestra, G.P., Browder, J.A., and Hoey, J.J., 1993, Exploring the association between swordfish catch rates and thermal fronts on US longline grounds in the Western North Atlantic. *Conti. Shelf. Res* , 13, 253-277.

57. Podesta G. P., Broder, J.A. and Hoey, J.J., 1993, Exploring the association between swordfish catch rates and thermal fronts on US long line grounds in the Western North Atlantic, *Cont. Shelf, Res.*, 13, 253-277.

58. Plant R. E. and Stone N.D., (1991), *Knowledge systems in agriculture*, McGraw Hill, New York.

59. Prestidge, M.C., R.P. Harris and A.H. Taylor (1995) A modelling investigation of copepod egg production in the Irish Sea. *ICES J. Mar. Sci.*, **52**, 693-703.

60. Qasim, 1982, Oceanography of the northern Arabian Sea, *Deep Sea Research*, 29A, 1041-1068.

61. Rao, T.S.S., 1973, Zooplankton studies in the Indian Ocean, In *The biology of Indian Ocean*, (Editor: Bernt Zeitzschel, Chapman & Hall

Limited, London) pp. 243-256.

62. Range J.A., 1984, Egg production of marine planktonic copepod *Calanus pacificus*: Laboratory observations, Jr. of experimental marine biology and ecology, 74, 53-66
63. Saitoh, S., Kosaka, S., and Lisaka, J , 1986, Satellite infrared observation of Kuroshio warm core ring and their application to study of Pacific saury migration. *Deep Sea Research*, 33, 1601-1615.
64. Schumacher, J.D., W.E. Barber, B. Holt and A. Liu 1991, Satellite observations of meso scale features in the lower Cook Inlet and Shelikoff Strait, Gulf of Alaska, NOAA, Technical Report, ERL 445-PMEL 40, 18pp
65. Seay, T.S., 1994, Two way data communication and position determination system using low earth orbit satellites, *Proceedings Marine Technology Society Conference, Challenges and Opportunity in the Marine Environment*, 472-478.
66. Smith R.L. (1968), Upwelling, *Ocean Mar. Bio. Ann. Rev.* 6: 11-46.
67. Smith S.L. and Whitledge T.K. , (1977), The role of zooplankton in the regeneration of nitrogen in a coastal upwelling system off

- NorthWest Africa, Deep Sea Research, 24, 49-56.
68. Solanki H U, Dwivedi R M & Nayak S R, Relationship between IRS MOS-B derived chlorophyll and NOAA AVHRR SST: A case study in the North West Arabian sea, India, in *Proceedings of 2nd International workshop on MOS-IRS and Ocean Colour*, (Institute of Space Sensor Technology, Berlin, Germany) 1998, pp. 327-339.
69. Solanki H U, Raman M, Kumari B, Dwivedi R M & Narain A., Seasonal trends in the fishery resources off Gujarat : salient observations using NOAA-AVHRR *Indian J. Mar Sci*, 27 (1998), 438-442
70. Solanki H.U. Dwivedi R.M., S.R. Nayak, V.S. Somvanshi, D.K. Gulati and S.K. Pattnayak 2003a, Fishery forecast using OCM chlorophyll concentration and AVHRR SST: Validation results Off Gujarat Coast, India, *International Jr. Of Remote Sensing*, Vol 24, No. 18. 3691-3699, September 20, 2003
71. Solanki H.U., Dwivedi R.M. and Nayak S.R. 2003b, Application of remotely sensed closely coupled biological and physical process for marine fishery resources exploration, cover page in *Jr. of society of Indian Remote sensing*, Vol. 31, No., 2, 67-70.

72. Somvanshi V.S., (2001), Marine fisheries of Gujarat: resources potentials and needed technologies , Proc. Experience sharing work shop organized by fisheries commissioner, Govt. of Gujarat and Indian Institute of Management, Ahmedabad: 51-56.
73. Stretta J.M , 1991, Forecasting models for tuna fishery with aeospatial remote sensing, *International Jr. of Remote Sensing*, 12, 771-779.
74. Strong, A., E. and McClain, E.P., 1984, Improved ocean surface temperature from space -comparison wiyh drift buoys, *Bull. Am. Meteolo. Sco.*, 65(2), 138-142.
75. Stunbury, K.B. and Starr, R.M. (1999), Applications of geographic information system (GIS) to habitat assessment and marine resources management, *Oceanologica Acta*, 22, 699-703.
76. Sugimoto T. and Tameish H. (1992), Warm core rings and their role on the fishing ground formation around Japan, *Deep Sea Research*, 39, Suppl. 1, S183-S201.
77. Sugimoto T. and Tameish H. (1992), Warm core rings and their role on the fishing ground formation around Japan, *Deep Sea Research*, 39, Suppl. 1, S183-S201.

78. *The Sea*, Volume 12, edited by Allan R. Robinson, James J. McCarthy, and Brian J. Rothschild, ISBN 0-471-18901-4 _ 2002
John Wiley & Sons, Inc., New York, pp.1-18
79. Wooster W.S., Scheafer and Robinson, M.K., 1967, Atlas of the Arabian Sea for fishery oceanography, University of California, Institute of Marine Resources, Ref. IMR 67-12, 35pp Charts
80. Wyrski K., 1971, Oceanographic Atlas of the International Ocean Expedition, National Science Foundation, Washington, D.C. 531pp.
81. Yamanaka, I., Ito, S., Niwa K., Tanabe, R., Yabuta, Y., and Chikuni, S., 1998, The fisheries forecasting system in Japan for coastal pelagic fish, *FAO Fisheries Technical Paper* No., 301 72pp.
82. Yelland, M., AND P.K. Taylor, Wind stress measurements from the open ocean, *J. Phys. Oceanogr.*, 26 (4), 541-558, 1996.