

LIST OF ILLUSTRATIONS

- Map - I Map of Kawant Range.
- Graph - 1 Ombrothermic diagram for Kawant Range.
- PLATE No. 1 Degraded forest of Tectona grandis and Holarrhena antidysenterica near Fanai.
- PLATE No. 2 Biotic interferences resulting in depleted forests along the roads.
- PLATE No. 3 Mohangadh - Lagerstroemia microcarpa along with Diospyros melanoxylon and Dalbergia latifolia.
- PLATE No. 4 Natural regeneration of Teak along with other species at Mohangadh.
- PLATE No. 5 Saidivasan - Biotic interference and the resulting degraded forests of scattered trees of Tectona, Butea and Diospyros. Saccharum spontaneum is conspicuous along the stream.
- PLATE No. 6 Bunjer - Degraded forests, cultivated plains and the Toddy trees indicating human habitation.
- PLATE No. 7 Ambadungar - Mining operations and the clearing of forests. In the background, forests of Adina, Tectona and Anogeissus may be noted.

- PLATE No. 8 Ambadungar - A patch of forest cleared for cultivation showing Mangifera in the centre. On the sides are Tectona, Diospyros, Terminalia and Lagerstroemia.
- PLATE No. 9 Ambadungar - Valleys support Tectona along with Terminalia, Cassine and Adina. Depleted forests on plains in the background.
- PLATE No. 10 Ambadungar - Holarrhena antidysenterica & Morinda tomentosa with Carvia callosa, few grasses as an undergrowth.
- PLATE No. 11 Highest peak of Handevdungar with Boswellia serrata in the foreground.
- PLATE No. 12 Handevdungar - Diospyros melanoxylon, Morinda tomentosa, Acacia chundra, Tectona grandis, Dendrocalamus strictus and depleted forest in the background (a summer aspect).
- PLATE No. 13 Adina cordifolia, the dominant tree species at Handevdungar.
- PLATE No. 14 Handevdungar - Cassia fistula, Morinda tomentosa and Tectona grandis association. Carvia callosa, Helicteres isora, Dendrocalamus strictus as an undergrowth.

- PLATE No. 15 Mahudabari - On the slopes Tectona grandis, Helicteres isora, Morinda tomentosa and Dendrocalamus strictus are noted.
- PLATE No. 16 Dominance of Hardwickia binata along Narmada banks at Hampsheshwar.
- PLATE No. 17 Hardwickia binata, Bombax ceiba and Tectona grandis on the banks of Narmada.
- PLATE No. 18 River beds show Cyperus. Elevated banks support Borassus flabellifer, Mangifera indica and Madhuca indica. Fanai hill in the background.
- PLATE No. 19 Stream beds support Vitex negundo. On the escarpments Morinda, Lannea, Bombax, Woodfordia and Eriophorum are noted.
- Map - 2 Demarcation line between western elements and Indo-Malayan-Indian elements as given by vegetation types. (after Meher-Homji, 1970).
- Graph - 2 Composition of various floristic elements.
- PLATE No. 20 Fruits of Oroxylum, Rhadernmachera and Stereospermum.

- PLATE No. 21 Fruits of Holarrhena, Wrightia and Dolichandrone.
- PLATE No. 22 Fruits of Tectona and Soymida.
- PLATE No. 23 Fruits of Hymenodictyon and Sterculia.
- PLATE No. 24 Fruits of Buchanania, Dalbergia and Lagerstroemia.
- PLATE No. 25 Fruits of Garuga, Boswellia and Schrebera.
- PLATE No. 26 Fruits of Terminalia and Syconia of Ficus.