



LIST OF FIGURES

Figure No.	TITLE	Page
Chapter 1	Review of literature and Introduction	
Figure 1.1	Global distribution of phosphorus reserve	2
Figure 1.2	Proposed Energy generation model in presence of Vhb	24
Figure 1.3	Proposed role of <i>vgb</i> during oxidative stress	26
Figure 1.4	Proposed model for genetic manipulation in <i>E. asburiae</i> PSI3	30
Chapter 3	Characterization of MPS abilities of <i>E. asburiae</i> PSI3 containing Gluconate dehydrogenase operon (<i>gad</i>) from <i>Pseudomonas putida</i> KT2440	
Figure 3.1	Restriction digestion pattern of pCNK11	59
Figure 3.2	Restriction digestion pattern of pCNK14	59
Figure 3.3	MPS ability of <i>E. asburiae</i> PSI3 (pTTQGm) and <i>E. asburiae</i> PSI3 (pCNK14) in TRP agar medium with glucose as carbon source.	60
Figure 3.4	Growth and pH profile of <i>E. asburiae</i> PSI3 (pTTQGm) and <i>E. asburiae</i> PSI3 (pCNK14) in TRP medium containing 45 mM glucose.	61
Figure 3.5	Effect of <i>E. asburiae</i> PSI3 (pCNK14) on the growth of mung bean (<i>Vigna radiata</i>)	63
Chapter 4	Characterization of MPS ability of <i>E. asburiae</i> PSI3 containing periplasmic invertases	
Figure 4.1	Map of plasmids used in study	72
Figure 4.2	Cloning strategy of pCNK1	79
Figure 4.3	Restriction digestion pattern of pCNK1	80
Figure 4.4	Schematic representation of recombinant PCR strategy of <i>tac</i> *SS- <i>invB</i> .	81
Figure 4.5	Recombinant PCR of constitutive <i>tac</i> promoter with <i>malE</i> signal sequence.	81
Figure 4.6	Schematic representation of cloning strategy of pCNK2	82
Figure 4.7	Restriction digestion pattern of pCNK2.	82
Figure 4.8	Schematic representation of cloning strategy of pCNK3	83
Figure 4.9	Restriction digestion pattern of pCNK3	83
Figure 4.10	Schematic representation of cloning strategy of pCNK4.	84
Figure 4.11	Restriction digestion pattern of pCNK4.	84
Figure 4.12	Schematic representation of cloning of <i>suc2</i> gene in pTTQGm ^r .	85

Figure 4.13	Restriction digestion pattern of pCNK5	85
Figure 4.14	Growth of <i>E. coli</i> transformant pCNK3 and pCNK4 on M9 minimal medium containing sucrose as carbon source	86
Figure 4.15	Growth of <i>E. coli</i> (pCNK5) on M9 minimal medium agar plate with sucrose as c source.	86
Figure 4.16	Utilization of different sugars by <i>E. coli</i> (pCNK5)	87
Figure 4.17	Invertase enzyme detection on semi native PAGE	87
Figure 4.18	MPS ability of <i>E. asburiae</i> PSI3 VC (pTTQGm) and <i>E. asburiae</i> PSI3 (pCNK4) in TRP agar medium with sucrose as carbon source	89
Figure 4.19	Growth and pH profile of <i>E. a.</i> PSI3 (pBBR1MCS-2) and <i>E. a.</i> PSI3 (pCNK4) in TRP containing 75mM Tris and 75mM sucrose	89
Figure 4.20	MPS ability of <i>E. a.</i> PSI3 (pTTQGm ^r) and <i>E. a.</i> PSI3 (pCNK5) in TRP containing varied Tris and carbon source.	90
Figure 4.21	Growth and pH profile of <i>E. a.</i> PSI3 (pTTQGm ^r) and <i>E. a.</i> PSI3 (pCNK5) in TRP containing 75mM Tris and 50mM sucrose.	90
Chapter 5	Phosphate mineralization and solubilisation abilities of <i>E. asburiae</i> PSI3 containing <i>Aspergillus fumigatus</i> phytase (<i>phyA</i>) gene and <i>C. braakii</i> Phytase (<i>appA</i>).	
Figure 5.1	Schematic representation of pCNK9 cloning strategy	103
Figure 5.2	Restriction digestion pattern of pCNK9	104
Figure 5.3	Enzyme activity of pCNK9 at pH-8.0 and 4.0	104
Figure 5.4	Schematic representation of pCNK10 cloning strategy	106
Figure 5.5	Restriction digestion pattern of pCNK10	106
Figure 5.6	Restriction digestion pattern of pCNK13	107
Figure 5.7	Growth and pH profile of <i>E. a.</i> PSI3 (pCNK9, pCNK10 and pCNK13) in 100 mM Tris buffered pH-8.0 phytate minimal medium containing 75 mM glucose.	107
Figure 5.8	Colonization of <i>E. asburiae</i> PSI3 (pCNK13) in non sterile soil after 5 day of inoculation	109
Figure 5.9	Colonization of <i>E. asburiae</i> PSI3 (pCNK13) in non sterile soil after 10day of inoculation	109
Chapter 6	Phosphate mobilization ability of <i>E. asburiae</i> PSI3 containing extracellular <i>Uromyces viciae-fabae</i> invertase (<i>inv1</i>), <i>C. braakii</i> phytase (<i>appA</i>) and <i>Vitreoscilla</i> haemoglobin (<i>vgb</i>) genes.	
Figure 6.1	Schematic representation of cloning of <i>vgb</i> gene in pTTQ18	121
Figure 6.2	Restriction digestion pattern of pCNK12	122
Figure 6.3	Restriction digestion pattern of pCNK15	122

Figure 6.4	Restriction digestion pattern of pCNK21	123
Figure 6.5	Restriction digestion pattern of pCNK22	124
Figure 6.6	MPS ability of <i>E. asburiae</i> PSI3 (pTTQGm ^r) and <i>E. asburiae</i> PSI3 (pCNK15) in TRP agar medium with glucose as carbon source	125
Figure 6.7	Growth and pH profile of <i>E. a.</i> PSI3 (pTTQGm ^r) and <i>E. a.</i> PSI3 (pCNK15)	125
Figure 6.8	GDH activity of <i>E. a.</i> PSI3 pTTQGm ^r and <i>E. a.</i> PSI3 pCNK15	126
Figure 6.9	Phytase and acid phosphatase activity of <i>E. a.</i> PSI3 (pCNK21) and <i>E. a.</i> PSI3 (pTTQGm ^r)	127
Figure 6.10	MPS ability of <i>E. asburiae</i> PSI3 (pTTQGm ^r) and <i>E. asburiae</i> PSI3 (pCNK21) in Tris phytate agar medium with glucose as carbon source.	128
Figure 6.11	Growth and pH profile of <i>E. a.</i> PSI3 (pCNK21) and vector control in unbuffered condition	128
Figure 6.12	Growth and pH profile of <i>E. a.</i> PSI3 (pCNK21) and vector control in buffered condition	129
Figure 6.13	Growth and pH profile of <i>E. a.</i> PSI3 (pTTQGm ^r) and <i>E. a.</i> PSI3 (pCNK22) in Tris phytate minimal medium containing 75mM Tris and 50mM sucrose.	130
Figure 6.14	Growth and pH profile of <i>E. a.</i> PSI3 (pTTQGm ^r) and <i>E. a.</i> PSI3 (pCNK22) in Tris phytate minimal medium containing 100 mM Tris and mixture of sugars (each at 10 mM).	131
Figure 6.15	Ca- phytate mineralization by <i>E. asburiae</i> PSI3 (pTTQGm ^r) and <i>E. asburiae</i> PSI3 (pCNK22) in buffered phytate minimal medium containing 50 mM glucose and mixture of sugars (each at 10 mM).	131

LIST OF TABLES

Table No.	TITLE	Page
Chapter 1		
Table 1.1	myo-inositol phosphate intermediate generated during enzymatic phytate degradation	12
Table 1.2	Phytase producing microorganisms used for plant growth promotion	14
Table 1.3	Phytase overexpressing transgenic plants	16
Table 1.4	Major organic acid secreted by PSBs	17
Table 1.5	Major organic acid secreted by PS fungi	18
Table 1.6	Genetic modifications for enhanced MPS by bacteria	21
Table 1.7	Application of VHb expression in metabolic engineering	26
Chapter 2		
Table 2.1	Recommended dozes of antibiotics used in this study	32
Table 2.2	Composition of SDS-PAGE reagents	40

Chapter 3		
Table 3.1	Bacterial strains and plasmids used in this study	51
Table 3.2	Activities of GAD and GDH enzymes	61
Table 3.3	Amount of Organic acid secretion and P release by <i>E. asburiae</i> PSI3 (pTTQGm) and <i>E. asburiae</i> PSI3 (pCNK14) in TRP medium containing 45 mM glucose	62
Table 3.4	Effect of <i>E. asburiae</i> PSI3 (pCNK14) inoculation on the mung bean (<i>Vigna radiata</i>) dry shoot weight, dry shoot /root weight ratio and shoot P content after 15 days	63
Chapter 4		
Table 4.1	Bacterial strains and plasmids used in this study	69
Table 4.2	Amount of Organic acid secretion and P release by <i>E. asburiae</i> PSI3 (pTTQGm), <i>E. asburiae</i> PSI3 (pCNK4) and <i>E. asburiae</i> PSI3 (pCNK5) in TRP medium containing different amount of sucrose and glucose.	91
Chapter 5		
Table 5.1	Bacterial strains and plasmid used in this study	98
Table 5.2	Organic acid secretion and P release in Tris buffered (pH-8) phytate minimal medium by <i>E. asburiae</i> PSI 3 (pCNK9, 10 and 11)	108
Table 5.3	Phytase overexpression using different expression system	111
Chapter 6		
Table 6.1	Bacterial strains and plasmids used in this study	116
Table 6.2	pH drop and Pi release in the medium after 96 h.	126
Table 6.3	pH drop and Pi release in varying Tris buffered phytate minimal medium with varying glucose.	129