

REFERENCES

- Agrawal, S.K. [1957], Kutch Mesozoic : A study of the Jurassic of Kutch with special reference to the Jhura dome, Jour. of the Paleont. Soc. India, Lucknow, V.2, P.119-130.
- Agrawal, S.K. and Tripathi [1980], A note on the Umrapur Section, Khadir Island, Dist. Kutch [Gujarat], Jour. Paleont. Soc. India, 23 & 24, P.71-80.
- Alexander, J., Nichols, G.J. and Leigh, S. [1990], The origins of marine conglomerates in the Pindus foreland basin, Greece, Sedimentary Geology, V.66, P.243-254.
- Alpert, S.P. [1974], Systematic review of the genus skolithos, Jour. Paleont., V.48, P.661-669.
- Asgaard, U. and Bromley, R.G. [1974], Sporofossiler fraden Mellem Miocene transgression; soby-fasterholt omradet, Geol. Surv. Den. Arsskr, 1973 : 1179.
- Aubouin, J. [1959], Contribution a l'etude geologique de la Grece Septentrionale : les confins de L'Epire et de la Thessalie Ann. Geol. Pays Helleniques, 9.
- Barthel, K.W. and Barth, W. [1972], Paleologic specimens from the Devonian of Bolivia, Neues. Jahrb. Geologic, Palaontologic, Monatsh, P.573-581.
- Barwis, J.H. [1985], Tubes of the modern Polychaete Diopatra Cuprea as current velocity indicators and as analogs for skolithos-monocraterion P.225-236. In H.A. Curren [ed.] Biogenic Structures : Their use in interpreting depositional environment, SEPM, Spec. Publication, No.35.
- Berner, R.A. [1981], A new geochemical classification of sedimentary environments, Jour. Sed. Petrol., V.51, P.359-365.

- Bhalla, S.N. and Abbas, Abbas [1980], The age and paleogeographical significance of Jurassic Foraminifera from Kutch, India, P.507-544.
- Biswas, S.K. [1971], Note on the Geology of Kutch, Quart. Jour. Geol. Min. Met. Soc. India, V.43, No.4, P.223.
- Biswas, S.K. [1977], Mesozoic rock-stratigraphy of Kutch, Gujarat, Quart. J. Geol. Min. Met. Soc. India, V.49, No.3 & 4, P.1-31.
- Biswas, S.K. [1981], Basin framework, paleo-environment and depositional history of the Mesozoic sediments of Kutch basin, Western India, Quart. Jour. Geol. Min. Met. Soc. India, V.53, No.1 & 2, P.56-85.
- Biswas, S.K. [1982], Rift basins in Western Margin of India with special reference to Kutch basin and its hydrocarbon prospects, Bull. A.A.P.G., V.66, No.10, P.1497-1513.
- Biswas, S.K. and Deshpande, S.V. [1968], The basement of the Mesozoic sediments of Kutch, Western India, Bull. Geol. Min. Met. Soc. India, No.40, P.1-7.
- Biswas, S.K. and Deshpande, S.V. [1970], Geological and tectonic maps of Kutch, Bull. ONGC, V.7, No.2, P.115-120.
- Borrello, A.V. [1966], Paleontografia Bonaerense Fasc S. Trazas, restos tubiformes cuerpos problematicos de la formacion La Tinta, Sierras Septentrionales-Provincia de Buenos Aires : Prov. Buenos, Aires, Gobernac, Comis. Investig., Cient, P.42.
- Boucek, B. and Elias, M. [1962], O novem Zajimavem bioglyfu I. Paleogenu Ceskoslovenskych flysovyh karpas, Geol. Prace Zpravy, V.25-26, P.145-151.
- Bromley, R.G. and Asgaard, U. [1972], Notes on Greenland trace fossils-III, a large radiating burrow system in Jurassic Micaceous Sandstones of Jameson Land, East Greenland, Greenlands Geol. Undersogelse,

- Report, V.49, P.23-30.
- Bromley, R.G. and Frey, R.W. [1974], Redescription of the trace fossil gyrolithes and taxonomic evaluation of Thalassinoides, Ophiomorpha and Spongeliomorpha, Bull. Geol. Soc. Denmark, V.23, P.311-335.
- Bull, W.B. [1972], Recognition of alluvial-fan deposits in the stratigraphic record, in Rigby, K.J. and Hamblin, W.K. [eds.] Recognition of ancient sedimentary environments, Soc. Econ. Paleont. Min. Spec. Publ. No.16, Tulsa, Okla; P.68-83.
- Burke, K. [1967], The Yallahs basin, a sedimentary basin southeast of Kingston, Jamaica, Marine Geology, V.5, P.45-60.
- Chamberlain, C.K. [1971], Morphology and ethology of trace fossils from the Quachita Mountains, southeast Oklahoma, Jour. Paleont. V.45, P.212-246.
- Chamberlain, C.K. and Clark, D.L. [1973], Trace fossils and conodonts as evidence for deep-water deposits in the Oquirrh basin of Central Utah, Jour. Paleont. V.47, No.4, P.663-682.
- Collinson, J.D. and Thompson, D.B. [1982], Sedimentary structures, George Allen & Unwin [Publishers] Ltd, London.
- Corbo, S. [1979], Vertical distribution of trace fossils in a turbidite sequence, Upper Devonian, New York State, Palaeogeogr. Palaeoclimatol. Palaeoecol. V.28, P.81-101.
- Crimes, T.P. [1973], From limestones to distal turbidities : a facies and trace fossil analysis in the Zumaya flysch [Paleocene-Eocene], North Spain, Sedimentology, V.20, P.105-131.
- Crimes, T.P. [1975], The stratigraphical significance of trace fossils, P.109-130. In R.W. Frey [ed.] the study of trace fossils, Springer-Verlag, New York.

- Crimes, T.P. and Anderson, M.M. [1985], Trace fossils from Late Precambrian-Early Cambrian Strata of south-eastern Newfoundland [Canada] : Temporal and environmental implications, Jour. Paleont. V.59, P.310-343.
- Crimes, T.P. and Grams, G.J.B. [1982], Trace fossils from the Nama Group [Precambrian-Cambrian] of southwest Africa [Namibia], Jour. Paleont. V.56, P.890-907.
- Craft, J.H. and Bridge, J.S. [1987], Shallow-marine sedimentary processes in the Late Devonian Catskill Sea, Newyork State, Geol. Soc. America Bull. V.98, P.338-355.
- Crosby, E.J. [1972], Classification of sedimentary environments in Righy, J.K. and Hamblin, W.K. eds. Recognition of ancient sedimentary environments, Soc. Econ. Paleont. Min. Spec. Pub. 16, P.4-11.
- Curren, H.A. [1985], Trace fossil assemblage of a cretaceous near-shore environment : Englishtown Formation of Delaware, USA, P.261-276. In H.A. Curren [ed.] Biogenic Structures : their use in interpreting depositional environments, SEPM Spec. Publication No.35.
- Curren, H.A. and Frey, R.W. [1977], Pleistocene trace fossils from North Carolina [USA] and their Holocene analogues. In, T.P. Crimes and J.C. Harper [eds.] trace fossils, Geological Journ. Spl. Issue 9, Seel House Press, Liverpool, P.139-162.
- Daily B., Moore, P.S. and Rust, B.R. [1980], Terrestrial - marine transition in the Cambrian rocks of Kangaroo Island, South Australia, Sedimentology, V.27, P.379-399.
- Dam Gregers [1990], Systematics of trace fossils from the marine Lower Jurassic Neill Klintor Formation, Central East Greenland, Bull. Geol. Soc. Den.38, P.119-114.

- Davies, I.C. and Walker, R.G. [1974], Transport and deposition of resedimented conglomerates : the Cap Enrage Formation, Gaspé, Quebec, J. Sediment Petrol. V.44, P.1200-1216.
- Davies, R.A. [1983], Depositional Systems : A Genetic approach to Sedimentary Geology, 1st Edition, Prentice - Hall International, P.667.
- Dott, R.H. Jr. and Bourgeois, J. [1982], Hummocky Stratification : Significance of its variable bedding sequences, Bull. Geol. Soc. America, V.93, P.663-680.
- Dutton, S.P. [1982], Pennsylvanian fan-delta and carbonate deposition, Mobeetie Field, Texas Panhandle, A.A.P.G. Bull. V.66, No.4, P.389-407.
- Ekdale, A.A. [1977], Abyssal trace fossils in worldwide deep sea drilling project cores. In, T.P. Crimes & J.C. Harper [eds.] Trace Fossils 2 : Geological Journal Special Issue 9, P.163-183.
- Ekdale, A.A. and Lewis, D.W. [1991], Trace fossils and paleoenvironmental control of ichnofacies in a Late Quaternary gravel and lower-fan-delta complex, Palaeogeogr. Palaeoclimatol. Paleoecology. V.81, P.253-279.
- Elliott, T. [1986], Deltas in H.G. Reading [Ed.] Sedimentary Environments and facies 2nd Edition, 1986, P.615, Blackwell Scientific Publications.
- Ethridge, F.G. and Wescott, W.A. [1984], Tectonic setting, recognition and hydrocarbon reservoir potential of fan-delta deposits. In sedimentology of gravels and conglomerates [Ed. by E.H. Koster and R.J. Steel] Mem. Can. Soc. Petrol. Geol. Calgary, 10, P.217-235.
- Fedonkin, M.A. [1977], Precambrian-Cambrian ichnocoenoses of the east European platform, P.183-194. In T.P. Crimes and J.C. Harper [Eds.] trace fossils, 2, Geol. J. Spec. Issue 9.
- Fillion, D. and Pickerill, R.K. [1984], On arthraria antiquata billings, 1872 and its relationship to Diplocraterion Torell, 1870 and Bifungites

- Desio, 1940, J. Palcontol. V.58, P.683-696.
- Fraser, G. and Suttner L.J. [1986], Alluvial fans and fan deltas : Boston, Internat. Human Resource.
- Frey, R.W. [1970], The lebensspuren of some common marine invertebrates near Beaufort, North Cavolina, 2, anemone burrows, Jour. Paleont. V.44, P.308-311.
- Frey, R.W., 1975, The study of trace fossils, P.562, Springer-verlag, New York.
- Frey, R.W. and Howard, J.D. [1970], Comparison of Upper Cretaceous ichnofaunas from siliceous sandstone and chalk, western interior region, USA. In T.P. Crimes and J.C. Harper [Eds.] Trace fossils, Geological Journal Special Issue 3.
- Frey, R.W. [1978], Howard, J.D. and Pryor, W.A., Ophiomorpha, its morphologic, taxonomic and environmental significance, Paleogeogr. Paleoclimatol. Paleoecol. V.25, P.199-299.
- Frey, R.W. and Howard, J.D. [1985], Trace fossils from the Panther Member, star point formation [Upper Cretaceous] Coal Creek Canyon, Utah, Jour. Paleont. V.59, P.370-404.
- Frey, R.W. and S. Pemberton [1984], Trace fossil facies models in R.G. Walker [Ed.] facies models, 2nd edition, P.189-207, Geoscience Canada Reprint Series, P.189-207.
- Frey, R.W. and Seilacher, A. [1980], Uniformity in marine invertebrata ichnology, Lethaia, V.13, P.183-207.
- Fursich, F.T. [1973], A revision of the trace fossils Spongiomorpha, Ophiomorpha and Thalassinoides. Neues Jr. Geol. Palaeont. Mh, 12, 719.

- Fursich, F.T. [1974b], Trace fossils as environmental indicators in the Corallian of England and Normandy, *Lethaia*, V.8, P.151-172.
- Fursich, F.T. [1975], Trace fossils as environmental indicators in the Corallian of England and Normandy, *Lethaia*, 8 : 151-172.
- Fursich, F.T. [1974c], On Diplocraterion Torell, 1870 and the significance of morphological features in vertical, spreiten-bearing, U-shaped trace fossils, *Jour. Paleont.* V.48, P.952-962.
- Fursich, F.T. [1981], Kennedy, W.J. and Palmer, T.J. : Trace fossils at a regional discontinuity surface : The Austin-Taylor [Upper Cretaceous] contact in Central Texas, *Jour. Paleont.* V.55, P.537-551.
- Gerrard, A.J. [1988], *Rocks and Landforms*, Unwin Hyman, London, P.319.
- Ghosh, D.N. [1969b], Biostratigraphic classification of the Patcham-Chari sequence at the Jumara section, Kutch, *Proc. 56th Ind. Soci. Cong.* Pt.III, P.214.
- Glennie, K.W. and Evans, G. [1976], A reconnaissance of the Recent sediments of the Rans of Kutch, India, *Sedimentology*, V.23, P.625-647.
- Hakes, W.G. [1976], Trace fossils and depositional environments of four clastic units, Upper Pennsylvanian, megacyclothems, northeast Kansas, *Univ. Kansas. Paleont. Contrib.* 63, P.46.
- Hallam, A. and Swett, K. [1966], Trace fossils from the Lower Cambrian pipe rock of the north-west highlands : *Scottish Jour. of Geology*, V.2, P.101-106.
- Hampton, M.A. [1972], The role of subaqueous debris flow in generating turbidity currents, *J. Sedim. Petrol.* V.42, P.775-793.
- Handford, C.R. [1980], Lower permian facies of the Palo Duro basin, texas, depositional systems, shelf-margin evolution, paleogeography,

and petroleum potential : Texas Univ. Bur. Econ. Geology Rept. Inv. 102, P.31.

Hantzschel, W. [1962], Trace fossils and problematics, P.W₁-W₂₆₉. In, I.C. Teichert [Ed.], Treatise on invertebrate paleontology, pt. W. Miscellanea, supplement 1, Geological Society of America and University of Kansas Press, Lawrence.

Hantzschel, W. [1975], Trace fossils and problematics, P.1-269. In C. Teichert [Ed.], Treatise on invertebrate paleontology, part W, Miscellanea, Suppl. I. Geological Society of America and University of Kansas Press, Lawrence.

Hattin, D.E. and Frey, R.W. [1969], Facies relations of Crossopodia Sp., a trace fossil from the Upper Cretaceous of Kansas, Iowa and Oklahoma, J. Paleont. V.43, No.6, P.1435-1440.

Hayword, A.B. [1983], Coastal alluvial fans and associated marine facies in the miocene of southwest Turkey, in Collinson, J.D. and Lenoir, J. eds. modern and ancient fluvial systems : International Association of sedimentologist Special Publication 6, P.323-326.

Hedberg, H.D. [1972], International code of stratigraphic nomenclature.

Heinberg, C. [1973], The internal structure of the trace fossils Gyrochorte and Curvolithos, Lethaia, 6 : 227-238.

Heinberg, C. Tove Birkelund [1984], Trace-fossil Assemblages and basin evolution of the vardekloft formation [Middle Jurassic, Central East Greenland], Jour. of Paleont. V.58, No. 2, P.362-397.

Hertweck, G. [1970a], The animal community of a muddy environment and the development of biofacies as effected by the life cycle of the characteristic species, Geol. Jour. Special Issue, 3, P.235-242.

Higgs, R. [1990], Sedimentology and tectonic implications of Cretaceous

- fan-delta conglomerates Queen Charlotte Islands, Canada, *Sedimentology*, V.37, P.83-103.
- Hiscott, R.N. and Middleton, G.V. [1979], Depositional mechanics of thick-bedded sandstones at the base of a submarine slope, Tourella Formation [Lower Oxfordian], Quebec, Canada, In Doyle, L.J. and Pilkey, O.H. [Eds.] *Geology of continental slopes*, S.E.P.M. Spec. Publ. 27, P.307-326.
- Holmes, A. [1965], *Principles of physical geology*, Thomas Nelson, London, P.1288.
- Howard, J.D. and Frey, R.W. [1975], Regional animal-sediment characteristics of Georgia estuaries, *Senckenbergiana Marit*, 7, P.33-103.
- Howard, J.D. and Reinck, H.E. [1972], Georgia coastal region, Sapel Island, USA : *Sedimentology and biology*, N. physical and biogenic sedimentary structures of the nearshore shelf Senckenberg, Mar. 4, 47, 79.
- Howell, B.F. [1957], New Cretaceous Scolelellid and annelid from Colorado, *J. Paleont. Soc. India*, V.2, P.149-152.
- Ineson, J.R. [1989], Coarse-grained submarine fan and slope apron deposits in a Cretaceous back arc basin, Antarctica, *Sedimentology*, V.36, P.793-819.
- Jaikrishna, I.B. Singh, J.D. Howard, Syed Abbas Jafar [1983], Implications of new data on Mesozoic rocks of Kachchh, Western Indian, *Nature*, V.305, P.790-792.
- Jaitley, A.K., Singh, C.S.P. 1983, A new species of *Pronovella* Fischer [Bivalvia] from the Bathonian [Middle Jurassic] rocks of Kaladongar, Patcham Island, Kachchh, *Jour. Geol. Soc. India*, V.24[9], P.496.
- Jenkins, D.A.L. [1972], Structural development of Western Greece, *Bull.*

- Am. Assoc. Pet. Geol. V.56, P.128-149.
- Johnson, A. [1970], Physical processes in Geology, Freeman, Cooper, San Francisco, P.577.
- Karaszewski, W. [1971], Some fossil traces from the lower liassic of the Holy Cross Mountains, Central Poland, Bull. Acad. Polon. Sci. Ser. Sci de la Terre, V.19, P.101-105
- Kennedy, W.J. [1967], Burrows and surface traces from the Lower Chalk of Southern England, Bull. Brit. Mus [Nat. Hist.], Geol. 15, P.127-167.
- Kern, J.P. [1978], Paleoenvironment of new trace fossils from the Eocene Mission Valley Formation, Jour. Paleont. V.52, P.186-194.
- Kern, J.P. and Warme, J.E. [1974], Trace fossils and bathymetry of the Upper Cretaceous Point Loma Formation, San Diego, California, Geol. Soc. Am. Bull. V.85, P.893-900.
- Kitchell, J.A. and Clerk, D.L. [1979], A multivariate approach to biofacies analysis of deep-sea traces from the Central Arctic, Jour. Paleont. V.53, P.1045-1067.
- Kreisa, R.D. [1981], Storm-generated sedimentary structures in subtidal marine facies with examples from the Middle and Upper Ordovician of Southwestern Virginia, J. Sedim. Petrol. V.51, No.3, P.823-848.
- Krishnan, M.S. [1968], Geology of India and Burma Heggabothara, Madras.
- Ksiazkiewicz, M. [1970], Observations on the ichnofauna of the Polish Carpathians, in Crimes, T.P. and Harper, J.C. [eds.] Trace Fossils, Geol. Jour. Spec. Issue 3, Liverpool, Seal House Press, P.283-322.
- Lowe, D.R. [1976a], Grain flow and grain flow deposits, J. Sedim. Petrol. V.45, P.5-15.

- Lowe, D.R. [1976b], Subaqueous liquified and fluidized sediment flows and their deposits, *sedimentology*, V.23, P.285-308.
- Lowe, D.R. [1982], Sediment gravity flows : II. Depositional Models with special reference to the deposits of high density turbidity currents, *J. Sedim. Petrol.* V.52, P.279-297.
- Markello, J.R. and Read, J.F. [1981], Carbonate ramp-to-deeper shale shelf transitions of an Upper Cambrian intrashelf basin, Nolichucky Formation, Southwest Virginia Appalachians, *sedimentology*, V.28, P.573-597.
- Martino, R.L. [1989], Trace fossils from marginal marine facies of the Kanawh Formation [Middle Pannsytranian], West Virginia, *J. Paleont.* V.63[4], P.389-403.
- Martinsson, A. [1965], Aspects of a Middle Cambrian thantotope on bland, *Geol. Foren. Stockholm Forhandl*, 87, P.181-230.
- Martintsch, E.J. and Finks, R.M. [1982], Lower Devonian ichnofacies at Highland Mills, New York, and their gradual replacement across environment gradients, *Jour. Paleont.* V.56, P.1050-1078.
- McCall, P.L. and Tevesz, M.J.S. [1983], Soft-bottom succession and the fossil record. In biotic interactions in recent and fossil benthic communities. M.J.S. Tevesz and P.L. McCall [eds.], P.157-194, New York, London : Plenum Press.
- McCann and Pickerill, R.K. [1988], Flysch trace fossils from the Cretaceous Kodiak Formation of Alaska, *Jour. Paleont.* V.62, P.330-348.
- McGowen, J.H. [1970], Gum Hollow fan-delta, Nueces Bay, Texas. Texas Bureau of Economic Geology, Report of Investigations, 69, 91.P.
- McGowen, J.H. and Groat, C.G. [1971], Van Horn Sandstone, West Texas : an alluvial fan model for mineral exploration Texas. *Bur. econ. Geol.*

Rep. Invest. 72, 57.PP.

McPherson, J.G., Shanmugam, G. and Moiola, R.J. [1987], Fan-deltas and Braid deltas : Varieties of coarse-grained deltas, Bull. Geol. Soc. Am. V.99, P.331-340.

Miller, M.F. and Johnson, K.G. [1981], Spirophyton in alluvial-tidal facies of the Catskill deltaic complex : Possible biological control of ichnofossil distribution, Jour. Paleont. V.55, P.1016-1027.

Miller, M.F. and Knox, L.W. [1985], Biogenic structures and depositional environments of a Lower Pennsylvanian coal-bearing sequence, Northern Cumberland Plateau, Tennessee, USA, P.67-98. In, H.A. Curren [ed.] Biogenic structures : their use in interpreting depositional environments, SEPM special Publication No.35.

Mount, J.F. [1984], Mixing of siliciclastic and carbonate sediments in shallow shelf environments, Geology, V.12, P.432-435.

Mount, J.F. [1985], Mixed siliciclastic and carbonate sediments : a proposal first-order textural and compositional classification, Sediment. V.32, P.435-450.

Mutti, F. [1977], Distinctive thin-bedded turbidite facies and related depositional environments in the Eocene Hecho Group [South-Central Pyrenees, Spain], Sedimentology, V.24, P.107-131.

Myers, A.C. [1970], Some Paleoichnological observations on the tube of *Diopatra cuprea* [Bosc] : Polychaete Onuphidae in Crimes, T.P. and Harper, J.C. [eds.] Trace fossils : Geol. Jour. Special Issue No.3, Liverpool, Seel House Press, P.345-360.

Nemec, W., Porebski, S.J. and Steel, R.J. [1980], Texture and structure of resedimented conglomerates : examples from Ksiaz Formation [Fammenian-Tournaisian], Poland, Sedimentology, V.27, P.519-538.

- Nemec, W. and Steel, R.J. [1984], Alluvial and coastal conglomerates : their significant features and some comments on gravelly mass-flow deposits. In sedimentology of Gravels and Conglomerates [(Ed.)] by E.H. Koster and R.J. Steel], Mem. Can. Soc. Petrol. Geol. Calgary, 10, P.1-31.
- Normark, W.R. [1970], Growth patterns of Deep-sea fans A.A.P.G. Bull. V.54, No.11, P.2170-2195.
- Normark, W.R. [1974], Submarine canyons and fan valleys : factors affecting growth patterns of deep-sea fans, in Modern and ancient geosynclinal, Sedimentation : SEPM Spec. Pub. 19, P.19, P.56-68.
- Normark, W.R. [1978], Fan valleys, channels and depositional lobes on modern submarine fans : characters for recognition of sandy turbidite environments., A.A.P.G. Bull. V.62, No.6, P.912-931.
- Osgood, R.G. Jr. [1970], Trace fossils from the Cincinnati area, Paleontographica, America, V.6, P.281-244.
- Osgood, R.G. and Szmuc, E.J. [1972], The trace fossil Zoophycos as an indicator of water depth, Bull. of American Paleontology, 62, P.1-22.
- Pascoe, E.H. [1959], A manual of Geology of India and Burma, V.II, 3rd Edition, Govt. of India Publ.
- Pemberton, S.G. and Frey, R.W. [1982], Trace fossils nomenclature and the Planolites-Paleophycus dilemma, Jour. Paleont. V.56, P.843-881.
- Pemberton, S.G.; Frey, R.W. and Bromley, R.G. [1988], The ichnotaxonomy of conostichus and other plug-shaped ichnofossils, Canad. Jour. Earth. Sci. V.25, P.866-892.
- Pettijohn, F.J. [1975], Sedimentary Rocks, 3rd Edi. Harper & Row, New York, 628.P.

- Pfeiffer, H. [1965], Volkichinium Volki n.gen. n.sp. [lebensspuren] aus den Phycodes Schichten Thuringens, Geologie 16, 781.
- Pickerill, R.K.; Fillion, D. and Harland, T.L. [1984], Middle Ordovician trace fossils in carbonates of the Trenton group between Montreal and Quebec City, St. Lawrence, Lowland, Eastern Canada, Jour. Paleont. V.58, P.416-439.
- Pickerill, R.K. and Paul F. Williams [1989], Deep burrowing in the early Palaeozoic deep sea : examples from the Cambrian - Early Ordovician Meguma Group of Nova Scotia, Can. J. Earth. Sci. V.26, P.1061-1068.
- Pickering, K.; Stow, D.; Watson, M.; and Hiscott, R. [1986], Deep-water facies, processes and models : a review and classification scheme for modern and ancient sediments, Earth. Sci. Rev. V.23, P.75-174.
- Pierson, T.C. [1981], Dominant particle support mechanisms in debris flows at Mt. Thomas, New Zealand and implications for flow mobility, Sedimentology, V.28, P.49-60.
- Pivnik, D.A. [1990], Thrust-generated fan-delta deposition : Little Muddy Creek Conglomerate, SW Wyoming, Jr. Sed. Petrol. V.60, No.4, P.489-503.
- Poddar, M.C. [1959], Stratigraphy and oil possibility in Kutch in Western India, Proc. 1st Symp. Devp. Pet. Res. of Asia and far east, No.10, P.146-148.
- Price I. [1977], Deposition and derivation of clastic carbonates on a Mesozoic Continental Margin, Orthis, Greece, Sedimentology, V.24, P.529-546.
- Prucha, J.J., Graham, J.A. and Nickelson, R.P. [1965], Basement controlled deformation in rocky mounts foreland, A.A.P.G. Bull. V.49[7], P.966-992.

- Purdy, E.G. [1964], Sediments as substrates, P.238-271. In Imbrie and N.D. Newell [ed.] Approaches to paleoecology, John Wiley, New York.
- Radwanski, A. [1970], Dependence of rock-borers and burrowers on the environmental conditions within the Tortonian littoral zone of Southern Poland, P.371-390. In T.P. Crimes and J.C. Harper [eds.] Trace fossils, Geol. Jour. Spec. Issue 3.
- Radwanski, A.; H. Friis and G. Larsen [1975], The miocene hagenor-borup sequence at Lillebaelt [Denmark]; its biogenic structures and depositional environment. Bull. Geol. Soc. Denmark, 24, P.229-260.
- Rajnath [1932], A contribution to the stratigraphy of Kutch. Quart. Jour. Geol. Min. Met. Soc. India, V.4, P.161-174.
- Rao, P.V. [1964], Geology and mineral resources of India. Intern. Geol. Cong. 22nd Session, India, P.1-43.
- Rhoads, D.C.; Boesch, D.F.; Zhican, T.; Fengshan, X.; Liquiang, H. and Nilsen, K.J. [1985], Macrobenthos and sedimentary facies on the Changjiang delta platform and adjacent continental shelf. Cont. Shelf. Res. 4, P.189-213.
- Richter, R. [1954], Fahrte eines "Riesenkrebse" im Rheinischen Schiefergebirge : Matur U. Volk, V.84, P.84, P.261-269.
- Rust, B.R. [1979], Coarse alluvial deposits. In facies models [Ed. by R.G. Walker], 1, 9-22, Geoscience, Canada, Reprint Series.
- Rust, B.R. and Koster, E.H. [1984], Coarse alluvial deposits in Walker, R.G. ed. facies models [2nd edition]. Geoscience Canada Reprint Series, P.53-69.
- Seilacher, A. [1953], Studien Zur Palichnologie II, Die fossilen Ruhespuren [Cubichnia], N. Jahrb, Geol. Palaeont. Abb. 98, P.87-124, Pl.7-13.

- Seilacher, A. [1964], Biogenic sedimentary structures. In J. Imbrie and N. Newell [Ed.] Approaches to Paleoeconology, Wiley, New York, PP.296-316.
- Seilacher, A. [1967], Bathymetry of trace fossils, Mar. Geol. 5, P.413-428.
- Seilacher, A. [1977], Pattern analysis of Paleodictyon and related trace fossils. In T.P. Crimes and Y.C. Harper [Eds.], Trace fossils II Seel House Press, Liverpool, PP.289-334.
- Seilacher, A. [1978], [Use of trace fossil assemblages for recognizing depositional environments. In P.B. Bason (Ed.) Trace fossil concepts. SEPM Short Course, 5, P.185-201] for 1978.
- Shringarpure, D.M. [1985], Ichnological studies of the rocks exposed between Chitrod and Oanfara, Eastern Kutch-Gujarat State. Unpubl. Ph.D. Thesis, M.S. University, Baroda, 397.P.
- Simpson, S. [1956], On the trace-fossil chondrites. Quart. Jour. Geol. Soc. London, V.112, P.475-499.
- Singh, C.S.P.; Jaitly, A.K. and Pandey, D.K. [1982], First Report of some Bajocian-Bathonian [Middle Jurassic] Ammonoids and the age of the oldest sediments from Kuchchh, W.India, Nenst. Stratigr. V.11[1], PP.37-40.
- Small, R.J. [1978], The study of landforms. A textbook of Geomorphology. Cambridge University Press, Cambridge IInd Edition, 502.PP.
- Smith, A.B. and Crimes, T.P. [1983], Trace fossils formed by heart urchins - a study of scolicia and related traces. Lethaia, 16, P.79-92.
- Sneh, A. [1979], Late Pleistocene fan-deltas along the Dead sea. Sediment. Petrol. V.49, P.541-552.

- Spath, L.F. [1924], On the blake collection of ammonites from Kutch, India. *Paleont. Indica*, N.S.V.9, Mem.1.
- Spath, L.F. [1927-33], Revision of the Jurassic cephalopod fauna of Kutch. India Geol. surv. Mem. *Paleontologic Indica* new Sr.V.9, Mem. Pt. 1-6, 945.P.
- Stanley, D.J. and Unrug, R. [1972], Submarine channel deposits, fluxoturbidites and other indicators of slope and base of slope environments in modern and ancient marine basins. In recognition of ancient sedimentary environments [Ed. by J.K. Rigby and W.K. Hamblin] Spec. Publ. Soc. econ. Paleont. Min. Tulsa. 16, P.287-340.
- Surlyk, F. [1978], Jurassic basin evolution of East Greenland - *Nature*, V.274, P.130-133.
- Surlyk, F. [1984], Fan-delta to submarine from conglomerates of the Volgian-Valanginian Wollaston Forland Group, East Greenland. In sedimentology of gravels and conglomerates [Ed. by E.H. Koster and R.J. Steel] Mem. Can. Soc. Petrol. Geol. Calgary, 10, P.359-382.
- Swift, D.Y.P.; Figueiredo, A.G.; Freeland, G.L. and Oertel, G.F. [1983], Hummocky cross-stratification and mega ripples : A geological double standard ? *J. Sed. Petrol.* Y.53, P.1295-1318.
- Swinbanks, D.D. and Murray, J.W. [1981], Biosedimentological zonation of boundary bay tidal flats, Fraser river delta, British Columbia. *Sedimentol.* V.28, P.201-237.
- Tankard, A.J. and Barwis, J.H. [1982], Wave-dominated deltaic sedimentation in the Devonian Bokkeveld basin of South Africa. *J. Sed. Petrol.* V.52, P.959-974.
- Tewari, B.S. [1948], The geology of the Habo hills, Cutch State [Abstract]. *Proc. 35th Ind. Sci. Cong. Pt.III*, P.148.

- Thornbury, W.D. [1969], Principles of Geomorphology. John Wiley & Sons, New York, IInd Edition, 594.PP.
- Thourton, S.E. [1984], Basin model for hemipeingic sedimentation in a tectonically active continental margin : Santa Barbara Basin, California Continental Margin. In D.A.V. slow Stow and D.J.W. Piper [Ed.] fine-grained sediments, deep-water processes and facies. Spec. Publ. Geol. Soc. London, 15, P.377-395.
- Troeh, F.R. [1965], Landform equations fitted to contour maps. Am. J. Sci. 263, P.616-627.
- Valentine, J.W. [1971], Resource supply and species diversity patterns, Lethaia, V.4, P.51-61.
- Vossler, S.M. and Pemberton, S.G. [1988], Superabundant chondrites : a response to storm buried organic material ? Lethaia, 21, P.94.
- Vyalov, O.S. [1971], Redkie problematiki iz mesozoya Pamira; Kavkaza; Vyp. Vtoroy No.7, P.85-93.
- Waagen, W. [1873-75], Jurassic fauna of Kutch : the cephalopoda. India Geol. Surv. Mem. Paleont. Indica Ser.9, V.1, Pt.1-4, 247.P.
- Walker, R.G. [1967], Turbidite sedimentary structures and their relationship to proximal and distal depositional environments. J. Sediment. Petrol. V.37, P.25-43.
- Walker, R.G. [1977], Facies and facies models. General introduction. In R.G. Walker [Ed.] facies models. Geosci. Can. Repr. Ser. 1, P.1-8.
- Walker, R.G. [1975], Generalized facies models for resedimented conglomerates of turbidite association. Geol. Sco. Amer. Bull. V.86, P.737-749.

- Walker, R.G. [1978], Deep-water sandstone facies and ancient submarine fans : Models for exploration for stratigraphic traps. A.A.P.G. Bull. V.62, No.6, P.932-966.
- Walker, R.G.; Duke, W.L. and Leckie, D.A. [1983], Hummocky Stratification : Significance of its variable bedding sequences : Discussion. Bull. Geol. Soc. America V.94, P.1245-1249.
- Webby, B.D. [1970a], Brookvalichnus, a new trace fossil from the triassic of the Sydney basin, Australia. In trace fossils, T.P. Crimes & J.C. Harper [Ed.], Geol. Jour. Spec. Issue No.3, P.527-530 Seel House Press [Liverpool].
- Weiss, W. [1940], Beobachtungen an zopfplatten : Deutsch. Geol. Gesell. Zeitschr. V.92, P.333-349.
- Weiss W. [1941], Die entstehung der "Zöpfe" im schwarzen and braunen jura : Matur U. Volk, V.71, P.179-184.
- Wescott, W.A. and Ethridge, F.G. [1980], Fan-delta sedimentology and tectonic setting - Yallahs fan-delta, Southeast Jamaica. A.A.P.G. Bull. V.64, P.374-399.
- Wescott, W.A. and Ethridge, F.G. [1983], Eocene fan delta - submarine deposition in the Wagwater trough, east-central Jamaica. Sedimentology, V.30, P.235-248.
- Wynne, A.B. [1872], Memoir on the geology of Kutch. To accompany a map compiled by A.B. Wynne and F. Fedden during the seasons 1867-68 and 1868-69. Memoir. Geol. Survey India, 9, P.1
- Young, A. [1972], Slopes. Edited by K.M. Clayton. Oliver & Boyd. Edinburgh, 288.PP.
- Zernitz, E.R. [1932], Drainage Patterns and their significant. Jour. Geology, V.40, P.498-521.