

LIST OF PLATES

Plate No.		Page No.
Plate-II.1	Folded amphibolites with southernly plunging F_2 folds (Loc-Mohmad Phalasia)	39
Plate-II.2	Interfolial shearing observed in amphibolites. AF_2 fold is coaxially folded by AF_3 fold. (Loc-Barvalli village)	39
Plate-II.3	Field photograph showing more amount of quartz-feldspar material in gneissic amphibolite (Loc-Barvalli village).	42
Plate-II.4	Basaltic dyke cutting across the amphibolitic gneiss (Loc- Kirat village)	42
Plate-II.5	Quartzite band within garnetiferous gneisses showing effect of AF_4 folds. (Loc- Kirat village)	47
Plate.II.6a	Fractured and sheared quartzite ridge in contact with schist. (Loc- Koliyari village)	47
Plate.II.6b	Sheared contact of schist and quartzite . (Loc- Dhikliya village)	49

Plate-II.6c	Quartzite lens in schistose rock	49
Plate-II.7	Duplex structure in schistose rock. (Loc- 10km to the south of Mohmad Phalasia village)	52
Plate-II.8	AF ₄ crinkles in schists. (Loc- Garanwas village)	52
Plate-III.1	IRS-FCC photo of North eastern Gujarat and southern Rajasthan.	56
Plate-III.2	Af ₂ fold hinge has been coaxially refolded by AF ₃ =DF ₂ fold (Loc- 5 km to the North of Phalasia)	71
Plate-III.3	AF ₂ fold coaxially refolded by AF ₃ .	71
Plate-III.4	AF ₂ isoclinal fold whose axis is trending due NNE-SSW direction.	73
Plate-III.5	AF ₂ =DF ₁ , recumbent fold in chlorite schist of Jharol group, showing 'a' lineation across the fold axis.	73
Plate-III.6	AF ₃ (=DF ₂) upright fold in Calc-schist. (Loc-10 km to the North of Kotra)	76
Plate-III.7	AF ₄ crinkles and puckers in gneisses (Loc- Ora village)	76

Plate-III.8	AF ₄ open fold in quartzite (Loc- Kirat village)	78
Plate-III.9a	Domal structure (sheath fold) developed on account of superposition of AF ₄ anticlinal fold on AF ₃ anticlinal fold in schistose rock.(Loc- 5km North of Phalasia)	80
Plate-III.9b	Basin structure (sheath fold) developed on account of superposition of AF ₄ synclinal fold on AF ₃ synclinal fold	80
Plate-III.10a	Laterally displaced quartz veins found in quartzites.	82
Plate-III-10b	Quartzites found slightly away from the wakal shear zone, showing initially sinistral local shear movement, which later followed by greater dextral shear movement.	82
Plate-III.11a	Uneven development of en-echelon vein system in quartzite.	86
Plate-III.11b	Sigmoidal quartz veins found along the periphery of wakal shear zone, showing dextral shear sense. (Loc-5 km to the west of Manpur village)	86

Plate-III.12a	The great Kotra shear zone (GKSZ). (Loc-5km to the east of Kotra village).	89
Plate-III.12b	Deep-slip slicken-side in Kotra shear zone.	89
Plate-III.12c	Wakal shear zone (WSZ), 5 km to the west of Manpur, developed within the Delhi quartzites. The shear sense is easterly down dip of dextral movement.	90
Plate-III.13	Sinistrally rotated porphyroblast of epidote within amphibolites	92
Plate-III.14	Rotation of porphyroclast in amphibolitic gneisses, showing sinistral sense of movement. (Loc- Kirat village)	92
Plate-III.15	Rotation of porphyroclast in gneisses showing sinistral sense of movement (Loc-Ammiwara village)	93
Plate-III.16	Mylonites showing S-shaped rotated quartzo-feldspathic porphyroclasts with broken tails as well as folded tails (Loc-Kirat village)	93
Plate-III.17	The porphyroclast of quartzo feldspathic material showing sinistral sense of movement	95

	One tail of the porphyroclast is detached. All the crinkles seen are of AF ₃ generation.	
Plate-III.18	Mylonites showing varieties of quartz-feldspathic porphyroclast with as well as without tails or broken tails into boudinaged fragments, giving sinistral sense of shear (Loc-1 km east of Ora village.	95
Plate-III.19	Gneisses showing complex refolded folds of quartzo-feldspathic bands. (Loc-Kirat village)	96
Plate-III.20	Assymetrical folds in gneisses giving sinistral sense of movement. (Loc-Kirat village)	96
Plate-III.21	Intrafolial folds in amphibolites showing sinistral sense of shearing. AF ₂ fold coaxially folded by AF ₃ during dextral shear deformation. The porphyroclasts developed during AF ₂ deformation have been affected by tail detachment during dextral shear Superposition. (Loc-Barvalli village)	97
Plate-III.22	Tuilage or tiling structures found in composite gneisses.	97

Plate-III.23	Twisted and rotated AF ₂ fold hinge of folded quartzite layer embedded in Jharol schist belt exhibiting AF ₂ fold hinge down folded vertically by AF ₃ folding. (Loc-5 km North of Phalasia)	99
Plate-III.24	Mylonitised micaceous quartzites on the river bed. (Loc-west of Ora village)	99
Plate-III.25	Highly sheared, mylonitised and sliced, micaceous quartzites. (Loc-East of Ora village)	103
Plate-III.26	Mylonites showing S-C fabrics in the shear zone running through the non-garnetiferous schists. (M-20x, XN)	104
Plate-III.27a	Formation of mica-fish by micro-faulting or micro-shearing from pre-existing mica-grain. The mica-fish shows sinistral sense of movement. (M-20x, XN)	109
Plate-III.27b	Antithetically detached mica-fish in serpentinites. The shear plane has also been serpentitized. The sense of movement is sinistral. (M-20x, PPL)	109

Plate-III.28	Antithetic splitting or pull-apart structure of mica-grain. Space created is filled by quartz grains. The grain is sinistrally sheared. (M-20x, XN)	111
Plate-III.29a	Micafish in schist indicating dextral shear sense. (M-50x, XN)	111
Plate-III.29b	Micafish in quartz-schist indicating dextral sense of shear. (M-20x, XN)	113
Plate-III.30a	Here plagioclase grain is synthetically sheared showing sinistral sense of movement (M-20x, XN)	114
Plate-III.30b	Sheared and pull-apart muscovite grain synthetically, filled by calcitic material. The sense of movement is sinistral. (M-20x, XN)	114
Plate-30c	Staurolite grain is antithetically sheared and filled up by chloritic material, the altered product of sheared staurolite. The sense of movement is sinistral. (M-20x, XN)	115
Plate-III.31a	SEM microphotograph illustrating the antithetic shear structure of mica flakes	117

showing dextral shear sense.
(M = 500)

Plate-31b	SEM microphotograph illustrating the close-up of the sheared structure shown in 36a, (M = 1000)	117
Plate-III.32	SEM microphotograph illustrating a tiny garnet porphyroblast in a mylonitised garnet mica schist showing dextral shear sense. (M=3500)	119
Plate-III.33	SEM microphotograph illustrating microshear structure in a micaceous mylonitic rock. It shows a sinistral shear sense with several microcrinkles. (M = 1000)	119
Plate-34	Rotated garnet. Inclusions of quartz are perpendicular to the main foliation. The rotation of garnet porphyroblast is sinistrally observed (M- 20x, XN)	129
Plate-III.35	Clockwise (dextral movement) rotation of garnet grain pushing the muscovite grain to rotate and inject into another grain. (M-20x, XN)	129
Plate-III.36	Flattening of garnet porphyroblast in sheared garnet mica schist.	130

Plate-III.37	Stretched garnet in gneisses along the shear plane developed at low angle oblique to the main foliation (M-20x, XN)	131
Plate-IV.1	Photomicrograph of serpen- tinite with chromite. (M = 20x, XN)	136
Plate-IV.2	Photomicrograph showing garnet, epidote and hornblende in amphibolites (M=20x, XN)	136
Plate-IV.3	Photomicrograph showing alternate rich layers of plagioclase and hornblende in amphibolites (M=20x, XN)	137
Plate-IV.4	Photomicrograph showing granoblastic structure in amphibolite (M=20x, PPL)	137
Plate-IV.5	Photomicrograph showing growth of micas along axial plane of AF_3 (M= 20x, XN)	141
Plate-IV.6	Photomicrograph of chlorite schist. (M-20x, PPL)	141
Plate-IV.7	Photomicrograph of sillimanite needles in muscovite (M-20x,XN)	144
Plate-IV.8	Photograph of AF_2 fold in schistose rock.	144
Plate-IV.9	Photomicrograph of the garnet	146

in the core portion of AF_2
 fold of plate-IV.8. Micas
 are crystallised along the
 axial plane of the fold.
 (M = 20x, XN)

Plate-IV.10	Photomicrograph of elongated garnet in schists. (M=20x, XN)	146
Plate-IV.11	Photomicrograph showing synkinematic growth of garnet (M = 20x, XN)	148
Plate-IV.12	Photomicrograph showing almandine garnet within staurolite (M = 20x, PPL)	148
Plate-IV.13	Photomicrograph showing ottrellite in staurolite schist (M = 20x, XN)	149
Plate-IV.14	Photomicrograph showing AF_4 crinkles of biotite (M = 20x, PPL)	149
Plate-IV.15	Photomicrograph showing garnet altering to chlorite (M = 20x, PPL)	153
Plate-IV.16	Photomicrograph showing sillimanite in quartzite (M = 20x, XN)	153